

**Supplementary Table 4.** Non-synonymous small variants or mutations (SNVs, MNVs and INDELs) found in LBs and leukocytes

patient	collection time (per patient)	cancer gene	AA change	MAF (%) <sup>1</sup>	saliva	plasma	leukocytes <sup>2</sup>
FA082	4	TP53	p.E180GfsTer67	0.1907		yes	
FA103	2	PDGFRA	p.A466VfsTer39	0.0955		yes	
FA122	1	TP53	p.R175H	7.2159	yes		
FA122	1	TP53	p.R175H	33.7177			yes
FA145	1	TP53	p.V157I	0.0768	yes		
FA145	1	TP53	p.W91L	0.0777	yes		
FA145	1	TP53	p.P92A	0.0932	yes		
FA145	3	TP53	p.V157I	0.0998	yes		
FA145	2	TP53	p.V157I	0.1148		yes	
FA145	4	PDGFRA	p.A466VfsTer39	0.2536			yes
FA228	1	TP53	p.P92A	0.0837	yes		
FA228	4	TP53	p.T102P	0.2742		yes	
FA452	2	TP53	p.W146SfsTer3	0.3911	yes		
FA452	2	TP53	p.W146SfsTer3	0.8229		yes	
FA467	1	TP53	p.Q38R	0.2323		yes	
FA531	2	TP53	p.G245S	0.0726	yes		
FA531	2	IDH2	p.R140Q	0.0758	yes		
FA531	3	TP53	p.G245S	0.2911	yes		
FA531	4	IDH2	p.R140Q	0.1121			yes
FA536	4	TP53	p.P92A	0.0928	yes		
FA536	4	TP53	p.W91L	0.0986	yes		

<sup>1</sup> MAF: molecular allele frequency

<sup>2</sup> Used to discard possible mutations in saliva/plasma as originated in white blood cells upon clonal hematopoiesis