- "Puberty age gap": New method of assessing pubertal timing and its association with
- 2 mental health problems
- 3 Niousha Dehestani<sup>1,2\*</sup>, Nandita Vijayakumar <sup>1</sup>, Gareth Ball<sup>3,4</sup>, Sina Mansour L.<sup>2,5</sup>, Sarah Whittle<sup>2,6</sup> &
- 4 Timothy J. Silk<sup>1,3,6\*</sup>

1

5

13

19

- 6 <sup>1</sup>School of Psychology, Deakin University, 221 Burwood Highway, Burwood, VIC 3125, Australia.
- <sup>2</sup>Melbourne Neuropsychiatry Centre, Department of Psychiatry, The University of Melbourne and
- 8 Melbourne Health, Victoria, Australia
- 9 <sup>3</sup>Developmental Imaging, Murdoch Children's Research Institute, Parkville 3052, Australia.
- <sup>4</sup>Department of Paediatrics, The University of Melbourne, Melbourne 3053, Australia.
- 11 <sup>5</sup>Department of Biomedical Engineering, The University of Melbourne, Victoria, Australia
- 12 <sup>6</sup>These authors jointly supervised this work: Sarah Whittle and Timothy J. Silk
- 14 Correspondence
- 15 Niousha Dehestani, School of Psychology, Deakin University, 221 Burwood Highway,
- Burwood, VIC 3125, Australia, ndehestanikolag@deakin.edu.au, +61422937273 or Tim Silk,
- 17 School of Psychology, Deakin University, 221 Burwood Highway, Burwood, VIC 3125,
- Australia, tim.silk@deakin.edu.au, +61408533160.

**Abstract** 

Puberty is linked to mental health problems during adolescence, and in particular, the timing of puberty is thought to be an important risk factor. This study developed a new measure of pubertal timing using multiple pubertal features and nonlinear associations with age and investigated its association with mental health problems. Using the Adolescent Brain Cognitive Development (ABCD) cohort, we implemented three models of pubertal timing by predicting chronological age from i) observed physical development, ii) hormones (testosterone and dehydroepiandrosterone [DHEA]), and iii) a combination of the two, using a supervised machine learning method (n ~9,900). Accuracy of the new models, and their associations with mental health problems were evaluated. The new pubertal timing measure performed better in capturing age variance compared to a commonly used method, and the physical measure accounted for more variance in mental health, such that earlier pubertal timing was associated with higher symptoms. This study demonstrates the utility for a new model of pubertal timing and suggests that physical maturation may play a predominant role in predicting mental health problems in early adolescence.

Introduction

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

Adolescence is the period between childhood and adulthood when individuals acquire the emotional and cognitive skills necessary to gain independence from their parents(1,2). Individuals also progress through puberty, the process of attaining reproductive maturity, during this period. This includes hormonal and physical changes such as body hair growth and gonadal maturation (along with the development of breasts and onset of menstruation in females)(3,4). The progression through puberty has been associated with an increase in susceptibility to a range of internalizing and externalizing mental health problems (5,6), which may reflect the effects of hormones on the central nervous system and/or psychosocial mechanisms related to physical differences from peers(7). Therefore, investigation of individual differences in pubertal processes that consider both hormonal and physical changes may lead to a better understanding of adolescent mental health problems. While all typically developing individuals progress through the same stages of puberty (based on observable physical changes), the onset and speed of progression can differ across individuals. As such, at any given period during adolescence, there is marked variability in pubertal stage, termed "pubertal timing" (8,9). Importantly, it is pubertal timing – not pubertal stage – that is often linked to the emergence and severity of mental health problems (10), though there are inconsistencies in the literature. A number of studies have found that in females, earlier pubertal timing is associated with internalizing problems (11,12), including depression(13,14), anxiety(15), eating disorders (16), and externalizing behaviors. On the other hand, a number of other studies report null findings regarding internalizing symptoms(14,15). Similarly in males, both early and late timing has been related to internalizing (11 VS 16) and externalizing problems (17 VS 18), while others have failed to

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

identify any associations (9, 15). Small sample sizes and methodological differences between studies may partially explain the inconsistent findings, with variation in the measures used to calculate pubertal timing likely important. Indeed, a meta-analysis of 101 studies found that the method of measuring puberty had a moderating role in the relationship between pubertal timing and mental health problems (4).

Different statistical approaches exist to measure the relative pubertal timing of individuals compared to same-aged peers. A common approach is to regress age from pubertal status (e.g., based on Pubertal Developmental Scale [PDS] scores) in a sample, with residuals reflecting earlier or later maturity relative to the group average (21). However, such approaches only capture individual differences in observable physical development and do not inform us about underlying biological mechanisms that are more reflected in hormone levels (22). Hormone levels provide valuable information regarding the endocrine processes of puberty (23,24). Previous studies have also demonstrated associations between changes in hormone levels and adolescent mental health problems. For example, increasing levels of dehydroepiandrosterone (DHEA) have been linked to internalizing symptoms (25), while testosterone has been associated with externalizing and disruptive behaviors (26). To the best of our knowledge, only one study has calculated pubertal timing with hormone data to predict internalizing behaviors in a sample of 174 females (27). Although this study did not find any significant associations, it is important to investigate this in a larger sample, of both females and males, and to assess associations across different dimensions of mental health problems. Additionally, no previous studies have combined both hormonal and physical measurements when calculating pubertal timing, which may be more sensitive to multiple underlying mechanisms that contribute to mental health problems. Thus, further investigation is needed

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

to develop pubertal timing methods that amalgamate different aspects of puberty in a single model.

In this study, we propose a multivariate method to calculate pubertal timing that

draws upon the "brain age" approach (28,29), where an association between multiple neuroimaging variables and chronological age is learned with supervised machine learning methods. Subtracting chronological age from brain age yields a "brain age gap" that reflects brain maturation relative to the group average (26,28). The major benefits of this model include being able to combine multiple features and reducing complex multivariate information to a single parameter. We propose to use a similar model to calculate estimates of pubertal timing by combining multiple puberty-related features. In contrast, alternative strategies either focus on single measures and only capture specific aspects or mechanisms of puberty or calculate the mean of multiple measures but may obscure the relative importance of each item. The second benefit of this method is that it can model nonlinear relationships between multiple measures of puberty and age, which is important given that nonlinear associations between specific features of puberty and chronological age have been observed in previous studies (30,31). In the current study, we aimed to create a normative model of pubertal timing that combines multiple measures of puberty capturing hormone levels and physical changes. By using a strict cross-validation approach, we ensure that model performance is robust. We compared model performance to a common linear model of pubertal timing (i.e., regressing age from total PDS score). We also compared the combined normative model against two unimodal models employing a similar supervised machine learning approach, one that uses hormonal measures alone and another that uses physical measures only. Finally, we examined the

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

association between each pubertal timing model and multiple dimensions of mental health problems. We hypothesized that the combined normative model of pubertal timing, which uses both hormonal and physical measures, would provide a more accurate prediction of age and better prediction of mental health problems compared to either of the unimodal (hormonal or physical) normative models or the (traditional) linear model of puberty. We additionally hypothesized that early pubertal timing would be associated with increased mental health problems in females and males. Based on a recent meta-analysis, we hypothesized significant effects of similar magnitude for internalizing and externalizing problems, while no significant effects were expected for attention problems (4). Prior findings of sex differences are mixed, and as such we did not propose specific hypotheses here. **Materials and Methods** To promote reproducible open research practices, all analyses conducted as part of this article are made publicly available in a git repository ("https://github.com/Niousha-Dehestani/Puberty-age"). In addition, more detail about participants, measurements and analysis is provided in the supplementary materials. **Participants** Participants were drawn from the ongoing, longitudinal, Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org/). Data was collected from ~11,500 children at baseline (47% females, age 9-10 years old) from 21 sites across the United States,

with annual data collection thereafter (see Supplementary Table 1 for demographic information). Data from baseline and 3 annual follow-up waves were used in the current analyses. We excluded participants who had a mismatch between their biological sex (collected per visit with the salivary sample) and their self-reported gender as well as those with missing values in biological sex (See Supplementary Information (SI) Appendix S1 for details of data cleaning procedures). The exact sample sizes utilized in each analysis are reported in detail below.

#### **Measures**

Pubertal Development Scale. The Pubertal Development Scale (PDS) measures observable physical signs of puberty. It includes items on height, body hair, and skin change in both sexes, as well as the onset of menarche and breast development for girls, and facial hair and voice changes for boys. Items are rated on a Likert scale from one to four points ("had not begun" to "already complete"). Onset of menarche was a binary variable (yes/no response) that was converted to one for "no" and four for "yes". The PDS can either be collected via self- or parent-report. However, each of these measures have their own limitations. The parent-report has good correspondence to clinician ratings, though this correspondence is lower in males (32). Conversely, the self-report is less accurate for individuals who are in the lower or upper pubertal stages as they tend to report toward the mid stages. Therefore, some studies have recommended using the parent-report PDS, especially in late childhood or early adolescence (33), and accordingly, the current study utilized this version. In this study we used individual PDS items rather than the average PDS score typically used in prior literature. This allowed us to capture the unique relationship between each item and our

outcomes of interest, consistent with previous literature that has found that separate PDS

items exhibit differential relationships with, e.g., brain structure (34).

"Puberty Age gap": A new method of pubertal timing

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

*Hormones.* DHEA and testosterone (TST) levels were measured via salivary hormone samples assayed by Salimetrics. Although estradiol was also measured, it was not used in the current analyses due to it only being available for females and having excessive missingness (n = 780). The hormone data was cleaned based on the protocol that published recently (35), which involved removing the confounding effects of collection time, duration of collection, wake-up time on collection day, having exercised before collection, and caffeine intake (with a linear mixed effect model, see SI for more details). Further information on the reliability and overall quality of the hormone data can be found in protocol (32). **Body Mass Index (BMI).** BMI was calculated as the average of two weight and height measurements per visit, assessed by the researcher. Next, BMI standard deviation scores (BMI z-scores) were calculated relative to age and sex, with reference to the CDC 2000 Growth Charts (36). Sociodemographic Variables. Five categories of race/ethnicity were coded: White, Black, Hispanic, Asian, and Other/Multi-race. Additionally, household income and education were obtained as measures of family socioeconomic status (SES) that were collected in a parent report. Mental health problems. The Child Behavior Checklist (CBCL) (age 6 to 18 form; (37)) was used to measure parent-reported mental health problems. CBCL includes 8 syndrome scales; 8

namely, Anxious/Depressed, Withdrawn/Depressed, Somatic Complaints, Social Problems, Thought Problems, Attention Problems, Aggressive Behavior, and Rule-breaking Behavior as well as three broad summary scales of externalizing, internalizing, and total problems.

#### **Statistical Analysis**

Calculating pubertal timing. Inheriting fundamental concepts from the literature on "brain age" (26), "puberty age" was computed using supervised machine learning. The model was trained to learn the relationship between physical and hormonal measurements of puberty (specifically, each PDS item, DHEA levels and TST levels) and chronological age, separately in males and females. Thereafter, the model was used to predict chronological age from pubertal measurements in an independent test sample. The effect of age was subsequently regressed from these predictions to adjust the bias created by regression toward the mean (RTM) (For more details see SI, Appendix S3, Figure S2 and Figure S3, and (38)). The bias-adjusted prediction of an individual's chronological age from pubertal measurements is termed "puberty age". Further, the residuals of the prediction model (after subtracting chronological age from puberty age) are referred to as the "puberty age gap", which we use as a dimensional measure indicative of relative pubertal timing. A positive puberty age gap is interpreted as a sign of earlier pubertal development compared to the normative age and sexmatched group, while a negative gap reflects relatively delayed pubertal development (see, Appendix 1, Figure S1).

We implemented a Generalized Additive Model (GAM) for the prediction of age (response variable) from pubertal measurements, which utilized TST, DHEA, and each of the PDS items as multivariate predictor variables. To remove the potential impacts of familial relations and repeat assessments of each individual, the sample was stratified to randomly

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

keep a single observation for each family (i.e., across waves and siblings). For details on data processing, such as the procedure for dealing with missing values, see SI Appendix S1 and Table S1. Additionally, to ensure the model was trained on a typically developing sample; using the CBCL DSM-oriented scales (consistent with DSM diagnostic categories), we excluded individuals above the threshold (symptom's score >60) for affective, anxiety, somatic, opposition defiant and conduct problems, as well as ADHD. The remaining participants who had symptom scores <60 were included in the training of models of puberty age (N = 4,949 (2,439 females)). However, a representation of the whole sample was used for testing (N = 9.919 (4.725 females)). As GAM fits smooth nonlinear curves in the form of spline functions, it is expected to outperform commonly utilized linear methods of pubertal timing measurement, given the nonlinear relations previously reported (30). Model training was performed within a nested 10-fold cross-validation design (for each fold in the outer loop, 90% of samples were used for model training and 10% for testing). Inner loop hyperparameter tuning was performed by a grid search for optimal regularization penalty on each term (i.e., using GAM) to minimize the estimated prediction error in the training sample (generalized cross-validation (GCV) score). To understand the contribution of different indices to the measurement of pubertal timing, we implemented three alternative models to estimate "puberty age". The first approach only used hormones (DHEA and TST) as features to predict chronological age, the second only used PDS items, and the third combined hormone and PDS items. The predictions of age were named *hormonal* puberty age, *physical* puberty age, and *combined* puberty age, respectively. In all three models, the residuals of age prediction indexed "puberty age gap", which reflects pubertal timing. We also used the partial dependence function in GAM models that can

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

performance.

reflect the importance of each feature in the combined puberty age model (for more detail, see SI, Appendix S4 and Figure S4). Comparison of "puberty age" models. We compared the performance of the three alternate puberty age models using Pearson's correlation between predicted and chronological age and median absolute error (MAE) averaged over the 10 cross-validation folds. Additionally, we used a non-parametric paired t-test (Wilcoxon signed-rank test) to statistically compare the model performance of different models based on the absolute error of predictions. "Physical puberty age" compared to a traditional pubertal timing model. The current study also compared the out-of-sample performance and accuracy of the physical puberty age model with the most common traditional pubertal timing model: linearly regressing age from average PDS score. In order to draw a comparison between the new model and the traditional approach, the performance of the physical puberty age model was contrasted with the traditional approach. The physical model was selected (rather than the hormonal or combined model) to ensure a fair comparison of two approaches that measure pubertal timing from the same input features (i.e., PDS). In order to conduct this comparison, a linear regression model was used to regress chronological age from the total PDS score (train) and used the fitted model coefficients in unseen data to measure timing (test). Similar to the puberty age model design, a 10-fold cross-validation design was used to measure traditional pubertal timing for the whole sample. This linear model provided an implementation of the traditional model in an out-of-sample prediction paradigm. Model performance was assessed based on the out-ofsample prediction accuracies (quantified by the absolute error of predictions). Similarly, we used the "Wilcoxon signed-rank test" to investigate the statistical differences in model

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

"Puberty age gap" associations with mental health problems. We used linear mixed-effect models (LMM) to investigate associations between each alternate "puberty age gap" measure and different dimensions of mental health problems, in males and females separately. The following formula was tested for each syndrome dimension (i.e., Anxious/Depressed, Withdrawn/Depressed, Somatic Complaints, Social Problems, Thought Problems, Attention Problems, Aggressive Behavior, and Rule-breaking Behavior), as well as three broad scales including total problems, externalizing, and internalizing problems. Sample size for this analysis was N = 9.919 (4,725 (females)) after removing the missing value in CBCL items. Mental health problems  $\sim puberty\ age\ gap + age + (1/site)$ . Age was included as a confound (fixed effect) and data collection sites were modelled as random effects. We corrected for multiple tests controlling for False Discovery Rate (FDR) at 5% and reported the FDR corrected p-values in the results. The different "pubertal age" LMMs were compared based on the Akaike Information Criterion (AIC) and a cut-off of 2 was used to indicate evidence for a better model, i.e., the model with an AIC that is at least 2 units smaller is considered a comparatively better model. Furthermore, to investigate whether associations between pubertal timing and mental health problems differed by age, we investigated the interaction effect of age and puberty age gap in predicting mental health problems using the following LMM: Mental health problems  $\sim puberty$  age gap + age + puberty age gap: age + (1/site)

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

Finally, given the known association of pubertal timing with BMI, SES, and race/ethnicity (Mendle et al., 2019), supplementary analyses repeated primary models while accounting for these variables as covariates (see SI, Appendix S5 and Table S2, for details of these analyses). This approach avoided potential complexity in our primary analyses due to the collinearity of confounding variables with our main variables of interest. Results Accuracy of puberty age models. All three models were able to provide significant and accurate out-of-sample predictions of age (Table 1, Figure 1). The performance of the hormonal model was better than the physical model in males, while on the contrary, physical puberty age was found to be a better predictor of age than hormonal puberty age in females. Across both sexes, combined puberty age explained the largest degree of variation in chronological age. Non-parametric statistical comparisons showed that the MAE between all models was significantly different (p<0.001). See the correlation between these three models in SI (See Appendix S2 and Figure S1). Approximate position of Table 1 Approximate position of Figure 1 "Puberty age model" vs. Traditional model. Comparing the out-of-sample prediction performance of the physical puberty age gap with the traditional model of pubertal timing showed that the physical puberty age model was significantly better at modelling the

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

relationship between puberty development and age in previously unseen data (See Table 2). Non-parametric statistical comparisons showed that the MAE between models were significantly different (p<0.001) Approximate position of Table 2 Pubertal timing measures predicting mental health problems. We investigated the association between each puberty age gap measure (hormonal, physical, and combined) and different dimensions of mental health problems (see Table 3, Figure 2). Hormonal puberty age gap was not significantly associated with any mental health problems in either males or females. Physical puberty age gap, however, was significantly positively associated with all dimensions of mental health problems in males and all dimensions except Anxiety-Depression in females. While similar associations were present for the combined puberty age gap, AIC differences indicated a better model fit for the physical puberty age gap in all cases. Finally, there were no significant interactions between age and any puberty age gap measure in predicting mental health problems in either females or males, suggesting that associations were stable across the age range. Approximate position of Table 3 Approximate position of Figure 2 **Discussion** In this study, we utilized biological and physical pubertal features and used supervised machine learning to model pubertal timing and its association with mental health

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

problems. Overall, a combined puberty age model predicted age more accurately than the unimodal (i.e., physical, or hormonal features alone). In males, the performance of the hormonal model was better than the physical model, which was not the case for females. This may be due to the fact that the two hormones tested (DHEA and testosterone) are more predominant in males, whilst female-dominant hormones (such as estradiol) were not considered. We showed that our novel pubertal timing model (based on physical pubertal features) had better accuracy than a conventional linear regression-based measurement in both sexes. Findings also highlight that our new method of assessing pubertal timing was significantly associated with most dimensions of mental health problems in males and females. Generally, we found that early pubertal maturation (based on physical and combined physical and hormonal features) was consistently predictive of various dimensions of mental health problems and could hence be an indicative risk factor for general mental health problems. Findings indicated that our novel normative puberty age models were able to provide significantly accurate out-of-sample predictions of age. This nonlinear modelling approach was found to be more accurate at predicting age than traditional measures of pubertal timing that rely on linear modelling of the relationship between (average) pubertal stage and age. These results highlight the presence of nonlinear relationships between age and pubertal development, consistent with previous literature reporting benefits of other nonlinear methods for calculating pubertal timing(21,31). Additionally, our examination of the collinearity between each of the PDS items showed that, although the items are significantly correlated, the correlations between them are less than 0.35 (for more details, see supplementary section 2). This suggests that each PDS item carried a significant amount of independent information. Our method thus used this independent information in a multivariate design instead of simplifying this information by averaging items to create a

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

single PDS score. Furthermore, our out-of-sample prediction provides an evaluation of the generalizability and replicability of study findings (39). We presented how a model that was trained on 90% of the sample could provide accurate out-of-sample predictions in the remaining 10% of the sample. This provides the opportunity for future research to measure pubertal timing in studies with smaller sample sizes, by taking advantage of this pre-trained normative model. This could potentially alleviate some study biases inherent to small sample sizes. Beyond its benefits for predicting age, our novel method of assessing pubertal timing was also significantly related to mental health problems in adolescents. Across most models of puberty age, we found that relatively early pubertal timing (i.e., positive puberty age gap) was associated with an increase in most dimensions of mental health problems. This is consistent with the maturation disparity hypothesis, whereby a mismatch between physical development and progression of emotional and cognitive development is purported to increase in those with early pubertal timing, which accounts for difficulties navigating the complexities and challenges of this period and may thus result in greater risk for mental health problems (7, 4). In particular, findings showed that physical puberty age gap outperformed combined and hormonal puberty age gap measures in the prediction of mental health problems in both females and males. Thus, although the prediction accuracy of the combined puberty age model was significantly better than physical puberty age, it was not better at explaining mental health problems. Moreover, we did not find any significant associations between hormonal puberty age gap and mental health problems in males or females. These findings are partially consistent with (40), who reported that pubertal timing measured from hormonal information (testosterone and DHEA) did not predict internalizing symptoms. Consistent with prior studies, these findings may suggest that psychosocial

mechanisms have a larger role (in contrast to biological mechanisms indexed by hormones)

"Puberty Age gap": A new method of pubertal timing

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

in predicting mental health problems in early adolescence (14). Likewise, prior work has suggested that earlier physical development impacts social functions such as difficulty maintaining friendship with peers who mature later, and a tendency to associate with older adolescents that engage in more externalizing behaviors (12). While this study has strengths in its large sample size, use of hormonal assays and physical measurement of puberty, as well as implementation of a novel, generalizable method for assessing pubertal timing, there are limitations that should be addressed in future work. First, the age range of the sample used was rather limited and a wider age range could have captured more variance in pubertal maturation and improved the out-of-sample model prediction of age. Future studies that implement our model in a wider age range encompassing the start to completion of puberty would be able to better capture the complete nonlinear relationship between age and pubertal development. Relatedly, while we did not observe any interactive effects of age with pubertal timing in predicting mental health problems symptoms in early adolescents, it is possible that such effects may be detectable across a larger age range. Further, different measures of puberty age gap may also differentially predict mental health problems in earlier versus later adolescence. Second, this study focused on a limited number of pubertal hormones, and the omission of other hormonal features may have impacted the predictive power of hormonal puberty age gap – particularly in females. Additionally, variations in the time for collecting hormones across the day could have also had an impact on the inferior performance of the hormonal age gap (although these were statistically accounted for). Thus, future studies could incorporate more detailed assessments of hormones to improve model prediction. Although the current study used the parent report version of PDS due to reportedly higher reliability in this particular age range, it would be valuable for future research to replicate this analysis in a sample with a wider age

range with self- or clinician-assessed pubertal development.

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

In conclusion, the current study proposes a smooth nonlinear puberty age model that facilitates generalizable investigations of pubertal timing in future studies. Our findings also highlight the importance of physical pubertal maturation, relative to hormonal changes, for mental health problems during early adolescence. This suggests that psychosocial mechanisms may play an important role in the relationship between early pubertal timing and mental health problems, which has implications for interventions aimed at reducing the risk of the emergence of mental health problems in adolescence. **Data Sharing:** This study used data from Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a longitudinal study which collected data from ~ 11000 children in age 9-10 and follow them through 10 years. A full list of supporters is available at <a href="https://abcdstudy.org/federal-partners.html">https://abcdstudy.org/federal-partners.html</a>. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/scientists/workgroups/. ABCD consortium investigators designed and implemented the study and/or provided data but did not participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. Acknowledgement

Niousha Dehestani was supported by a Deakin University Postgraduate

425

427

431

432

Research Scholarships (DUPRS). Additionally, this study was supported by the MASSIVE 426 high-performance computing facility (www.massive.org.au). 428 **Conflict of Interest** 429 The authors declare that they have no financial interests or personal relationships that could 430 impact the work reported in this paper.

457

433 434 References 435 1. Blakemore S-J, Choudhury S. Development of the adolescent brain: implications for 436 executive function and social cognition. J Child Psychol Psychiatry [Internet]. 2006 437 Mar;47(3–4):296–312. Available from: 438 https://onlinelibrary.wiley.com/doi/10.1111/j.1469-7610.2006.01611.x 439 2. Sawyer SM, Azzopardi PS, Wickremarathne D, Patton GC. The age of adolescence. 440 Lancet Child Adolesc Heal. 2018;2(3):223–8. 441 3. Banerjee I, Clayton P. The Genetic Basis for the Timing of Human Puberty. J 442 Neuroendocrinol [Internet]. 2007 Nov;19(11):831–8. Available from: 443 https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2826.2007.01598.x 444 4. Kohen-Raz R. PHYSIOLOGICAL MATURATION AND MENTAL GROWTH AT 445 PRE-ADOLESCENCE AND PUBERTY. J Child Psychol Psychiatry [Internet]. 1974 446 Jul;15(3):199–213. Available from: 447 https://onlinelibrary.wiley.com/doi/10.1111/j.1469-7610.1974.tb01245.x 448 5. Ullsperger JM, Nikolas MA. A meta-analytic review of the association between 449 pubertal timing and psychopathology in adolescence: Are there sex differences in risk? 450 Psychol Bull [Internet]. 2017 Sep;143(9):903–38. Available from: 451 http://doi.apa.org/getdoi.cfm?doi=10.1037/bul0000106 452 6. Natsuaki MN, Klimes-Dougan B, Ge X, Shirtcliff EA, Hastings PD, Zahn-Waxler C. 453 Early pubertal maturation and internalizing problems in adolescence: Sex differences 454 in the role of cortisol reactivity to interpersonal stress. J Clin Child Adolesc Psychol. 455 2009;38(4):513–24. 456 7. Ge X, Natsuaki MN. In search of explanations for early pubertal timing effects on

developmental psychopathology. Curr Dir Psychol Sci. 2009;18(6):327–31.

A W, TANNER JM. VARIATIONS IN PATTERN OF PUBERTAL CHANGES IN 458 8. 459 GIRLS. Obstet Gynecol Surv [Internet]. 1970 Jul;25(7):694–5. Available from: 460 http://journals.lww.com/00006254-197007000-00018 461 9. Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal 462 status: Reliability, validity, and initial norms. J Youth Adolesc [Internet]. 1988 463 Apr;17(2):117–33. Available from: http://link.springer.com/10.1007/BF01537962 464 10. Graber JA. Pubertal timing and the development of psychopathology in adolescence 465 and beyond. Horm Behav [Internet]. 2013 Jul;64(2):262–9. Available from: 466 http://dx.doi.org/10.1016/j.yhbeh.2013.04.003 467 11. Crockett LJ, Carlo G, Wolff JM, Hope MO. The role of pubertal timing and 468 temperamental vulnerability in adolescents' internalizing symptoms. Dev 469 Psychopathol [Internet]. 2013 May 30;25(2):377–89. Available from: 470 https://www.cambridge.org/core/product/identifier/S0954579412001125/type/journal\_ 471 article 472 12. Marceau K, Neiderhiser JM, Lichtenstein P, Reiss D. Genetic and Environmental 473 Influences on the Association Between Pubertal Maturation and Internalizing 474 Symptoms. J Youth Adolesc. 2012;41(9):1111–26. 475 13. Hamlat EJ, McCormick KC, Young JF, Hankin BL. Early pubertal timing predicts 476 onset and recurrence of depressive episodes in boys and girls. J Child Psychol 477 Psychiatry Allied Discip. 2020;61(11):1266–74. 478 14. Mendle J, Ferrero J. Detrimental psychological outcomes associated with pubertal 479 timing in adolescent boys. Dev Rev [Internet]. 2012;32(1):49–66. Available from: 480 http://dx.doi.org/10.1016/j.dr.2011.11.001 481 15. Weingarden H, Renshaw KD. Early and late perceived pubertal timing as risk factors 482 for anxiety disorders in adult women. J Psychiatr Res [Internet]. 2012;46(11):1524–9.

483		Available from: http://dx.doi.org/10.1016/j.jpsychires.2012.07.015
484	16.	Klump KL. Puberty as a critical risk period for eating disorders: A review of human
485		and animal studies. Horm Behav [Internet]. 2013;64(2):399-410. Available from:
486		http://dx.doi.org/10.1016/j.yhbeh.2013.02.019
487	17.	Winer JP, Parent J, Forehand R, Breslend NL. Interactive Effects of Psychosocial
488		Stress and Early Pubertal Timing on Youth Depression and Anxiety: Contextual
489		Amplification in Family and Peer Environments. J Child Fam Stud [Internet]. 2016
490		May 7;25(5):1375–84. Available from: http://link.springer.com/10.1007/s10826-015-
491		0318-0
492	18.	R. K-H, M. M, P. R, M. R. Early puberty is associated with mental health problems in
493		middle adolescence. Soc Sci Med [Internet]. 2003;57(6):1055-64. Available from:
494		http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L368
495		76071%5Cnhttp://dx.doi.org/10.1016/S0277-9536(02)00480-
496		X%5Cnhttp://sfxhosted.exlibrisgroup.com/galway?sid=EMBASE&issn=02779536&id
497		=doi:10.1016%2FS0277-9536%2802%2900480-X&atitle=E
498	19.	Dorn LD, Dahl RE, Williamson DE, Birmaher B, Axelson D, Perel J, et al.
499		Developmental Markers in Adolescence: Implications for Studies of Pubertal
500		Processes. J Youth Adolesc. 2003;32(5):315–24.
501	20.	Ge X, Brody GH, Conger RD, Simons RL, Murry VMB. Contextual amplification of
502		pubertal transition effects on deviant peer affiliation and externalizing behavior among
503		African American children. Dev Psychol. 2002;38(1):42–54.
504	21.	Mendle J, Beltz AM, Carter R, Dorn LD. Understanding Puberty and Its Measurement:
505		Ideas for Research in a New Generation. J Res Adolesc. 2019;29(1):82–95.
506	22.	Dorn LD, Biro FM. Puberty and its measurement: A decade in review. J Res Adolesc.
507		2011;21(1):180–95.

508	23.	Byrne ML, Whittle S, Vijayakumar N, Dennison M, Simmons JG, Allen NB. A
509		systematic review of adrenarche as a sensitive period in neurobiological development
510		and mental health. Dev Cogn Neurosci [Internet]. 2017 Jun;25:12–28. Available from:
511		http://dx.doi.org/10.1016/j.dcn.2016.12.004
512	24.	Shirtcliff EA, Zahn-Waxler C, Klimes-Dougan B, Slattery M. Salivary
513		dehydroepiandrosterone responsiveness to social challenge in adolescents with
514		internalizing problems. J Child Psychol Psychiatry Allied Discip. 2007;48(6):580–91.
515	25.	Peixoto C, José Grande A, Gomes Carrilho C, Egidio Nardi A, Cardoso A, Barciela
516		Veras A. Dehydroepiandrosterone for depressive symptoms: A systematic review and
517		meta-analysis of randomized controlled trials. J Neurosci Res. 2020;98(12):2510–28.
518	26.	Han G, Miller JG, Cole PM, Zahn-Waxler C, Hastings PD. Adolescents' internalizing
519		and externalizing problems predict their affect-specific HPA and HPG axes reactivity.
520		Dev Psychobiol. 2015;57(6):769–85.
521	27.	Barendse MEA, Cosme D, Flournoy JC, Vijayakumar N, Cheng TW, Allen NB, et al.
522		Neural correlates of self-evaluation in relation to age and pubertal development in
523		early adolescent girls. Dev Cogn Neurosci [Internet]. 2020 Aug;44:100799. Available
524		from: https://doi.org/10.1016/j.dcn.2020.100799
525	28.	Cole JH, Franke K. Predicting Age Using Neuroimaging: Innovative Brain Ageing
526		Biomarkers. Trends Neurosci [Internet]. 2017 Dec;40(12):681–90. Available from:
527		http://dx.doi.org/10.1016/j.tins.2017.10.001
528	29.	Kaufmann T, van der Meer D, Doan NT, Schwarz E, Lund MJ, Agartz I, et al.
529		Common brain disorders are associated with heritable patterns of apparent aging of the
530		brain. Nat Neurosci [Internet]. 2019 Oct 24;22(10):1617–23. Available from:
531		http://www.nature.com/articles/s41593-019-0471-7
532	30.	Gracia-Tabuenca Z, Moreno MB, Barrios FA, Alcauter S. Development of the brain

533		functional connectome follows puberty-dependent nonlinear trajectories. Neuroimage
534		[Internet]. 2021;229(January):117769. Available from:
535		https://doi.org/10.1016/j.neuroimage.2021.117769
536	31.	Marceau K, Ram N, Houts RM, Grimm KJ, Susman EJ. Individual differences in
537		boys' and girls' timing and tempo of puberty: Modeling development with nonlinear
538		growth models. Dev Psychol [Internet]. 2011 Sep;47(5):1389–409. Available from:
539		http://doi.apa.org/getdoi.cfm?doi=10.1037/a0023838
540	32.	Rasmussen AR, Wohlfahrt-Veje C, De Renzy-Martin KT, Hagen CP, Tinggaard J,
541		Mouritsen A, et al. Validity of self-assessment of pubertal maturation. Pediatrics.
542		2015;135(1):86–93.
543	33.	Huang B, Hillman J, Biro FM, Ding L, Dorn LD, Susman EJ. Correspondence
544		Between Gonadal Steroid Hormone Concentrations and Secondary Sexual
545		Characteristics Assessed by Clinicians, Adolescents, and Parents. J Res Adolesc.
546		2012;22(2):381–91.
547	34.	Hu S, Pruessner JC, Coupé P, Collins DL. Volumetric analysis of medial temporal
548		lobe structures in brain development from childhood to adolescence. Neuroimage
549		[Internet]. 2013;74:276–87. Available from:
550		http://dx.doi.org/10.1016/j.neuroimage.2013.02.032
551	35.	Herting MM, Uban KA, Gonzalez MR, Baker FC, Kan EC, Thompson WK, et al.
552		Correspondence Between Perceived Pubertal Development and Hormone Levels in 9-
553		10 Year-Olds From the Adolescent Brain Cognitive Development Study. Front
554		Endocrinol (Lausanne) [Internet]. 2021 Feb 18;11(April 2020):1–22. Available from:
555		https://www.frontiersin.org/articles/10.3389/fendo.2020.549928/full
556	36.	Gray JC, Schvey NA, Tanofsky-Kraff M. Demographic, psychological, behavioral,
557		and cognitive correlates of BMI in youth: Findings from the Adolescent Brain

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

Cognitive Development (ABCD) study. Psychol Med. 2020;50(9):1539–47. 37. Achenbach TM, Ruffle TM. The child behavior checklist and related forms for assessing behavioral/emotional problems and competencies. Pediatr Rev. 2000;21(8):265-71. 38. Liang H, Zhang F, Niu X. Investigating systematic bias in brain age estimation with application to post-traumatic stress disorders. Hum Brain Mapp. 2019;40(11):3143–52. 39. Rosenberg MD, Casey BJ, Holmes AJ. Prediction complements explanation in understanding the developing brain. Nat Commun [Internet]. 2018;9(1):1–13. Available from: http://dx.doi.org/10.1038/s41467-018-02887-9 40. Barendse MEA, Byrne ML, Flournoy JC, Mcneilly EA, Williamson VG, Barrett MY, et al. Journal of Abnormal Psychology Internalizing Psychopathology in Early Adolescent Girls Multimethod Assessment of Pubertal Timing and Associations With Internalizing Psychopathology in Early Adolescent Girls. 2021;

# **Tables and Figures:**

Table 1. Comparison of the out-of-sample prediction performance of alternative puberty age gap

## 577 models

574

575

576

578

579

	Fer	nale	Male			
	r	mae	r	mae		
Combined puberty age	0.58	0.65 year	0.54	0.67 year		
Hormonal puberty age	0.37	0.76 year	0.45	0.72 year		
Physical development puberty age	0.55	0.67 year	0.43	0.72 year		

r = correlation, mae = mean absolute error

**Figure 1.** Performance of puberty age gap models. For all three models (combined, physical development and hormonal), the prediction of the age of each individual is plotted against chronological age (in years). Red dots indicate individuals with a positive puberty age gap (early timing) and blue dots indicate individuals with a negative puberty age gap (late timing).

587

588

589

590

591

# Table 2. Comparison of the out-of-sample prediction performance of the traditional pubertal timing

## model and the physical development puberty age model

	Fen	nale	Male		
	r	mae	r	тае	
Traditional pubertal timing	0.49	0.71	0.32	0.78	
Physical development Puberty age Model	0.55	0.68	0.43	0.72	

r = correlation, mae = mean absolute error

**Table 3.** Associations between puberty age gap and mental health problems measured by linear mixed effect regression

"Puberty Age gap": A new method of pubertal timing

	Female							Male					
	Physical		Combined		Hormonal		Physical		Combined		Hormonal		
	t	aic											
Total	7.16***	31382.19	5.06***	31407.62	-1.61	31430.59	6.19***	34308.27	6.03***	34330.51	-0.44	34346.23	
Internalising	5.81***	31145.29	3.93**	31163.47	-1.52	31176.60	6.33***	33654.96	5.29***	33677.32	-0.41	33694.64	
Externalising	6.80***	30528.71	5.04***	30549.40	-1.150	30573.33	6.54***	33492.17	6.07***	33515.11	0.23	33534.59	
Anxiety_Depression	1.82	26535.61	1.03	26537.87	-1.20	26537.47	5.27***	29000.46	4.78**	29019.26	-1.20	29026.61	
Withdrawn_depression	6.89***	25451.83	5.67***	25467.03	0.84	25498.29	5.07***	29052.81	3.98**	29063.78	0.89	29077.61	
Somatic_Complaints	4.73**	26644.49	2.68*	26659.56	-2.38	26661.07	6.35***	28592.79	5.17**	28619.53	-0.90	28632.09	
Attention_problems	3.73**	26187.00	3.26*	26190.28	0.05	26200.82	2.86*	29291.46	1.89*	29296.84	-0.39	29299.45	
Rule_breaking	5.38**	24226.11	3.93*	24239.47	-0.72	24254.27	6.29***	26982.27	5.31**	27005.78	-0.08	27021.57	
Aggressive_Behaviour	4.82**	24915.41	3.64*	24925.27	-0.51	24938.20	5.78***	28625.15	4.46**	28646.70	-0.28	28658.31	
Thought_Problems	3.10*	25920.98	2.25*	25925.51	-1.36	25928.69	5.02***	29131.42	3.61**	29147.13	-0.49	29156.23	
Social_Problems	4.96**	23988.88	3.67*	24000.00	-1.24	24011.85	5.50***	26908.05	3.88**	26923.08	0.51	26937.86	

Note: Statistical significance is indicated by asterisks (\*: p<0.05; \*\*: p<0.01; \*\*\*: p<0.001; FDR corrected). The bold numbers reflect better model

597 performance.

598

**Figure 2.** Bar plot showing the T-statistic from the prediction of each dimension of mental health problems from different puberty age models. Hormonal puberty age gap is not significantly associated with any dimension of mental health problems.



