

**Supplementary Table 2: Enriched Biocarta pathway gene sets in Psoriasis patients vs. Healthy Controls as determined by Gene Set Enrichment Analysis.**

Pathways that are also enriched in Darier Disease patients are highlighted in bold. FDR, false discovery rate

Biocarta pathway name	No of genes in Nanostring panel	normalized enrichment score	FDR q-value
<b>IL-17 signaling</b>	12	2.32	0.174
<b>Cytokine Network</b>	17	2.27	0.174
<b>Regulation of hematopoiesis by cytokines</b>	12	2.11	0.174
<b>NFkB activation by Nontypeable Hemophilus influenzae</b>	14	2.06	0.174
<b>Cytokines and inflammatory response</b>	21	1.82	0.174
<b>Free Radical Induced Apoptosis</b>	4	1.82	0.174
IL22 Soluble Receptor Signaling	9	1.76	0.174
<b>Adhesion and Diapedesis of Granulocytes</b>	13	1.74	0.174
Mechanism of Gene Regulation by Peroxisome Proliferators via PPARa	8	1.72	0.174
<b>Local accute inflammatory response</b>	15	1.6	0.224
<b>Adhesion and Diapedesis of Lymphocytes</b>	12	1.59	0.22
Antigen Processing and Presentation	11	1.52	0.231