## **Supplementary Figures**

## A systematic review on machine learning approaches in the diagnosis of rare genetic diseases

Roman-Naranjo P 1,2,3, Parra-Perez AM 1,2,3,4, Lopez-Escamez JA 1,2,3,4

<sup>1</sup>Otology and Neurotology Group CTS495, Department of Genomic Medicine, GENYO - Centre for Genomics and Oncological Research - Pfizer, University of Granada, Junta de Andalucía, PTS, Granada, Spain.

<sup>2</sup>Division of Otolaryngology, Department of Surgery, Instituto de Investigación Biosanitaria, ibs.GRANADA, Granada, Universidad de Granada, Granada, Spain.

<sup>3</sup>Sensorineural Pathology Programme, Centro de Investigación Biomédica en Red en Enfermedades Raras, CIBERER, Madrid, Spain


<sup>4</sup>Meniere's Disease Neuroscience Research Program, Faculty of Medicine & Health, School of Medical Sciences, The Kolling Institute, University of Sydney, Sydney, New South Wales, Australia

## Supplementary Figure 1: Category and journal where selected articles were published.




JIF: Journal Impact Factor

**Supplementary Figure 2**: A) Authors from selected articles showed a low level of collaboration, with only 3 authors participating in 2 or more articles (Rachel Karchin, Dimitrios Vitsios and Slavé Petrovski). B) Co-occurrence analysis on abstracts of selected articles.



Supplementary Figure 3: Sample sizes of studies using A) their own cohorts and B) external cohorts.



**Supplementary Figure 4:** Studies using external cohorts showed higher variability in their methods, using different sequencing platforms, read lengths and variant callers in the same study.

