

Executability and repeatability of a study setup using wearable and computerized technology to examine a wide range of physiological and cognitive functions of a patient outside hospital

Kiti Müller et al.

SUPPLEMENT 3: Clinical neuropsychological tests and subjective questionnaires

Table 1. Clinical neuropsychological test battery

Cognitive domain	Neuropsychological test
Processing speed	WAIS-IV Coding [1] Stroop test, colour-congruent [2] FAT, perceptual motor speed [3,4]
Executive functions	Stroop test, colour-incongruent [2] Dual task [5] Verbal fluency test, phonemic [2] FAT, set shifting [3,4]
Working memory	WMS-III Letter-Number Sequencing [6] WMS-III Digit Span [6] FAT, visuospatial span [4]
Memory and learning	WMS-III Word Lists [6] WMS-III Logical Memory [6] WMS-III Visual Reproduction [6]
Visuospatial perception	WAIS-IV Block Design [1] Visual search for parallel lines [7]
Verbal functions	WAIS-IV Similarities [1] Boston naming test [8] Verbal fluency test, semantic [2]
Questionnaires	Checklist for cognitive and emotional consequences following stroke (CLCE-24) [9] Cognitive Reserve Scale (CRS) [10] Resilience Scale (RS-14) [11]

Table 2. Subjective Questionnaires

Patients answered the following validated subjective questionnaires via an internet portal.

Questionnaires and references	Short description
Oslo Social Support Scale [12]	Three-item scale used to determine the social support of individual's family, friends, and neighbours
Alcohol Use Disorders Identification Test (AUDIT) [13]	Widely used 10-item screening tool developed to evaluate individual's alcohol-related problems
WHO Alcohol, Smoking and Substance Involvement Screening Test (ASSIST) [13]	A brief screening tool used to discover the use of psychoactive substances
International Physical Activity Questionnaire (IPAQ) [14]	A brief questionnaire for assessing individual's health-related physical activity comprehensively and multidimensionally
Mediterranean Diet Score (MDS) [15]	Nine-item score to evaluate the healthiness of individual's diet habits
Shirom-Melamed Burnout Measure (SMBM) [16]	Fourteen-item tool to evaluate individual's burnout level
Absenteeism and Presenteeism questions of the World Health Organization's Health and Work Performance Questionnaire (HPQ) [17]	Seven-item questionnaire extracted from WHO's HPQ. Describes individual's work performance and the number of sick leaves and other work absences
Perceived Stress Scale [18]	Ten-item scale to describe stress experiences
Mobile Modified Rankin Scale [19]	Mobile-based scale used to evaluate performance and independency mostly in stroke patients
Beck Depression Inventory (BDI) [20]	Widely used questionnaire to evaluate the level of depression symptoms
Beck Anxiety Inventory (BAI) [21]	Widely used questionnaire to evaluate the level of anxiety symptoms
Epworth Sleepiness Scale (ESS) [22]	Ten-item scale describing day-time sleepiness
Basic Nordic Sleep Questionnaire (BNSQ) [23]	A questionnaire determining a wide range of sleep complaints
Fatigue Assessment Scale (FAS) [24]	Ten-item scale evaluating the levels of cognitive performance and fatigue
EuroHis-8 [25]	Eight-item questionnaire to evaluate an individual's quality of life
Checklist for Cognitive and Emotional consequences (CLCE-24) [9]	A screening tool to identify cognitive and emotional problems
Brain Work Questionnaire (BWQ) [26]	A comprehensive questionnaire evaluating cognitive demands at work and work-related cognitive strain

Table 3. Evaluation form filled after home recordings. Estimates are median values (0-10) with interquartile range (IQR). The items of “Number of tests” and “Number of questionnaires” are graded from 0 = too many to 10 = too few and 5 represents not too many or too few. Otherwise, the higher the value the easier or more positive experience. (N= number of patients answering.)

Evaluation about the study protocol	Preoperatively (N=8)	1st postoperatively (N=11)	2nd postoperatively (N=11)
Mental demand	7.5 (7-10)	10 (4-10)	6 (4-8)
Physical demand	8.5 (7-10)	9 (6-10)	6 (5-9)
Effort	7 (4-8.5)	9 (4-10)	6 (5-8)
Frustration	8.5 (7-9.5)	10 (4-10)	5 (4-10)
Saliva samples	10 (9-10)	10 (10-10)	10 (5-10)
Sleep diary	10 (8.5)	10 (10-10)	10 (8-10)
Cognitive tablet tests	9 (5-10)	10 (8-10)	9.5 (9-10)
Physiological measurements	9.5 (9-10)	10 (10-10)	10 (8-10)
Instructions			
• Sufficiency	10 (9.5-10)	10 (9-10)	10 (10-10)
• Clearness	10 (10-10)	10 (9-10)	10 (9-10)
Opportunity to ask help	10 (8-10)	10 (10-10)	10 (10-10)
Number of tests	5 (4-5)	5 (5-5)	5 (3-5)
Number of questionnaires	4.5 (3.5-5)	5 (4-5)	5 (2-5)
Re-participation	10 (8-10)	10 (6-10)	10 (5-10)

References

- 1 Wechsler D. *WAIS-IV: Wechsler adult intelligence scale*. San Antonio, Tex.: : Pearson 2008.
- 2 Lezak MD, Howieson, Diane B., Bigler Erin D., *et al.*, editors. *Neuropsychological assessment*. 5th ed. Oxford ; New York: : Oxford University Press 2012.
- 3 Reitan RM. Validity of the Trail Making Test as an Indicator of Organic Brain Damage. *Perceptual and Motor Skills* 1958;**8**:271–6. doi:10.2466/pms.1958.8.3.271
- 4 Kessels RP, van Zandvoort MJ, Postma A, *et al.* The Corsi Block-Tapping Task: standardization and normative data. *Appl Neuropsychol* 2000;**7**:252–8. doi:10.1207/S15324826AN0704_8
- 5 Vilkki J, Virtanen S, Surma-Aho O, *et al.* Dual task performance after focal cerebral lesions and closed head injuries. *Neuropsychologia* 1996;**34**:1051–6. doi:10.1016/0028-3932(96)00028-0

- 6 Wechsler D. Wechsler Adult Intelligence Scale--Third Edition. 2019. doi:10.1037/t49755-000
- 7 Vilkki J. Hemi-inattention in visual search for parallel lines after focal cerebral lesions. *J Clin Exp Neuropsychol* 1989;**11**:319–31. doi:10.1080/01688638908400892
- 8 Laine M, Goodglass H, Niemi J, *et al.* Adaptation of the Boston Diagnostic Aphasia Examination and the Boston Naming Test into Finnish. *Scandinavian Journal of Logopedics and Phoniatics* 1993;**18**:83–92. doi:10.3109/14015439309101353
- 9 van Heugten C, Rasquin S, Winkens I, *et al.* Checklist for cognitive and emotional consequences following stroke (CLCE-24): development, usability and quality of the self-report version. *Clin Neurol Neurosurg* 2007;**109**:257–62. doi:10.1016/j.clineuro.2006.10.002
- 10 León I, García-García J, Roldán-Tapia L. Estimating cognitive reserve in healthy adults using the Cognitive Reserve Scale. *PLoS One* 2014;**9**:e102632. doi:10.1371/journal.pone.0102632
- 11 Losoi H, Turunen S, Wäljas M, *et al.* Psychometric Properties of the Finnish Version of the Resilience Scale and its Short Version. *PCH* 2013;**2**:1–10. doi:10.5964/pch.v2i1.40
- 12 Kocalevent R-D, Berg L, Beutel ME, *et al.* Social support in the general population: standardization of the Oslo social support scale (OSSS-3). *BMC Psychol* 2018;**6**:31. doi:10.1186/s40359-018-0249-9
- 13 Saunders JB, Aasland OG, Babor TF, *et al.* Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-II. *Addiction* 1993;**88**:791–804. doi:10.1111/j.1360-0443.1993.tb02093.x
- 14 Booth M. Assessment of Physical Activity: An International Perspective. *Research Quarterly for Exercise and Sport* 2000;**71**:114–20. doi:10.1080/02701367.2000.11082794
- 15 Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, *et al.* Diet and overall survival in elderly people. *BMJ* 1995;**311**:1457–60. doi:10.1136/bmj.311.7018.1457
- 16 Gerber M, Colledge F, Mücke M, *et al.* Psychometric properties of the Shirom-Melamed Burnout Measure (SMBM) among adolescents: results from three cross-sectional studies. *BMC Psychiatry* 2018;**18**:266. doi:10.1186/s12888-018-1841-5
- 17 Kessler RC, Barber C, Beck A, *et al.* The World Health Organization Health and Work Performance Questionnaire (HPQ): *Journal of Occupational and Environmental Medicine* 2003;**45**:156–74. doi:10.1097/01.jom.0000052967.43131.51
- 18 Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;**24**:385–96.
- 19 van Swieten JC, Koudstaal PJ, Visser MC, *et al.* Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 1988;**19**:604–7. doi:10.1161/01.str.19.5.604
- 20 Beck AT, Ward CH, Mendelson M, *et al.* An inventory for measuring depression. *Arch Gen Psychiatry* 1961;**4**:561–71. doi:10.1001/archpsyc.1961.01710120031004

- 21 Beck AT, Epstein N, Brown G, *et al.* An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology* 1988;**56**:893–7. doi:10.1037/0022-006X.56.6.893
- 22 Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;**14**:540–5. doi:10.1093/sleep/14.6.540
- 23 Partinen null, Gislason null. Basic Nordic Sleep Questionnaire (BNSQ): a quantitated measure of subjective sleep complaints. *J Sleep Res* 1995;**4**:150–5. doi:10.1111/j.1365-2869.1995.tb00205.x
- 24 Michielsen HJ, De Vries J, Van Heck GL. Psychometric qualities of a brief self-rated fatigue measure: The Fatigue Assessment Scale. *J Psychosom Res* 2003;**54**:345–52. doi:10.1016/s0022-3999(02)00392-6
- 25 da Rocha NS, Power MJ, Bushnell DM, *et al.* The EUROHIS-QOL 8-item index: comparative psychometric properties to its parent WHOQOL-BREF. *Value Health* 2012;**15**:449–57. doi:10.1016/j.jval.2011.11.035
- 26 Kalakoski V, Selinheimo S, Valtonen T, *et al.* Effects of a cognitive ergonomics workplace intervention (CogErg) on cognitive strain and well-being: a cluster-randomized controlled trial. A study protocol. *BMC Psychol* 2020;**8**:1. doi:10.1186/s40359-019-0349-1