
## Supplementary Material

## SUMMIT-FA: A new resource for improved transcriptome

imputation using functional annotations

Phenotypes Analyzed by SUMMIT-FA		
Asthma	Hypertension	Schizophrenia
Coronary Artery Disease	${ m Hyperparathyroidism}$	Systemic Lupus Erythematosus
Diastolic Pressure	Irritable Bowel Disease	Systolic Pressure
Deep Vein Thrombosis	Insomnia	Type-I Diabetes
Eosinophil Count	Lymphocyte Count	Type-II Diabetes
Eczema	Monocyte Count	Triglycerides
Gout	Psoriasis	Ulcerative Colitis
High Cholesterol	Rheumatoid Arthritis	Waist Circumference

Table 1: 24 phenotypes used in TWAS analysis.



Method 🚔 SUMMIT-FA 📫 SUMMIT 📫 Lassosum 🚔 TWAS-FUSION 📫 PrediXcan




Figure 1: Simulation performance comparison based on gene *CHURC1*. Plots of imputation  $R^2$  in test samples by SUMMIT-FA, SUMMIT, Lassosum, TWAS-FUSION, and PrediXcan for varied proportion of causal SNPs  $p_{causal}$ , DNAm heritability  $h_e^2$ , and phenotypic heritability  $h_p^2$ .

0.05

Pcausal Method  $\Rightarrow$  SUMMIT-FA  $\Rightarrow$  SUMMIT  $\Rightarrow$  Lassosum  $\Rightarrow$  TWAS-FUSION  $\Rightarrow$  PrediXcan (c)  $h_p^2 = 0.8$ 

0.1

0.2

0.00

0.01

0.05

0.1 Pcausal 0.2

0.000

0.01

0.0000

0.01

0.05

p<sub>causal</sub>

0.1

0.2




Figure 2: Simulation performance comparison based on gene *CHURC1*. Plots of statistical power for SUMMIT-FA, SUMMIT, Lassosum, TWAS-FUSION, and PrediXcan for varied proportion of causal SNPs  $p_{causal}$ , DNAm heritability  $h_e^2$ , and phenotypic heritability  $h_p^2$ . Empirical power was determined by the proportion of p-values  $< 2.5 \times 10^{-6}$ .




Figure 3: QQ plots of p-values under H<sub>0</sub> based on the *CHURC1* gene. 20,000 replicates were run under the null hypothesis to examine type I error rates, with  $h_p^2 = 0.2$ ,  $h_e^2 = 0.005$ ,  $p_{\text{causal}} = 0.05$ .




Figure 4: Simulation performance comparison based on gene KRIT1. Plots of imputation  $\mathbb{R}^2$  in test samples by SUMMIT-FA, SUMMIT, Lasso sum, TWAS-FUSION, and PrediXcan for varied proportion of causal SNPs  $p_{causal}$ , DNAm heritability  $h_e^2$ , and phenotypic heritability  $h_p^2$ .




Figure 5: Simulation performance comparison based on gene *KRIT1*. Plots of statistical power for SUMMIT-FA, SUMMIT, Lassosum, TWAS-FUSION, and PrediXcan for varied proportion of causal SNPs  $p_{causal}$ , DNAm heritability  $h_e^2$ , and phenotypic heritability  $h_p^2$ . Empirical power was determined by the proportion of p-values  $< 2.5 \times 10^{-6}$ .