

1 *Effects of joint mobilization combined with acupuncture on pain,*
2 *physical function, and depression in stroke patients with chronic*
3 *neuropathic pain: a randomized controlled trial*

4
5 Ji-Eun Lee¹, Takayuki Akimoto², Ho-Seong Lee^{1,3*}

6 ¹Department of Exercise and Medical Science, Graduate School, Dankook University,
7 Cheonan, Republic of Korea

8 ²Laboratory of Muscle Biology, Faculty of Sport Sciences, Waseda University,
9 Tokorozawa, Japan

10 ³Institute of Medical-Sports, Dankook University, Cheonan, Republic of Korea

11

12 * Corresponding author

13 E-mail: hoseh28@dankook.ac.kr (HSL)

15

16 **Abstract**

17 Objective: To investigate the effectiveness of joint mobilization (JM) combined
18 with acupuncture (AC) for treatment pain, physical function and depression in
19 poststroke patients. Methods: A total of 69 poststroke patients were randomly
20 assigned to the JM+AC group (n = 23), the JM group (n = 23), and the control group
21 (n = 23). Patients in the JM+AC group and the JM group received JM for 30 minutes,
22 twice a week for 12 weeks, and the JM+AC group received AC for 30 minutes
23 separately once a week. The control group did not receive JM or AC. Pain (visual
24 analog scale, shoulder pain and disability index, Western Ontario and McMaster
25 universities osteoarthritis index), physical function (range of motion, 10-m walking
26 speed test, functional gait assessment, manual function test, activities of daily living
27 scale, instrumental activities of daily living scale), and depression (center for
28 epidemiologic studies depression scale, Beck depression inventory) were assessed for
29 each patient before and after the 12 weeks of intervention. Results: Pain and physical
30 function were improved significantly in the JM+AC group compared with the JM and
31 control groups. Physical function and depression were improved significantly in the
32 JM+AC and JM groups compared with the control group. Conclusion: The treatment
33 of JM combined with AC improved pain, depression, and physical function of
34 poststroke patients with chronic neuropathic pain in this study. This valuable finding

35 provides empirical evidence for the designing therapeutic interventions and
36 identifying potential therapeutic targets.

37 **Keywords:** acupuncture, joint mobilization, neuropathic pain, poststroke

38

39 **Introduction**

40 Stroke is one of global healthcare problems that is common, serious, and
41 disabling. Motor recovery from stroke is frequently poor or insufficient, with only
42 one-third of poststroke patients regaining dexterity within the first 6 months [1]. Less
43 than 45% of stroke patients are likely to achieve complete functional recovery, and
44 the majority of patients has some residual impairment and an inability to perform
45 activities of daily living (ADL) [2-4]. In addition to motor control and sensory
46 deficits, stroke survivors also develop complications such as pain, spasticity, joint
47 constraint, skin or vascular damage, and chronic neuropathic pain caused by nerve
48 damage, which are the main challenges in poststroke management [5,6].

49 Shoulder pain experienced by more than 50% of poststroke patients, is a
50 common sequelae [7], interfering with rehabilitation and reducing arm function and
51 ADL [8]. Knee pain is the second most common pain in patients after stroke, and it
52 has been reported to cause motor dysfunction and abnormal gait [9]. The recovery of
53 gait function is a primary goal of rehabilitation, with the majority of poststroke

54 patients suffering from a reduction in gait velocity and independent gait is crucial in
55 carrying out ADL following stroke [10]. Additionally, sensorimotor dysfunction
56 causes limited joint range of motion (ROM) and muscle weakness in the lower
57 extremities of the affected side, leading to difficulties in performing functional
58 activities, such as sit to stand and gait [11,12]. Therefore, it is necessary to reduce
59 pain and increase the joint ROM to improve physical function in poststroke patients.
60 To tackle these problems and to help restore and improve the function of upper and
61 lower limbs, a wide range of treatments has been proposed to therapists, so far.

62 Mulligan first proposed mobilization-with-movement treatment as a joint
63 mobilization (JM) technique [13]. JM is an orthopedic manipulation therapy used for
64 its mechanical effects to relieve pain, improve mobility, and treat contractures via the
65 third stage of the translatory movement and the convex–concave rule [14]. According
66 to the convex–concave rule, gliding on a convex joint surface is performed in the
67 opposite direction to the bone’s movement; gliding on a concave joint surface is
68 performed in the same direction as the bone’s movement. When a concave joint
69 surface is distracted, the joint is pulled along the long axis of the bone; alternatively,
70 when a convex joint surface is distracted, the joint surface must be separated [14]. JM
71 has the mechanical effect of disrupting the contracture by direct movement of the joint
72 area, and the arthrokinetic reflex effect of suppressing or facilitating the related
73 muscles is achieved by stimulating the joint receptors [15]. However, there appears to

74 be a limit to how JM alone can reduce pain and restore physical function in poststroke
75 patients.

76 Acupuncture (AC) is commonly used to treat/manage chronic diseases in Korea
77 [16]. It is recognized to be effective in regulating physiological balance by controlling
78 muscles and acupuncture points or by stimulating the affected area [17]. And AC is
79 effective in reducing pain by stimulating the nerves in stroke patients and help to
80 restore motor function and improve the ADL score [18]. Studies suggest that AC is
81 also effective in improving pain and depression in poststroke patients [19].
82 Particularly, AC combined with other therapies is more effective in poststroke
83 patients than AC alone [20]. Therefore, it is possible that AC combined with JM is
84 more effective to improve pain, depression, and physical function in poststroke
85 patients. Although there are many studies about effectiveness of JM for poststroke
86 patients, a few studies have examined the effects of JM combined with AC for these
87 patients.

88 In this work, we examined the changes in pain, physical function, and depression
89 before and after the 12 weeks of intervention of JM combined with AC in poststroke
90 patients. We hypothesized that combining JM with AC would improve physical
91 function more compared with JM alone.

92

93 **Methods**

94 **1. Study design**

95 The study had a random sampling design. Participants were randomized
96 electronically using block sizes of 3 across groups. All investigators and data analysts
97 were blinded until the study and analysis were completed.

98 Written informed consent was obtained from all subjects and the signed informed
99 consent documents were kept on file. The protocol was approved by the Dankook
100 University ethics committee (IRB number: 2021-05-046) and it adhered to the tenets
101 of the Declaration of Helsinki and its later amendments or comparable ethical
102 standards. The individual in this manuscript has given written informed consent (as
103 outlined in PLOS consent form) to publish these case details.

104 This clinical trial was registered with the Korean Clinical Research Information
105 Service approved by the WHO. Registration number: KCT0007924. Date registered:
106 22/11/2022, retrospectively registered. No dropouts or adverse events were reported.

107 **2. Participants**

108 This study included 69 adults who were diagnosed with stroke and were enrolled
109 in a public health center and welfare center located in D city, South Korea. This group
110 was further divided into the JM+AC, the JM, and the control groups (n = 23 each).

111 The inclusion criteria were: diagnosis of stroke \geq 6 months ago; pain in the shoulders
112 and knees for more than 6 months; and a Korean Mini Mental State Examination
113 score of \geq 24. The exclusion criteria were: a risk of tumors or infections; a previous
114 history of orthopedic disease or ankle surgery; a previous history of orthopedic
115 disease or shoulder and knee joint surgery; and a length of \geq 6.3 mm in the line
116 bisection test; taking medications for diseases other than stroke, such as antiplatelet
117 drugs and anticoagulants. The demographics and clinical characteristics of the study
118 patients were summarized at inclusion (Table 1). No significant differences in the
119 pretraining status of the control and experimental groups were noted.

120 **Table 1. Participant characteristics.**

Variables	JM+AC (n=23)	JM (n=23)	CON (n=23)	p
Age (years)	73.78 \pm 6.52	74.13 \pm 5.63	76.74 \pm 6.62	0.23
Sex (male/female)	12 / 11	17 / 6	13 / 10	0.29
Affected side (right/left)	13 / 10	10 / 13	13 / 10	0.60
Onset time (month)	52.57 \pm 24.39	61.35 \pm 21.00	61.09 \pm 22.20	0.33
MMSE-K (score)	26.17 \pm 1.34	26.35 \pm 1.44	25.87 \pm 1.40	0.51

Values are means \pm standard deviation.

MMSE-K: Korean mini mental state examination.

121 **3. Interventions**

122 **3.1. Joint mobilization**

123 For 12 weeks, the JM+AC and JM groups received grades II and III JM for 30
124 min, twice weekly, to relieve pain in the shoulder and knee joints and to improve the
125 ROM [14]. Shoulder JM was performed on the glenohumeral joint and
126 scapulothoracic joint in the supine and prone positions. Lateral distraction, downward
127 glide, posterior glide, and anterior glide were performed on the glenohumeral joint.
128 Elevation, depression, protraction, retraction, upward rotation, and downward rotation
129 were performed on the scapulothoracic joint. Knee JM was performed on the
130 tibiofemoral and patellofemoral joints in the supine and prone positions. Lower
131 distraction, posterior glide, and anterior glide were performed on the tibiofemoral
132 joint. Downward, lateral, and medial glides were performed on the patellofemoral
133 joint.

134 **3.2 Acupuncture**

135 The JM+AC group received acupuncture and moxibustion treatments around the
136 shoulder and the knee joints for 30 min, once a week for 12 weeks. Acupuncture
137 treatment was performed using disposable sterile stainless-steel needles (0.20 × 30
138 mm, Dongbang, Korea) for 15 min on average (figure 1). The penetration depth
139 varied according to the site and was 20 mm deep on average. For moxibustion
140 treatment, mini moxibustion (Taegeuk, Korea) was performed for 15 min (fig 1).

141 **Fig 1. Acupuncture and moxibustion treatment materials.**

142 Disposable sterile needles (left picture). Disposable mini moxa rolls (right picture).

143 **4. Outcome Measures**

144 **4.1 Pain**

145 Pain severity was measured using a visual analog scale (VAS), the shoulder pain
146 and disability index (SPADI), and the Korean Western Ontario McMaster Universities
147 osteoarthritis index (KWOMAC) before and after 12 weeks of the experimental
148 period. The VAS used a 100-mm line on which a pain-free state was 0 and severe pain
149 was 100, and the reliability is 0.99 [21]. SPADI is an evaluation tool for assessing the
150 degree of shoulder pain and disability that comprises 13 evaluation items scored on a
151 scale of 0–10 [22]. A score above 100 (out of 130) indicates more shoulder pain and
152 disability. The reliability is 0.94 [23]. KWOMAC, a Korean version of WOMAC
153 [24], is an evaluation tool for assessing the degree of knee pain and function [25]. It
154 consists of 24 evaluation items scored on a scale of 0–4 points and a total score of 96
155 points. Higher scores indicate a worse knee condition. The reliability is 0.96 [25].

156 **4.2 Physical function**

157 The range of motion (ROM), 10-meter walk test (10-MWT), functional gait
158 assessment (FGA), manual function test (MFT) for measuring physical function,
159 ADL, and instrumental ADL (IADL) were evaluated before and after 12 weeks of the
160 experimental period.

161 A goniometer (Saehan Corporation, Korea) was used to measure the ROM of the
162 shoulder and knee joint. Active ROM for shoulder flexion (SF), shoulder extension
163 (SE), shoulder abduction (SAB), shoulder adduction (SAD), and knee flexion (KF)
164 were measured. The measurement was performed three times, and the average value
165 was used. The reliability is 0.89 [26].

166 The 10-MWT requires the participant to walk independently over the middle 10
167 m portion on a 14-m path; the time is measured in seconds and the test is repeated
168 three times. The reliability is 0.99 [27].

169 FGA was developed by Wrisley et al. [28] to evaluate gait and comprises 10
170 items scored on a scale of 0–3 points and a total score of 30 points. Specific items
171 include walking on level surfaces, changing walking speed, turning the head to the
172 side while walking, moving the head up and down while walking, walking with one
173 foot as an axis, walking around obstacles, gait with a narrow base of support, walking
174 with eyes closed, walking backward, and climbing stairs. The reliability is 0.93 [28].

175 MFT is a test tool developed to assess the overall condition of arm function in a
176 stroke patient. It consists of eight items, and if subitem inspection is possible, it is
177 measured as 1 point. The reliability is 0.95 [29].

178 ADL and IADL are tools developed by Won et al. [30] ADL comprising seven
179 items covering dressing, washing the face, bathing, eating, moving, using the toilet,

180 and controlling the bladder, with a total of 21 points. IADL comprises 30 items scored
181 on a scale of 1–3. Ten items cover grooming, chores, preparing meals, washing
182 clothes, going out a short distance, using transport, buying goods, managing money,
183 using a phone, and taking medicine. Higher scores indicate lower dependence in
184 evaluating the independence level in ADL. The reliability is 0.93 [30].

185 **4.3 Depression**

186 Depression was measured using the center for epidemiologic studies depression
187 scale (CES-D) and the Beck depression inventory (BDI) before and after 12 weeks of
188 the experimental period.

189 CES-D is a self-reported depression evaluation tool [31], comprising 20
190 evaluation items scored on a scale of 0–3 points. Scores higher than 21 (out of 60)
191 indicate a higher frequency of depressive experiences. The reliability is 0.90 [31].

192 BDI is a tool for evaluating emotional, cognitive, motivational, and physiological
193 depression, and the Korean version was used [32]. It comprises 21 evaluation items
194 scored on a scale of 0–3 points and a total score of 63 points. The reliability is 0.84
195 [32].

196 **5. Statistical Analysis**

197 The data were analyzed statistically using SPSS 23.0 software (SPSS, Chicago,
198 IL, USA). General characteristics prior to the interventions were analyzed using

199 descriptive statistics. Two-way analysis of variance (2-way ANOVA) with repeated
200 measures was performed to verify the difference between groups and time periods.
201 Multiple comparisons were performed according to the post hoc test (Tukey HSD)
202 when there was a significant difference between groups and time periods. A paired t
203 test was used to compare variations in pain, physical function, and depression before
204 and after experimental period for each group. Statistical significance was reached
205 when $p < 0.05$.

206

207 **Results**

208 **1. Pain**

209 The changes in pain severity are shown in Table 2. The VAS was not statistically
210 significant between groups, but SPADI decreased significantly in the JM+AC group
211 compared to the JM group and the control group ($p < 0.05$), and KWOMAC
212 decreased significantly in the JM+AC group compared to the control group ($p < 0.05$).
213 The VAS, SPADI, and KWOMAC were decreased significantly in the JM+AC and
214 JM groups after 12 weeks of intervention ($p < 0.05$).

215 **Table 2. Pain before and after joint mobilization and joint mobilization**

216 **combined with acupuncture.**

Variables	JM+AC (n=23)			JM (n=23)			CON (n=23)			F	ES
	pre	post	t	pre	post	t	pre	post	t		
	75.65 ±	56.13 ±	5.18*	63.48 ±	56.30 ±	2.32*	62.83	67.17 ±	-1.50	0.97	0.28
VAS (mm)	18.55	5.80		3.45	4.35		±	18.52			
								14.99			
SPADI	68.90 ±	63.71 ±	6.11*	58.16 ±	56.79 ±	3.32*	58.56	58.86 ±	-1.31	4.78*	0.13
(score)	0.12	8.40†‡		11.92	11.98		±	10.27			
								10.32			
KWOMAC	73.40 ±	51.88 ±	5.93*	68.65 ±	59.57 ±	3.07*	68.61	71.43 ±	-1.70	3.32*	0.09
(score)	5.22	12.46†		11.95	9.80		±	12.87			
								11.40			

Values are means ± standard deviation.

VAS: visual analog scale, SPADI: shoulder pain and disability index, KWOMAC: Korean Western

Ontario and McMaster universities osteoarthritis index.

*p < 0.05 vs. pre-exercise. †p < 0.05 vs. CON. ‡p < 0.05 vs. JM.

217 **2. Physical function**

218 The changes in physical functions are shown in Table 3.

219 **Table 3. Physical function before and after joint mobilization and joint**

220 **mobilization combined with acupuncture.**

Variables		JM+AC (n=23)			JM (n=23)			CON (n=23)			F	ES
Period	pre	post	t	pre	post	t	pre	post	t			
	119.91	125.26	-4.74*	128.17	129.17	-1.17	128.6	124.48	4.18	1.12	0.03	
SF	± 14.00	± 13.26		± 14.46	± 13.21		5 ±	± 3.30				
							3.02					
SE	33.70 ±	38.96 ±	-5.79*	37.52 ±	37.83 ±	-0.36	34.48	32.39 ±	2.36	2.03	0.06	
	8.13	6.64		8.63	1.40		± 1.76	1.40				
SAB	109.13	114.70	-6.26*	117.44	118.52	-1.38	112.2	110.22	2.26	2.56	0.07	
ROM	± 11.93	± 10.17		± 2.46	± 2.02		1 ±	± 2.64				
(°)							2.41					
SAD	40.83 ±	47.65 ±	-7.13*	39.91 ±	42.39 ±	-1.89	41.96	40.00 ±	2.20	2.28	0.07	
(°)	6.34	6.77		1.64	1.47		± 1.10	0.89				
KF	117.31	127.17	-5.49*	119.91	125.65	-4.37*	112.8	112.39	0.44	7.27*	0.18	
	± 11.68	± 8.71†		± 2.02	± 1.70†		3 ±	± 2.65				
(°)							2.61					
10-M	25.83 ±	14.81 ±	7.44*	23.23 ±	18.13 ±	4.97*	24.55	24.04 ±	0.98	1.54	0.45	
GAIT WT	11.49	8.45		6.23	6.49		± 9.58	9.92				
(s)												

FGA	18.78 ± 23.13 ± -5.90*	19.04 ± 22.00 ± -4.22*	20.04	19.74 ± 0.79	0.75	0.22
(score	3.28	2.70	3.04	2.70	± 3.83	4.17
)						
MFT	20.78 ± 22.91 ± -6.16*	20.43 ± 21.52 ± -4.64*	20.00	19.52 ± 1.85	11.65	0.26
(score	1.57	1.73†‡	1.41	1.62†	± 1.71	1.68
)						*
ADL	17.57 ± 20.22 ± -5.42*	17.13 ± 19.30 ± -7.61*	16.65	16.57 ± 0.33	13.93	0.30
ADL (score	0.52	1.88†	0.39	0.19†	± 0.36	0.40
)						*
IADL	24.35 ± 26.96 ± -1.13*	23.13 ± 23.57 ± -0.93	22.96	22.70 ± 1.03	4.77*	0.13
(score	0.90	0.39†‡	0.81	0.72	± 0.80	0.70
)						

Values are means ± standard deviation.

ROM: range of motion, SF: shoulder flexion, SE: shoulder extension, SAB: shoulder abduction, SAD: shoulder adduction, KF: knee flexion, KE: knee extension, 10-MWT: 10-meter walking speed test, FGA: functional gait assessment, MFT: manual function test, ADL: activities of daily living scale, IADL: instrumental activities of daily living scale.

*p < 0.05 vs. pre-exercise. †p < 0.05 vs. CON. ‡p < 0.05 vs. JM.

222 SF, SE, SAB, and SAD were not statistically significant between groups, but KF
223 increased significantly in the JM+AC and JM groups compared to the control group
224 ($p < 0.05$). Within the JM+AC group, SF, SE, SAB, and SAD increased significantly
225 ($p < 0.05$), whereas KF increased significantly in the JM+AC and JM groups after 12
226 weeks of intervention ($p < 0.05$).

227 **2.2 Gait**

228 The 10-MWT and FGA were not statistically significant between groups, but
229 10-MWT and FGA improved significantly in the JM+AC and JM groups after 12
230 weeks of intervention ($p < 0.05$).

231 **2.3 Activities of daily living**

232 MFT increased significantly in the JM+AC group compared to the JM group and
233 control groups ($p < 0.05$), as well as in the JM group compared to the control group (p
234 < 0.05). ADL increased significantly in the JM+AC and JM groups compared to the
235 control group ($p < 0.05$), and IADL increased significantly in the JM+AC group
236 compared to the JM group and the control group ($p < 0.05$). Within the JM+AC and
237 JM groups, MFT and ADL increased significantly ($p < 0.05$), whereas IADL
238 increased significantly in the JM+AC group after 12 weeks of intervention ($p < 0.05$).

239 **3. Depression**

240 The changes in depression are shown in Table 4. CES-D and BDI decreased
 241 significantly in the JM+AC and JM groups compared to the control group ($p < 0.05$),
 242 and CES-D and BDI decreased significantly in the JM+AC group after 12 weeks of
 243 intervention ($p < 0.05$).

244 **Table 4. Depression before and after joint mobilization and joint mobilization**
 245 **combined with acupuncture.**

Variables	JM+AC (n=23)			JM (n=23)			CON (n=23)			F	ES
	pre	post	t	pre	post	t	pre	post	t		
CES-D	26.83 ± 8.63	20.30 ± 8.87†	5.24*	20.48 ± 9.57	20.43 ± 8.88	0.06	27.91 ± 8.96	33.00 ± 8.77	-3.60	7.11*	0.18
BDI (score)	32.48 ± 11.04	22.35 ± 7.97†	5.49*	30.61 ± 12.23	29.96 ± 10.99	1.19	33.70 ± 9.49	39.43 ± 9.15	-3.33	5.97*	0.15

Values are means ± standard deviation.

CES-D: center for epidemiologic studies depression scale, BDI: Beck depression inventory.

* $p < 0.05$ vs. pre-exercise. † $p < 0.05$ vs. CON. ‡ $p < 0.05$ vs. JM.

246

247 **Discussion**

248 The main finding of this study was that JM combined with AC treatment
249 improved the subjective pain, depression, and physical function of poststroke patients
250 more effectively than JM alone.

251 Shoulder and knee pain impedes rehabilitation and interferes with daily activities
252 in poststroke patients [7–9]. Kaltenborn et al. [14] indicated that JM was effective in
253 relieves shoulder and knee pain through direct movement of the joints. In AC
254 treatments, acupuncture treatment is reported to increase pain threshold by stimulating
255 the central nervous system, and moxibustion treatment has an analgesic effect through
256 thermal stimulation [18,33]. Additionally, previous studies reported that AC treatment
257 reduced SPADI and KOMAC [34,35]. Taken together, it was confirmed that JM
258 combined with AC treatment had a more positive effect on pain in poststroke patients
259 than JM alone.

260 Kluding and Santos [36] stated that the JM technique improved movement
261 without changing in joint ROM. Alternatively, Choi et al. [37] reported that AC
262 treatment reduces pain and increases joint flexibility by regenerating the surrounding
263 soft tissue. We found that there was a significant difference in ROM before and after
264 the intervention in the JM+ AC group, but not in the JM group in this study. These
265 results may imply that it is difficult to improve joint ROM with JM alone.

266 As mentioned in the introduction, gait is an important factor for independent
267 living and social participation. JM showed a positive effect on gait by improving
268 active ROM which is the ROM that can be achieved by patient himself/herself [38],
269 and AC treatment stimulated muscles and nerves to improve gait [39]. Additionally,
270 previous studies stated that acupuncture had a beneficial impact on the energy
271 production of joints during walking [40]. In this study, gait was significantly
272 improved after the intervention in the JM+ AC and JM groups, but there was no
273 difference between groups. Generally, gait necessitates movement of hip, knee, and
274 ankle joints. The lower extremity interventions in this study were limited to the knee
275 joint, which may not have been sufficient to influence group effect comparisons. In
276 the future, it would be prudent to intervene in all joints of lower extremity.

277 We employed several measurements to assess physical function, which is
278 important for daily living. MFT is an assessment of overall arm function [29], SPADI
279 is for shoulder pain and disability [22]. IADL is for evaluation of dressing, toilet
280 using, and movement [30]. It is thought that the reduction of SPADI restored arm
281 function in poststroke patients and affected MFT and IADL. Previous research has
282 shown that acupuncture has little effect on improving the lifestyle of stroke patients
283 [41], but it is believed that it had a positive effect on IADL in this study because joint
284 mobilization was combined with acupuncture. Arm function and daily life activities of
285 the patients in the JM+ AC group might be improved through the increase in muscle
286 activity induced by JM and the recovery of neurological damage by AC treatment.

287 Depression is associated with pain in poststroke patients [7], and depression in
288 these patients are associated with mortality, suicide risk, and quality of life [42–44].
289 Thus, depression must be verified in rehabilitation process for functional recovery
290 after stroke, in general. The CES-D and BDI scores in the JM+ AC group were
291 significantly decreased after 12 weeks of the intervention, but not in the JM group.
292 This is most likely due to AC treatment to alleviate pain and improve ADL. A
293 previous study showed that JM improved depression by reducing stress factors
294 through improvement of unstable posture [45]. Furthermore, it has been reported that
295 acupuncture treatment was effective in suppressing changes in the autonomic nervous
296 system caused by mental stress [46]. It has also been reported that moxibustion
297 treatment relaxed and stabilized the body through its thermal action [18]. Thus, it is
298 considered that acupuncture and moxibustion treatment of AC reduced depression
299 through autonomic nerve inhibition and thermal stimulation.

300 **Limitation**

301 There are several limitations in our study. The major part of subjects in this study
302 was older farmers living in rural areas, and it is not uncertain whether the results of
303 this study can be applied to other elderly population living in urban areas.
304 Additionally, it might be better if we assessed the effect of the treatment more
305 frequently because the results of this study were obtained before and after the 12
306 weeks of intervention period.

307

308 **Conclusion**

309 Improvements in pain, depression, and physical function of poststroke patients in
310 the JM+ AC group were significantly greater than those in the JM and the control
311 groups, suggesting JM combined with AC was more effective in improving not only
312 pain but also the ability to physical function. JM combined with AC may also be
313 helpful for poststroke patients who have depression and decreased physical function
314 due to chronic neuropathic pain. Future studies should reveal the physiological
315 assessment of muscle strength and the effect of JM combined with AC on muscle tone
316 based on changes in physical function in stroke patients with chronic neuropathic
317 pain.

318

319 **Author Contributions**

320 Conceived and designed the experiments: JEL. Performed the experiments: JEL.
321 Analyzed the data: JEL, TA. Wrote the paper: JEL, TA. Revised paper: TA, HSL.

322

323 **Conflicts of Interest**

324 The authors declare that there are no conflicts of interest regarding the publication of
325 this paper.

326

327 **Acknowledgments**

328 We appreciate Kim Yeon-oh and Choi Jong-bin of the Dangjin City Public Health
329 Center for their assistance with this experiment.

330

331 **Data Availability Statement**

332 Access to data was prepared using the following data set and is available at the free,
333 public repository Figshare (<https://doi.org/10.6084/m9.figshare.21789092>).

334

335 **Funding**

336 This research did not receive any specific grant from funding agencies in the public,
337 commercial, or not-for-profit sectors.

338

339 **References**

- 340 1. Dobkin BH. Rehabilitation after stroke. *N Engl J Med*. 2005;352(16):1677–84.
- 341 2. Flynn RW V, MacWalter RSM, Doney ASF. The cost of cerebral ischaemia.
342 *Neuropharmacology*. 2008;55(3):250–6.
- 343 3. Kelly-Hayes M, Beiser A, Kase CS, Scaramucci A, D’Agostino RB, Wolf PA.
344 The influence of gender and age on disability following ischemic stroke: the
345 Framingham study. *J Stroke Cerebrovasc Dis*. 2003;12(3):119–26.
- 346 4. Marini C, Baldassarre M, Russo T, De Santis F, Sacco S, Ciancarelli I, et al.
347 Burden of first-ever ischemic stroke in the oldest old: evidence from a
348 population-based study. *Neurology*. 2004;62(1):77–81.
- 349 5. De Ryck A, Brouns R, Franssen E, Geurden M, Van Gestel G, Wilssens I, et al. A
350 prospective study on the prevalence and risk factors of poststroke depression.
351 *Cerebrovasc Dis Extra*. 2013 Jan;3(1):1–13.
- 352 6. Vér C, Hofgárt G, Menyhárt L, Kardos L, Csiba L. Ankle-foot continuous
353 passive motion device for mobilization of acute stroke patients. *Open J Ther*
354 *Rehabil*. 2015;3(02):23.
- 355 7. Klit H, Finnerup NB, Overvad K, Andersen G, Jensen TS. Pain Following
356 Stroke: A Population-Based Follow-Up Study. *PLoS One*. 2011 Nov
357 15;6(11):e27607.

- 358 8. Rand D. Proprioception deficits in chronic stroke—Upper extremity function
359 and daily living. *PLoS One*. 2018 Mar 30;13(3):e0195043.
- 360 9. Caglar NS, Akin T, Aytakin E, Komut EA, Ustabasioglu F, Okur S, et al. Pain
361 syndromes in hemiplegic patients and their effects on rehabilitation results. *J*
362 *Phys Ther Sci*. 2016 Mar;28(3):731–7.
- 363 10. Yoo C, Yong M hyun, Chung J, Yang Y. Effect of computerized cognitive
364 rehabilitation program on cognitive function and activities of living in stroke
365 patients. *J Phys Ther Sci*. 2015;27(8):2487–9.
- 366 11. Fujita T, Iwata M, Fukuda M, Ikeda M. Relationship between lower extremity
367 muscle mass, leg extension strength and muscle power of hemiplegic stroke
368 patients. *J Phys Ther Sci*. 2011;23(2):277–82.
- 369 12. Lin PY, Yang YR, Cheng SJ, Wang RY. The relation between ankle
370 impairments and gait velocity and symmetry in people with stroke. *Arch Phys*
371 *Med Rehabil*. 2006;87(4):562–8.
- 372 13. Mulligan BR. Mobilisations with movement (MWMS) for the hip joint to restore
373 internal rotation and flexion. *J Man Manip Ther*. 1996;4(1):35–6.
- 374 14. Kaltenborn FM, Evjenth O. Manual Mobilization of the Joints: The Kaltenborn
375 Method of Joint Examination and Treatment. Vol 1. spine 5th ed Oslo Norli.
376 2009;

- 377 15. Ersoy U, Kocak UZ, Unuvar E, Unver B. The Acute Effect of Talocrural Joint
378 Mobilization on Dorsiflexor Muscle Strength in Healthy Individuals: A
379 Randomized Controlled Single-Blind Study. *J Sport Rehabil.* 2019;28(6):601–5.
- 380 16. Kim JH, Kismali G, Gupta SC. Natural Products for the Prevention and
381 Treatment of Chronic Inflammatory Diseases: Integrating Traditional Medicine
382 into Modern Chronic Diseases Care. *Evidence-Based Complement Altern Med.*
383 2018;2018:9837863.
- 384 17. Yang A, Wu HM, Tang J, Xu L, Yang M, Liu GJ. Acupuncture for stroke
385 rehabilitation. *Cochrane Database Syst Rev.* 2016;(8).
- 386 18. Zuo CY, Lv P, Zhang CS, Lei RX, Zhou W, Wu QF, et al. Ipsi- and Contralateral
387 Moxibustion Generate Similar Analgesic Effect on Inflammatory Pain. *Olajide*
388 *OA*, editor. *Evidence-Based Complement Altern Med.* 2019;2019:1807287.
- 389 19. Lee JA, Park SW, Hwang PW, Lim SM, Kook S, Choi KI, et al. Acupuncture for
390 shoulder pain after stroke: a systematic review. *J Altern Complement Med.* 2012
391 Sep;18(9):818–23.
- 392 20. Heo KH, Hwang EH, Cho HW, Lee I, Hong JW, Shin YI, et al. An analysis of
393 the effectiveness of stroke between east-west integrative medicine hospital and
394 western medicine hospital by the data of brain rehabilitation registry. *J Korean*
395 *Med Rehabil.* 2013;23(3):117–24.

- 396 21. Wagner DR, Tatsugawa K, Parker D, Young TA. Reliability and utility of a
397 visual analog scale for the assessment of acute mountain sickness. *High Alt Med*
398 *Biol.* 2007;8(1):27–31.
- 399 22. Roach KE, Budiman-Mak E, Songsiridej N, Lertratanakul Y. Development of a
400 shoulder pain and disability index. *Arthritis Rheum Off J Am Coll Rheumatol.*
401 1991;4(4):143–9.
- 402 23. Angst F, Goldhahn J, Pap G, Mannion AF, Roach KE, Siebertz D, et al.
403 Cross-cultural adaptation, reliability and validity of the German Shoulder Pain
404 and Disability Index (SPADI). *Rheumatology.* 2007;46(1):87–92.
- 405 24. Bellamy N. WOMAC Osteoarthritis Index: User Guide IV. WOMAC.
406 Queensland, Aust. 2000;
- 407 25. Bae SC, Lee HS, Yun HR, Kim TH, Yoo DH, Kim SY. Cross-cultural adaptation
408 and validation of Korean Western Ontario and McMaster Universities
409 (WOMAC) and Lequesne osteoarthritis indices for clinical research. *Osteoarthr*
410 *Cartil.* 2001;9(8):746–50.
- 411 26. Riddle, D. L., Rothstein, J. M., Lamb RL. Goniometric reliability in a clinical
412 setting: shoulder measurements. *Phys Ther.* 67(5):668–73.
- 413 27. Steffen TM, Hacker TA, Mollinger L. Age- and Gender-Related Test
414 Performance in Community-Dwelling Elderly People: Six-Minute Walk Test,

- 415 Berg Balance Scale, Timed Up & Go Test, and Gait Speeds. *Phys Ther.*
416 2002 Feb 1;82(2):128–37.
- 417 28. Wrisley DM, Marchetti GF, Kuharsky DK, Whitney SL. Reliability, internal
418 consistency, and validity of data obtained with the functional gait assessment.
419 *Phys Ther.* 2004;84(10):906–18.
- 420 29. Miyamoto S, Kondo T, Suzukamo Y, Michimata A, Izumi SI. Reliability and
421 validity of the Manual Function Test in patients with stroke. *Am J Phys Med*
422 *Rehabil.* 2009;88(3):247–55.
- 423 30. Chang-Won WON, Keum-Yeol Y, Yong-Gyun RHO, Soo-Young KIM,
424 Kyung-Hwan CHO, Ho-Cheol S, et al. The development of Korean activities of
425 daily living (K-ADL) and Korean instrumental activities of daily living
426 (K-IADL) scale. *J Korean Geriatr Soc.* 2002;107–20.
- 427 31. Radloff LS. The CES-D scale: A self-report depression scale for research in the
428 general population. *Appl Psychol Meas.* 1977;1(3):385–401.
- 429 32. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for
430 measuring depression. *Arch Gen Psychiatry.* 1961;4(6):561–71.
- 431 33. Lee JD, Chon JS, Jeong HK, Kim HJ, Yun M, Kim DY, et al. The
432 cerebrovascular response to traditional acupuncture after stroke. *Neuroradiology.*
433 2003;45(11):780–4.

- 434 34. Dyson-Hudson TA, Shiflett SC, Kirshblum SC, Bowen JE, Druin EL.
435 Acupuncture and Trager psychophysical integration in the treatment of
436 wheelchair user's shoulder pain in individuals with spinal cord injury. Arch Phys
437 Med Rehabil. 2001;82(8):1038–46.
- 438 35. Lo MY, Wu CH, Luh JJ, Wang TG, Fu LC, Lin JG, et al. The effect of
439 electroacupuncture merged with rehabilitation for frozen shoulder syndrome: A
440 single-blind randomized sham-acupuncture controlled study. J Formos Med
441 Assoc. 2020;119(1, Part 1):81–8.
- 442 36. Kluding PM, Santos M. Effects of ankle joint mobilizations in adults poststroke:
443 a pilot study. Arch Phys Med Rehabil. 2008 Mar;89(3):449–56.
- 444 37. Choi G cheol, Lee J won, Bae JE, Kim D jin, Hong J su, Kyung D hyun. The
445 Clinical Effect of Complex Korean Medical Admission Treatment in Patients
446 with Fractures of Distal Radius by Traffic Accident: 2 Cases Series Report. J
447 Korean Med Rehabil. 2021;31(1):187–93.
- 448 38. Ravichandran H, Janakiraman B. The Effects of Ankle Joint Mobilization on
449 Dorsiflexion Range and Gait Parameters in Chronic Stroke Survivors: A
450 Systematic Review and Meta-analysis. J Stroke Med. 2021;4(1):15–24.
- 451 39. Kim MS, Moon BS. Effect of comprehensive Korean medicine with
452 rehabilitation in stroke patients: a retrospective study. J Physiol Pathol Korean
453 Med. 2016;30(5):355–9.

- 454 40. Hauer K, Wendt I, Schwenk M, Rohr C, Oster P, Greten J. Stimulation of
455 acupoint ST-34 acutely improves gait performance in geriatric patients during
456 rehabilitation: a randomized controlled trial. *Arch Phys Med Rehabil.*
457 2011;92(1):7–14.
- 458 41. Sibbritt D, Peng W, Lauche R, Ferguson C, Frawley J, Adams J. Efficacy of
459 acupuncture for lifestyle risk factors for stroke: A systematic review. *PLoS One.*
460 2018;13(10):e0206288.
- 461 42. Kellermann M, Berecz R, Bereczki D. Does the severity of depressive symptoms
462 after stroke affect long-term survival? An 18-year follow-up. *PLoS One.*
463 2018;13(12):e0209157.
- 464 43. Wilkins SS, Akhtar N, Salam A, Bourke P, Joseph S, Santos M, et al. Acute post
465 stroke depression at a Primary Stroke Center in the Middle East. *PLoS One.*
466 2018;13(12):e0208708.
- 467 44. Kang SW, Lee SH, Lee YS, Kwon S, Bosch P, Lee YJ, et al. Association
468 between depression and quality of life in stroke patients: The Korea National
469 Health and Nutrition Examination Survey (KNHANES) IV–VII (2008–2018).
470 *PLoS One.* 2022;17(6):e0269010.
- 471 45. Park D, Lee E, Lee K. Effects of Thoracic Flexibility Exercise Program on Pain,
472 Heart Rate Variability, and Depression in Patients with Chronic Low Back Pain.
473 *J Korean Soc Integr Med.* 2019;7(4):161–70.

- 474 46. Cho SY, Ha SY, Jang JY, Nam SS, Kim YS. Original Article: Effect of
475 Hwangryunhaedoktang Pharmacopuncture at CV17 (Jeonjung) for Mental
476 Stress on Short-term Analysis of Heart Rate Variability. J Korean Acupunct
477 Moxibustion Soc. 2009;26(5):50.

479

480 **Supporting Information**

481 **S1 Fig. Acupuncture and moxibustion treatment materials.** Disposable sterile
482 needles (left picture). Disposable mini moxa rolls (right picture).

483 **S1 Table. Participant characteristics.** MMSE-K: Korean mini mental state
484 examination.

485 **S2 Table. Pain before and after joint mobilization and joint mobilization
486 combined with acupuncture.** VAS: visual analog scale, SPADI: shoulder pain and
487 disability index, KWOMAC: Korean Western Ontario and McMaster universities
488 osteoarthritis index.

489 **S3 Table. Physical function before and after joint mobilization and joint
490 mobilization combined with acupuncture.** ROM: range of motion, SF: shoulder
491 flexion, SE: shoulder extension, SAB: shoulder abduction, SAD: shoulder adduction,
492 KF: knee flexion, KE: knee extension, 10-MWT: 10-meter walking speed test, FGA:
493 functional gait assessment, MFT: manual function test, ADL: activities of daily living
494 scale, IADL: instrumental activities of daily living scale.

495 **S4 Table. Depression before and after joint mobilization and joint mobilization
496 combined with acupuncture.** CES-D: center for epidemiologic studies depression
497 scale, BDI: Beck depression inventory.

498

499 **S1 Checklist. CONSORT Checklist.** Checklist of information to include when

500 reporting a randomized trial.

501 **S2 Checklist. CONSORT Flow Diagram.** Flow diagram.



Figure