

Supplementary Table S1. Association analysis of each *HLA-DRB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=160)	P _{uncorr} ¹	OR (95%CI)	log-rank P _{uncorr} ²
	n (%)	n (%)			
<i>DRB1*01:01</i>	5 (12.8)	28 (17.5)	0.63	0.69 (0.25-1.93)	0.53
<i>DRB1*04:01</i>	1 (2.6)	1 (0.6)	0.35	4.18 (0.26-68.4)	0.37
<i>DRB1*04:05</i>	7 (17.9)	35 (21.9)	0.67	0.78 (0.32-1.92)	0.58
<i>DRB1*04:06</i>	4 (10.3)	8 (5.0)	0.26	2.17 (0.62-7.62)	0.25
<i>DRB1*04:10</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.65
<i>DRB1*08:02</i> ³	7 (17.9)	11 (6.9)	0.055	2.96 (1.07-8.23)	0.040
<i>DRB1*08:03</i>	4 (10.3)	15 (9.4)	0.77	1.10 (0.35-3.53)	0.83
<i>DRB1*09:01</i> ³	22 (56.4)	62 (38.8)	0.049	2.05 (1.01-4.15)	0.049
<i>DRB1*11:01</i>	1 (2.6)	8 (5.0)	1.00	0.50 (0.06-4.12)	0.47
<i>DRB1*12:01</i>	3 (7.7)	11 (6.9)	0.74	1.13 (0.30-4.26)	0.87
<i>DRB1*12:02</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.58
<i>DRB1*13:02</i>	1 (2.6)	14 (8.8)	0.31	0.27 (0.03-2.15)	0.22
<i>DRB1*14:05</i>	1 (2.6)	5 (3.1)	1.00	0.82 (0.09-7.19)	0.79
<i>DRB1*14*54</i>	3 (7.7)	15 (9.4)	1.00	0.81 (0.22-2.93)	0.79
<i>DRB1*15:01</i>	5 (12.8)	16 (10.0)	0.57	1.32 (0.45-3.86)	0.60
<i>DRB1*15:02</i>	9 (23.1)	34 (21.3)	0.83	1.11 (0.48-2.56)	0.80
<i>DRB1*16:02</i>	1 (2.6)	3 (1.9)	0.59	1.38 (0.14-13.6)	0.74

OR: odds ratio, CI: confidence interval.

*¹Carrier frequency (%) of each *HLA-DRB1* allele was compared between the patients with and without relapse by Fisher's exact test. Uncorrected P values (P_{uncorr}) are shown.

*²Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DRB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³For *DRB1*08:02* and *DRB1*09:01*, FDR Q values are shown in the main document (Table 4).

Supplementary Table S2. Association analysis of each *HLA-DQA1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39) n (%)	Relapse (-: n=159) n (%)	P_{uncorr}^{*1}	OR (95%CI)	log-rank P_{uncorr}^{*2}
<i>DQA1*01:01</i>	5 (12.8)	28 (17.6)	0.63	0.69 (0.25-1.91)	0.52
<i>DQA1*01:02</i>	7 (17.9)	30 (18.9)	1.0	0.94 (0.38-2.33)	0.96
<i>DQA1*01:03</i>	13 (33.3)	47 (29.6)	0.70	1.19 (0.56-2.52)	0.65
<i>DQA1*01:04</i>	3 (7.7)	21 (13.2)	0.42	0.55 (0.15-1.94)	0.38
<i>DQA1*03:01</i>	7 (17.9)	27 (17)	1.0	1.07 (0.43-2.67)	0.94
<i>DQA1*03:02</i> ^{*3}	24 (61.5)	64 (40.3)	0.020	2.38 (1.16-4.87)	0.020
<i>DQA1*03:03</i>	9 (23.1)	37 (23.3)	1.0	0.99 (0.43-2.27)	0.97
<i>DQA1*04:01</i>	4 (10.3)	5 (3.1)	0.077	3.52 (0.90-13.8)	0.075
<i>DQA1*05:05</i>	1 (2.6)	14 (8.8)	0.31	0.27 (0.03-2.14)	0.18
<i>DQA1*05:08</i>	2 (5.1)	3 (1.9)	0.26	2.81 (0.45-17.4)	0.18
<i>DQA1*06:01</i>	1 (2.6)	4 (2.5)	1.0	1.02 (0.11-9.39)	0.70

OR: odds ratio, CI: confidence interval.

^{*1}Carrier frequency (%) of each *HLA-DQA1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

^{*2}Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DQA1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³ For *DQA1*03:02*, FDR Q value is shown in the main document (Table 4).

Supplementary Table S3. Association analysis of each *HLA-DQB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39) n (%)	Relapse (-: n=160) n (%)	P_{uncorr}^{*1}	OR (95%CI)	log-rank P_{uncorr}^{*2}
<i>DQB1*03:01</i>	5 (12.8)	36 (22.5)	0.27	0.51 (0.18-1.39)	0.21
<i>DQB1*03:02</i>	7 (17.9)	27 (16.9)	0.82	1.08 (0.43-2.69)	0.93
<i>DQB1*03:03</i> ^{*3}	23 (59.0)	65 (40.6)	0.048	2.10 (1.03-4.28)	0.043
<i>DQB1*04:01</i>	7 (17.9)	34 (21.3)	0.83	0.81 (0.33-2.00)	0.63
<i>DQB1*04:02</i>	5 (12.8)	10 (6.3)	0.18	2.21 (0.71-6.87)	0.18
<i>DQB1*05:01</i>	5 (12.8)	28 (17.5)	0.63	0.69 (0.25-1.93)	0.53
<i>DQB1*05:02</i>	2 (5.1)	12 (7.5)	1.0	0.67 (0.14-3.11)	0.65
<i>DQB1*05:03</i>	2 (5.1)	12 (7.5)	1.0	0.67 (0.14-3.11)	0.63
<i>DQB1*06:01</i>	13 (33.3)	47 (29.4)	0.70	1.20 (0.57-2.54)	0.63
<i>DQB1*06:02</i>	5 (12.8)	15 (9.4)	0.55	1.42 (0.48-4.18)	0.50
<i>DQB1*06:04</i>	1 (2.6)	13 (8.1)	0.31	0.30 (0.04-2.35)	0.26

OR: odds ratio, CI: confidence interval.

^{*1}Carrier frequency (%) of each *HLA-DQB1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

^{*2}Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DQB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

*³For *DQB1*03:03*, FDR Q value is shown in the main document (Table 4).

Supplementary Table S4. Association analysis of each *HLA-DPB1* allele with occurrence of relapse and time to relapse

	Carrier frequency		Relapse-free survival		
	Relapse (+: n=39)	Relapse (-: n=160)	P _{uncorr} ^{*1}	OR (95%CI)	log-rank P _{uncorr} ^{*2}
	n (%)	n (%)			
<i>DPB1*02:01</i>	22 (56.4)	72 (45.0)	0.22	1.58 (0.78-3.20)	0.27
<i>DPB1*02:02</i>	2 (5.1)	13 (8.1)	0.74	0.61 (0.13-2.83)	0.58
<i>DPB1*03:01</i>	5 (12.8)	14 (8.8)	0.54	1.53 (0.52-4.55)	0.38
<i>DPB1*04:01</i>	1 (2.6)	10 (6.3)	0.70	0.39 (0.05-3.18)	0.41
<i>DPB1*04:02</i>	8 (20.5)	42 (26.3)	0.54	0.73 (0.31-1.70)	0.54
<i>DPB1*05:01</i>	22 (56.4)	87(54.4)	0.86	1.09 (0.54-2.20)	0.81
<i>DPB1*09:01</i>	8 (20.5)	31 (19.4)	0.83	1.07 (0.45-2.56)	0.89
<i>DPB1*13:01</i>	2 (5.1)	4 (2.5)	0.33	2.11 (0.37-11.9)	0.45

OR: odds ratio, CI: confidence interval.

*¹Carrier frequency (%) of each *HLA-DPB1* allele was compared between the patients with and without relapse by Fisher's exact test.

Uncorrected P values (P_{uncorr}) are shown.

*²Time to relapse after achievement of remission was compared between the patients carrying and not-carrying each *HLA-DPB1* allele. Log-rank P_{uncorr} values for the relapse-free survival rate calculated by Kaplan-Meier method are shown.

Supplementary Table S5 Association analyses of HLA-DRβ1_13S encoding alleles with susceptibility and relapse-free survival.

HLA amino acid	susceptibility			relapse-free survival		
	MPO-AAV (2n=856)	healthy controls (2n=1558)	P _{uncorr}	OR / 95% CI	log-rank P _{uncorr}	HR / 95%CI
DRβ1_13S	130 (15.2)	312 (20.0)	0.0029	0.71 / 0.56-0.89	0.015	0.31 / 0.12-0.80
<i>HLA-allele encoding DRβ1_13S</i>						
<i>DRB1*11:01</i>	24 (2.8)	30 (1.9)	0.16	1.48 / 0.85-2.57	0.47	0.49 / 0.07-3.55
<i>DRB1*13:02</i>	32 (3.7)	126 (8.1)	6.9x10 ⁻⁵	0.45 / 0.30-0.66	0.22	0.31 / 0.04-2.24
<i>DRB1*14:03</i>	22 (2.6)	32 (2.1)	0.41	1.26 / 0.72-2.19	-	-
<i>DRB1*14:05</i>	15 (1.8)	48 (3.1)	0.054	0.56 / 0.30-0.98	0.79	0.77 / 0.11-5.59
<i>DRB1*14:06</i>	5 (0.6)	16 (1.0)	0.27	0.56 / 0.18-1.45	-	-
<i>DRB1*14:54</i>	31 (3.6)	50 (3.2)	0.59	1.13 / 0.71-1.78	0.79	0.85 / 0.26-2.76

Associations with susceptibility and relapse-free survival of HLA-DRβ1_13S encoding alleles in the subjects of this study. Allele frequencies are shown in parentheses. Association with susceptibility was examined using logistic regression analysis under the additive model. Only the alleles with minor allele frequency >0.01 are listed.

The data on relapse-free survival are derived from Table 5 (main text) and Supplementary Table S1. Carriers of *DRB1*14:03* and *DRB1*14:06* were not present in the subjects with relapse.

OR: odds ratio, CI: confidence interval, HR: hazard ratio.