

Supplemental data

Gene	Primer sequences	Product size	Tm	# cycles
<i>Ccr5</i> (qPCR)	F: GTT CCT GAA AGC GGC TGT AAA R: GCA GTC AGG CAC ATC CAT AGA C	74	-	-
<i>Cxcr4</i> (qPCR)	F: TGG CAT AGT CGG CAA TGG A R: CGT CAT GCT CCT TAG CTT CTT CT	66	-	-
<i>Tbp</i> (qPCR)	F: TTGACCTAAAGACCATTGCACTTC R: TTCTCATGATGACTGCAGCAAA	78	-	-
<i>Rps9</i> (qPCR)	F: GAC CAG GAG CTA AAG TTG ATT GGA R: TCT TGG CCA GGG TAA ACT TGA	81	-	-
<i>Ccl3</i>	F: ATG AAG GTC TCC ACC ACT GC R: CCC AGG TCT CTT TGG AGT CA	279	60	32
<i>Ccl4</i>	F: GCC CTC TCT CTC CTC TTG CT R: GTC TGC CTC TTT TGG TCA GG	196	60	32
<i>Ccl5</i>	F: CCC TCA CCA TCA TCC TCA CT R: CCT TCG AGT GAC AAA CAC GA	185	60	32
<i>L32</i>	F: GTG AAG CCC AAG ATC GTC AA R: TTG GTG ACT CTG ATG GCC AG	349	58	26
<i>Ccr5</i> promoter site A	F: CTT GAC TCA AAT GTG GGC TTT R: GGC TGA GAG GTG ACT TAC CA	400	58	40
<i>Ccr5</i> promoter site B	F: TTT AAA CAG GGC AAG CCA GT R: TTC TCA GCT GCA AGA AGC AA	379	58	40

Supplemental table 1. Oligonucleotide primers and conditions used for real-time PCR (qPCR) and semi-quantitative end-point RT-PCR. Ccl: Chemokine (C-C motif) ligand; Ccr5: C-C chemokine receptor type 5; Rps9: Ribosomal Protein S9; Tbp: TATA-binding protein

	Cognitively normal subjects		All type of dementia		Alzheimer's disease		Vascular and mixed dementia ^A	
Age ^{B,C}	80.3	7.8	81.1	7.0	81.8	6.1	80.3	7.6
Female ^C	238	67.7%	592	73.2%	306	78.1%	276	68.5%
ApoEε4 / CCR5 ⁺	261	72.1%	414	51.2%	197	50.3%	207	51.4%
ApoEε4 / CCR5-Δ32	49	13.5%	86	10.6%	41	10.5%	41	10.2
ApoEε4 ⁺ / CCR5 ⁺	46	12.7%	271	33.5%	138	35.2%	133	33%
ApoEε4 ⁺ / CCR5-Δ32	6	1.7%	38	4.7%	16	4.1%	22	5.5%
Total	362	100.0%	809	100.0%	392	100.0%	403	100.0%

Supplemental table 2. A comparison of CCR5-Δ32 and ApoEε4 allele frequencies between dementia patients, including the various dementia aetiologies, and subjects that are cognitively normal in the Italian and Swiss samples. To measure the impact of cerebral vascular lesions, patients with vascular and mixed dementia were analysed in the same group. Data are expressed as number of cases and %. ^aData are expressed as means ± SD.

ApoEε4⁺: one or two copies of ApoEε4; ApoEε4⁻: no copies of ApoEε4; CCR5Δ32⁺: one or two copies of the CCR5-32 bp deleted allele; CCR5Δ32⁻: two copies of the CCR5 wild-type allele.

	<i>Dementia</i> (n = 809)			<i>Alzheimer's disease</i> (n = 392)			<i>Vascular or mixed dementia</i> (n = 403)		
	Crude OR	95% CI	p	Crude OR	95% CI	p	Crude OR	95% CI	p
Univariate logistic regression									
ApoEε4 ⁺ ^A	3.08^C	2.20-4.31	<0.001	3.21	2.20-4.69	<0.001	3.25	2.25-4.71	<0.001
Italy vs Swiss	3.80	2.90-4.97	<0.001	5.11	3.65-7.15	<0.001	3.48	2.54-4.77	<0.001
CCR5-Δ32 ^B	1.37	0.95-1.99	0.095	1.22	0.78-1.91	0.374	1.49	0.97-2.28	0.068
Italy vs Swiss	4.44	3.40-5.81	<0.001	5.80	7.17-8.08	<0.001	4.08	2.98-5.59	<0.001
Age	1.08	1.06-1.11	<0.001	1.14	1.11-1.18	<0.001	1.07	1.04-1.10	<0.001
Italy vs Swiss	7.84	5.64-10.9	<0.001	17.41	10.9-27.8	<0.001	6.98	4.71-10.3	<0.001
Male vs female	0.63	0.48-0.84	<0.001	0.42	0.29-0.60	<0.001	0.79	0.57-1.08	0.141
Italy vs Swiss	4.41	3.38-5.76	<0.001	6.37	4.53-8.94	<0.001	3.93	2.89-5.35	<0.001
Multiple logistic regression									
	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p
<i>Association ApoEε4 / CCR5Δ32</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5-Δ32	1.44	0.95-2.16	0.083	1.39	0.85-2.29	0.191	1.46	0.90-2.38	0.123
ApoEε4 ⁺ / CCR5 ⁺	3.11	2.17-4.46	<0.001	3.35	2.23-5.02	<0.001	3.18	2.14-4.73	<0.001
ApoEε4 ⁺ / CCR5-Δ32	4.51	1.83-11.1	0.001	3.68	1.32-10.3	0.013	5.76	2.21-15.0	<0.001
Italy vs Swiss	3.95	3.00-5.20	<0.001	5.23	3.72-7.34	<0.001	3.70	2.68-5.12	<0.001
<i>Adjusted for age and sex</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5Δ32	1.42	0.93-2.17	0.108	1.25	0.73-2.15	0.414	1.50	0.91-2.48	0.109
ApoEε4 ⁺ / CCR5 ⁺	3.21	2.22-4.63	<0.001	3.70	2.39-5.72	<0.001	3.22	2.15-4.82	<0.001
ApoEε4 ⁺ / CCR5Δ32	4.71	1.82-12.2	0.001	4.03	1.15-14.1	0.029	5.94	2.19-16.1	<0.001
Age	1.08	1.06-1.10	<0.001	1.14	1.10-1.18	<0.001	1.07	1.04-1.10	<0.001
Male vs female	0.69	0.51-0.93	0.014	0.42	0.28-0.62	<0.001	0.83	0.59-1.16	0.28
Italy vs Swiss	7.29	5.18-10.3	<0.001	16.67	10.3-27.0	<0.001	6.78	4.49-10.2	<0.001

Supplemental Table 3. Univariate and multiple logistic regression models of the risk of dementia and of its main aetiologies with cognitively normal subjects as the reference group (n = 362) according to CCR5 and ApoE genotypes in the Italian and Swiss population. Models are adjusted by country.

^AApoEε4⁺: one or two copies of ApoEε4; ApoEε4⁻: no copies of ApoEε4; ^BCCR5Δ32⁺: one or two copies of the CCR5-32 bp deleted allele; CCR5Δ32⁻: two copies of the CCR5 wild-type allele. ^CBold entries = relevant results.

	<i>Dementia</i> (n = 317)			<i>Alzheimer's disease</i> (n = 126)			<i>Vascular or mixed dementia</i> (n = 186)		
	Crude OR	95% CI	p	Crude OR	95% CI	p	Crude OR	95% CI	p
Univariate logistic regression									
ApoEε4 ⁺ A	4.19^C	2.60-6.76	<0.001	4.46	2.56-7.77	<0.001	4.17	2.49-6.99	<0.001
Italy vs Swiss	2.87	1.70-4.86	<0.001	3.81	1.75-8.30	0.001	3.34	1.75-6.36	<0.001
CCR5-Δ32 ^B	0.95	0.53-1.69	0.859	0.77	0.35-1.69	0.509	1.13	0.59-2.18	0.706
Italy vs Swiss	3.66	2.18-6.14	<0.001	4.59	2.13-9.90	<0.001	4.36	2.30-8.29	<0.001
Age	1.05	1.00-1.10	0.036	1.18	1.08-1.28	<0.001	1.03	0.98-1.08	0.248
Italy vs Swiss	3.96	2.37-6.62	<0.001	6.61	2.96-14.77	<0.001	4.45	2.37-8.36	<0.001
Male vs female	0.94	0.64-1.39	0.759	0.67	0.40-1.10	0.114	1.05	0.68-1.63	0.828
Italy vs Swiss	3.70	2.23-6.14	<0.001	5.18	2.42-11.1	<0.001	4.24	2.27-7.91	<0.001
Multiple logistic regression									
	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p
<i>Association ApoEε4 / CCR5Δ32</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5-Δ32	1.14	0.59-2.18	0.699	1.00	0.40-2.45	0.994	1.32	0.62-2.80	0.471
ApoEε4 ⁺ / CCR5 ⁺	4.53	2.71-7.59	<0.001	4.80	2.66-8.66	<0.001	4.47	2.57-7.79	<0.001
ApoEε4 ⁺ / CCR5-Δ32	2.72	0.82-9.05	0.103	2.22	0.44-11.14	0.331	3.43	0.95-12.4	0.059
Italy vs Swiss	2.86	1.67-4.89	<0.001	3.57	1.61-7.90	0.002	3.45	1.78-6.70	<0.001
<i>Adjusted for age and sex</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5Δ32	1.09	0.57-2.11	0.788	1.01	0.40-2.56	0.980	1.31	0.61-2.79	0.488
ApoEε4 ⁺ / CCR5 ⁺	4.39	2.61-7.37	<0.001	4.44	2.42-8.16	<0.001	4.42	2.53-7.75	<0.001
ApoEε4 ⁺ / CCR5Δ32	2.65	0.79-8.93	0.116	2.36	0.41-13.47	0.334	3.40	0.94-12.3	0.063
Age	1.03	0.98-1.08	0.207	1.14	1.04-1.24	0.003	1.01	0.96-1.06	0.764
Male vs female	0.90	0.60-1.37	0.630	0.61	0.35-1.07	0.084	0.99	0.62-1.58	0.964
Italy vs Swiss	2.99	1.74-5.15	<0.001	4.96	2.14-11.52	<0.001	3.49	1.79-6.82	<0.001

Supplemental Table 4. Bivariate and multiple logistic regression models of the risk of dementia and of its main aetiologies with cognitively normal subjects as the reference group (n = 177) according to CCR5 and ApoE genotypes in the Italian and Swiss populations aged < 80. Models are adjusted by country.

	Dementia (n = 189)			Alzheimer's disease (n = 73)			Vascular or mixed dementia (n = 102)		
	Univariate logistic regression								
	Crude OR	95% CI	p	Crude OR	95% CI	p	Crude OR	95% CI	p
ApoEε4 ⁺ A	2.51^C	1.53-4.12	<0.001	2.63	1.48-4.69	0.001	2.75	1.58-4.78	<0.001
Italy vs Swiss	10.88	6.95-17.0	<0.001	16.44	10.01-27.0	<0.001	8.12	4.96-13.3	<0.001
CCR5-Δ32 ^B	1.70	1.02-2.84	0.041	1.34	0.71-2.51	0.370	1.98	1.10-3.56	0.022
Italy vs Swiss	12.16	7.77-19.0	0.000	17.85	10.9-29.2	<0.001	9.19	5.59-15.1	<0.001
Age	1.04	0.99-1.09	0.099	1.04	0.99-1.10	0.135	1.05	1.00-1.11	0.056
Italy vs Swiss	13.37	8.30-21.5	<0.001	20.85	12.1-36.0	<0.001	10.26	6.00-17.5	<0.001
Male vs female	0.53	0.34-0.82	0.005	0.33	0.18-0.59	<0.001	0.63	0.38-1.04	0.072
Italy vs Swiss	11.85	7.57-18.5	<0.001	19.73	11.7-33.1	<0.001	8.66	5.30-14.1	<0.001
	Multiple logistic regression								
	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p	Adjusted OR	95% CI	p
<i>Association ApoEε4 / CCR5Δ32</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5-Δ32	1.62	0.92-2.82	0.093	1.41	0.70-2.82	0.336	1.72	0.89-3.33	0.106
ApoEε4 ⁺ / CCR5 ⁺	2.35	1.38-4.01	0.002	2.63	1.42-4.87	0.002	2.42	1.32-4.42	0.004
ApoEε4 ⁺ / CCR5-Δ32	7.65	1.70-34.5	0.008	4.86	0.88-26.8	0.07	11.19	2.36-53.0	0.002
Italy vs Swiss	11.40	7.26-17.9	<0.001	16.6	10.08-27.3	<0.001	8.91	5.39-14.7	<0.001
<i>Adjusted for age and sex</i>									
ApoEε4 ⁻ / CCR5 ⁺	1.00	--	--	1.00	--	--	1.00	--	--
ApoEε4 ⁻ / CCR5Δ32	1.64	0.93-2.90	0.086	1.39	0.68-2.81	0.364	1.76	0.90-3.43	0.098
ApoEε4 ⁺ / CCR5 ⁺	2.53	1.48-4.34	0.001	3.15	1.65-6.00	<0.001	2.57	1.40-4.74	0.002
ApoEε4 ⁺ / CCR5Δ32	7.48	1.64-34.1	0.009	6.11	0.94-39.6	0.058	10.55	2.21-50.2	0.003
Age	1.05	1.00-1.10	0.056	1.05	0.99-1.11	0.1	1.06	1.00-1.12	0.038
Male vs female	0.51	0.33-0.81	0.004	0.29	0.16-0.53	0.000	0.64	0.38-1.08	0.093
Italy vs Swiss	13.73	8.38-22.5	<0.001	22.68	12.6-40.8	<0.001	11.60	6.58-20.4	<0.001

Supplemental Table 5. Bivariate and multiple logistic regression models of the risk of dementia and of its main aetiologies with cognitively normal subjects as the reference group (n = 185) according to CCR5 and ApoE genotypes in the Italian and Swiss populations aged ≥ 80. Models are adjusted by country.