



Supplemental Figure 1



Filter	Genes	Percentage
Total RNAs	60,754	100%
Total number of non-coding RNAs	16,892	26.8%
≥ 0.1 TPM $\geq 10\%$ threshold	7,537	12.4%


Supplemental Figure 1: lncRNA distribution in sequenced neurons. Left panel, distribution of lncRNA expression (including all lncRNAs). Right panel, after applying thresholds of 0.1 TPM expression in 10% of samples, histogram captures normal distribution. Table summarizing the numbers of genes at each stage. *TPM: transcripts per kilobase Million. Normalized for gene length and for sequence depth

Supplemental Figure 2



Supplemental Figure 2: *Norad* is differentially expressed in *MAPT* p.P301L mice.
Normalized read counts (TPM) of *Norad* in WT mice and Tau-P301L mice. *, $p < 0.05$.

Supplemental Figure 3



Supplemental Figure 3: Ectopic expression of *lncRNA-SNHG8* in HEK293T cells. A. Representative fluorescence images of HEK293T cells transfected with pCDNA3.1+-EGFP and pCDNA3.1+-SNHG8-EGFP plasmid constructs and images were taken 48hrs post transfection. The fluorescence images were overlaid with specific phase contrast image. B. Relative expression of *SNHG8* in HEK293T cells transfected with pCDNA3.1+-EGFP or pCDNA3.1+-SNHG8-EGFP plasmids. Expression normalized to GAPDH. n=4. Data are represented as mean \pm SEM. **** p < 0.0001; unpaired Student's t-test.