

Supplemental material

Table S1: Characteristics of the group of participants who had two pre-surgery sessions

	Baseline 1	Baseline 2	<i>p</i> value
<i>N</i>	19	19	
Sexe (F:M)	13 : 6	13 : 6	
Age (years)	43.8 ± 10.2	43.9 ± 9.0	0.9884 ; t=2.02619
Diabetic (Y:N)	6 : 13	6 : 13	
BMI (kg/m ²)	41.7 ± 3.8	42.0 ± 3.6	0.9703; t=2.02809
Waist circumference (cm)	126.2 ± 10.5	127.1 ± 11.0	0.8000 ; t=2.02619
Neck circumference (cm)	41.7 ± 3.8	41.8 ± 4.3	0.9286 ; t=2.02619
EWL (%)	-	-0.008 ± 0.04	0.3037; t=2.02809
TWL (%)	-	-0.001 ± 0.02	0.4885 ; t=2.02619
Systolic blood pressure (mmHg)	128.0 ± 12.7	123.5 ± 12.6	0.2844 ; t=2.03011
Diastolic blood pressure (mmHg)	76.2 ± 11.9	74.0 ± 10.2	0.5551 ; t=2.03011

Results are presented as mean ± SD. **F**, female; **M**, male; **Y**, yes; **N**, no; **BMI**, body mass index; **SG**, sleeve gastrectomy; **RYGB**, Roux-en-Y gastric by-pass; **BPD-DS**, biliopancreatic derivation with duodenal switch; **EWL**, excess weight loss; **TWL**, total weight loss.