

Sport	Title	Authors & Coauthors	Research Area of Focus	Journal Published	Year Published	# Authors	Sample Size Reported	Years studied	Type of Study	Country Performed	Goal of Research
Volleyball	Injury Epidemiology and Time Lost From Participation in Women's NCAA Division I Indoor Versus Beach Volleyball Players	Tristan Juhan, Ioanna K Bolla, Hyunwoo P Kang, Andrew Homere, Russ Romano, James E Thomas, Gabe P. Connors, Alessandro	Concussion Prevalence	Orthopedic Journal of Sports Med	2021	8	161 (53 beach 108 outdoor)	2015-2016	Cohort study	USA	To compare the epidemiology of injuries and time lost from participation between female NCAA Division I athletes who participate in indoor versus beach volleyball.
Basketball	Concussion incidence and Return-to-play time in National Basketball Association Players	Ajay S Padaki, Brian J Cole, Christopher S Ahmad	Concussion Prevalence	American Journal of Sports Medicine	2016	3	315	2006-2010; 2011-2014	Descriptive Epidemiological Study	USA	The incidence and return-to-play statistics were generated by synthesizing information from publicly available records
Equestrian	Horse-related incidents and factors for predicting injuries to the head	Lauren Meredith, Robert Ekman, Robert Thomson	Concussion Prevalence	BMJ Open Sport & Exercise Science	2018	3	7815	2001-2016	Retrospective Review	Sweden	Investigate the association between incidence type and head injury in equestrian-related sports.
Equestrian	Concussion History and Knowledge Base in Competitive Equestrian Athletes	Heather N Kuhl, David Ritchie, Angela C Tavera-Dick, Katie A Hoelling, Stephen A Russo	Concussion Prevalence	Sage Journals	2013	6	94	2010	Cross-sectional level 2	USA	Concussions per rider
Equestrian	Adult sports-related traumatic brain injury in United States trauma centers	Ethan A Winkler, John K Yue, John F Burke, Andrew K Chan, Sanjay S Dhali, Mitchell S Berger, Geoffrey T Nathan A Jacobson, David Buzas, Lawrence G. Morawa	Concussion Prevalence via Hospital/Urgent Care Records	Neurosurg Focus	2016	8	4788	2003-2012	Retrospective	USA	Utilizing the National Sample Program of the National Trauma Data Bank (NTDB), the authors retrospectively analyzed sports-related TBI data from adults (age ≥ 18 years) Over an 11-year span from January 2002 to December 2012, the authors reviewed the concussions sustained by athletes aged 5 to 13 years while playing football, as evaluated in emergency departments (EDs) in the United States and compared to the National Participants were put into two groups, (<14 years) or late (> or equal to 14 years) specialization, in addition to whether they sustained a concussion during competitive play, and risk factors related to horse riding, and to define groups of safer riders and those at greater risk
Football	Concussions From Youth Football: Results From NEISS Hospitals Over an 11-Year Time Frame, 2002-2012	Nathan A Jacobson, David Buzas, Lawrence G. Morawa	Concussion Prevalence via Hospital/Urgent Care Records	Orthopedic J Sports Med	2013	3	2028	2002-2012	Descriptive Epidemiological Study	USA	Utilizing the National Sample Program of the National Trauma Data Bank (NTDB), the authors retrospectively analyzed sports-related TBI data from adults (age ≥ 18 years) Over an 11-year span from January 2002 to December 2012, the authors reviewed the concussions sustained by athletes aged 5 to 13 years while playing football, as evaluated in emergency departments (EDs) in the United States and compared to the National Participants were put into two groups, (<14 years) or late (> or equal to 14 years) specialization, in addition to whether they sustained a concussion during competitive play, and risk factors related to horse riding, and to define groups of safer riders and those at greater risk
Gymnastics	Health outcomes among former female collegiate gymnasts: the influence of sport specialization, concussion, and disordered eating	Emily Sweeney, David R Howell, Corinne N Seehusen, David Tilley, Ellen Casey	Concussion Prevalence	The Physician and Sports Medicine	2020	5	437	Not Reported	Survey	USA	To analyze head/face injury diagnoses and to identify the mechanisms associated with such injuries.
Horse Riding	Protective and risk factors in amateur equestrians and description of injury patterns: retrospective data analysis and case-control survey	Rebecca M Hasler, Lena Gyselar, Lorin Benneker, Luca Martini, Andreas Schöttau, Heinz Zimmermann, and Aristomenis K Exadaktylos	Concussion Prevalence via Hospital/Urgent Care Records	Journal of Trauma Management & Outcomes	2011	7	365	2000-2006	Retrospective / case control survey	Switzerland	This study aimed to identify situational factors associated with concussion in elite Australian male and female cricket
Luge/Skeleton	Concussions in Sliding Sports and the Unrecognized "Sled Head": A Systematic Review	Melissa D McCradden, Michael D Cusimano	Concussion Prevalence Systematic Review	Frontiers in Neurology	2018	2	40	1946-2017	Systematic Review	Canada	To summarize our knowledge of the prevalence of concussion and related symptoms in sliding sports;
Snowboarding/skiing	Head Injuries in Skiers and Snowboarders in British Columbia	S Hentschel, W Hader, M Boyd	Concussion Prevalence via Hospital/Urgent Care Records	The Canadian Journal of Neurological Sciences	2000	3	54	1992-1997	Retrospective	Canada	The British Columbia Trauma Registry was searched and analyzed for patients incurring head injuries and length of hospital stay while skiing or snowboarding from January 1992 to December 1997.
Snowboarding/skiing	Skiing, Snowboarding, and Sledding Injuries in a Northwestern State	Carol S Federluk, Jamie L Schlueter, Annette L Adams	Concussion Prevalence via Hospital/Urgent Care Records	Wilderness and Environmental Medicine	2002	3	132	1992-1999	Retrospective	USA	State trauma registry data from the 1992-93 through 1999-00 ski seasons on all snow sports participants transported to tertiary trauma hospitals in Oregon were analyzed. A study to examine the incidence and characteristics of concussions among Canadian university athletes during 1 full year of football and soccer participation.
Soccer/Football	Concussions among university football and soccer players	J Scott Delaney, Vincent J Lacroix, Suzanne Leclerc, Karen M Johnston	Concussion Prevalence	Clinical Journal of Sports Med	2002	4	529	1999-2000	Retrospective survey	Canada	A study to examine the incidence and characteristics of concussions among Canadian university athletes during 1 full year of football and soccer participation.
Softball	A Retrospective Analysis of Softball-Related Head and Facial Injuries Treated in United States Emergency Departments, 2013-2017. Situational factors associated with concussion in cricket identified from video analysis	John S. Strickland, Marie Crandall, and Grant R. Bevil Anna E Saw, David J Howard, Alex Kountouris, Andrew S McIntosh, John W Orchard, Richard Saw and Thomas Hill	Concussion Prevalence	Orthopedic Journal of Sports Med	2019	3	3324 injuries	2013-2017	Descriptive Epidemiological Study	USA	To analyze head/face injury diagnoses and to identify the mechanisms associated with such injuries.
Cricket	A study to examine the incidence and characteristics of concussions among Canadian university athletes during 1 full year of football and soccer participation.	Anna E Saw, David J Howard, Alex Kountouris, Andrew S McIntosh, John W Orchard, Richard Saw and Thomas Hill	Concussion Prevalence	Journal of Concussion	2021	5	485	2015-2019	retrospective case series analysis method	Australia	This study aimed to identify situational factors associated with concussion in elite Australian male and female cricket
Waterpolo	The Epidemiology of Sports-Related Head Injury and Concussion in Water Polo	Robert S. Blumenfeld, Jessica C. Winsell, James W. Hicks and Steven L. Small	Concussion Prevalence	Frontiers in Neurology	2016	4	1470	2015	Retrospective survey	USA	An electronic survey to the 44,000+ members of USA Water Polo, asking questions about concussions, head impacts, and symptoms commonly associated with prior concussion.
Snowboarding/skiing	Emergency department reported head injuries from skiing and snowboarding among children and adolescents, 1996-2010	Janessa M Graves, Jennifer M Whitehill, Joshua O Stream, Monica S Vavilala, Frederick P Rivara	Concussion Prevalence via Hospital/Urgent Care Records	BMJ Publishing Group	2014	5	78538	1996-2010	Retrospective	USA	To evaluate the incidence of snow-sports-related head injuries among children and adolescents reported to emergency departments (EDs), and to examine the trend from 1996 to 2010 in ED visits for snow-sports-related injuries.
Snowboarding/Skiing	Concussion Among Youth Skiers and Snowboarders: A Review of the National Trauma Data Bank From 2009 to 2010	Kelly R Bergmann, Andrew Flood, Nathaniel S Kreykes, Anupam B Kharbanda	Concussion Prevalence	Pediatric Emergency Care	2015	4	1001	2009-2010	Retrospective / cross sectional	USA	Subjects 15 years or younger with a ski- or snowboard-related injury were studied using data from the National Trauma Data Bank to determine concussion risk between helmet and non-helmet user
Baseball	Concussion rates and effects on player performance in Major League Baseball players	Vani J Sabesan, Beau Prey, Ryan Smith, Daniel J Lombardo, Wilfredo J Borroto, and James D Whaley	Concussion Prevalence	J Sports Med	2018	5	112	2005-2016	Descriptive Epidemiological Study	USA	Relative risk of concussion per 100,000 athlete exposures, independent t-test compare averages between groups. Paired sample t-tests were performed to compare pre- and postinjury statistics
Field Hockey	Concussion in field hockey: a retrospective analysis into the incidence rates, mechanisms, symptoms and recovery of concussive injuries sustained by elite field hockey players	Michael Rossiter and Michael Challis	Concussion Prevalence	BMJ Open Sport and Exercise Science	2017	2	28	Not Reported	Retrospective	UK	To identify the incidence and mechanisms of concussion in elite Field Hockey in different age groups and also the post concussion symptoms and recovery times.
Hockey	Detailed description of Division I ice hockey concussions: Findings from the NCAA and Department of Defense CARE Consortium	Kathryn L. Van Pelt, Jaclyn B. Caccese, James T. Eckner, Margot Pulkusian, M. Alison Brooks, Kenneth L. Cameron, Megan N. Houston, Matthew A. Posner, Jonathan C. Mischa Fausen, Gueten Nyberg, Cecilia Tegner, Yelverton Tegner	Concussion Impact Factors	Journal of Sport Health Science	2021	15	332	2014-2017	Retrospective cohort study	USA	elucidate risk factors, specific mechanisms, and clinical presentations of concussion in men's and women's ice hockey.
Hockey	Concussion in Ice Hockey—A Cohort Study Across 29 Seasons	Matthew A. Posner, Jonathan C. Mischa Fausen, Gueten Nyberg, Cecilia Tegner, Yelverton Tegner	Concussion Prevalence	Clinical Journal of Sports Med	2017	4	267	1984-2013	Cohort Study	Sweden	Incidence rate ratio for concussion and rehabilitation periods due to concussion were calculated and analyzed.
Skiing, Freestyle Skiing, snowboarding, motocross, dinghy sailing, sliding sports (luge, bobsleigh, skeleton), skating sports (figure, short track, speed)	The Incidence of Pediatric and Adolescent Concussion in Action Sports: A Systematic Review and Meta-Analysis	Francesco Feletti, and Matteo Bonato	Concussion Prevalence Systematic Review	Int J environmental ResPublic health	2020	2	14 studies	1980-2020	Systematic/Meta	USA	This was a systematic review and meta-analysis of the incidence of concussion risk in youth athletes involved in action sports (AS)
Soccer	Head and neck injuries in professional soccer	Michael Nilsson, Martin Hägglund, Jan Ekstrand, Markus Waldén	Concussion Prevalence	Clinical Journal of Sports Med	2013	4	136 injuries	2001-2010	Prospective cohort	Sweden	Injury rate (number of time loss injuries per 1000 hours)
Luge	Injuries in the Sport of Luge: Epidemiology and Analysis	R S Cummings Jr 1, A T Shurtland, J A Prodoehl, K Moody, H H Sheik	Concussion Prevalence	The American Journal of Sports Medicine	1997	5	1034	1985-1992	Retrospective	USA	Between years 1985 -1992, data obtained from the athlete injury and liveness report forms at the US Training Center Sports Medicine Clinic in Lake Placid, New York
football, cheerleading, wrestling, volleyball, baseball, softball, boys' and girls' basketball, boys' and girls' soccer, and boys' and girls' track	Incidence and risk factors for concussion in high school athletes, North Carolina, 1996-1999	Mark R Schulz, Stephen W Marshall, Frederick O Mueller, Jingzhen Yang, Nancy L Weaver, William D Kalsbeek, J Michael Bowling	Concussion Prevalence	American Journal of Epid	2004	7	15802	1996-1990	Prospective cohort	USA	Quantify risk factors for sports concussions
Baseball	Mild traumatic brain injury in major and minor league baseball players	Gary A. Green, Keshia M. Pollack, John D'Angelo, Mark S. Schickendantz, Roger Caplinger, Kathleen Weber, Alex Valadka, Thomas W. McAllister, Randall W. Dick, Bert Mandelbaum, Frank C. Curriero,	Concussion Prevalence	American Journal of Sports Medicine	2015	11	307	2011-2012	Descriptive Epidemiology Study	USA	association between MTBI and return to play using several different measures.
Soccer	Mechanisms of head injuries in elite football	T E Andersen, A Arnason, L Engebretsen, R Bahr	Concussion Prevalence	British Journal of Sports Medicine	2004	3	192 incidents	1999-2000	Prospective cohort	Norway	the aim of this study was to describe, using video analysis, the mechanisms of head injuries and of incidents with a high risk of head injury in elite football within Norway and Iceland.
Soccer	Concussion incidence and recovery in Swedish elite soccer—Prolonged recovery in female players	Fredrik Vedung, Sofie Hänni, Yelverton Tegner, Jakob Johansson, Niklas Marklund	Concussion Prevalence	Scandinavian Journal of Medicine & Science in Sports	2020	5	959	2017	Descriptive epidemiological study.	Sweden	In the 1st and 2nd soccer leagues for men and women, a Sport Concussion Assessment Tool (SCAT)-based questionnaire study was performed at seven professional and from
Soccer	Head injuries in the female football player: incidence, mechanisms, risk factors and management	Jiri Dvorak, Paul McCrory, Donald T Kirkendall	Concussion Prevalence	British Journal of Sports Med	2007	2		1998-2004	Descriptive Epidemiological Study	USA	Exploring why females experience a higher rate of head injury, a different spectrum of injury and different mechanism of injury compared with male footballer
Soccer	Football injuries during the 2014 FIFA World Cup	Astrid Junge, Jiri Dvořák	Concussion Prevalence	British Journal of Sports Medicine	2015	2	736	2014	Retrospective	Switzerland	To analyse the incidence and characteristics of match injuries incurred during the 2014 FIFA World Cup in comparison to previous FIFA World Cups.
Soccer	Time Trends of Head Injuries Over Multiple Seasons in Professional Male Football (Soccer)	Florian Beaudouin, Karen Aus der Fütten, Tobias Tröb, Claus Reinsberger, Tim Meyer	Concussion Prevalence	Sports Med International Open	2019	5	238	2006/07/ 2016-2017	Prospective injury surveillance study	Germany	The present study aimed to investigate time trends of head injuries and their injury mechanisms since a rule change as monitoring may help to identify causes of head injuries and may advance head injury prevention efforts
Snowboarding/skiing	Head injuries among FIS World Cup alpine and freestyle skiers and snowboarders: a 7-year cohort study	Sophie E Steenstrup, Tone Bere, Roald Bahr	Concussion Prevalence	BJSM	2013	3	2080	2006-2013	Retrospective	Norway	retrospective interviews with FIS WC athletes at the end of seven consecutive seasons (2006-2013) to register injuries sustained during the competitive season to calculate the exposure of head and face injuries
Cricket	Incidence of Concussion and Head Impacts in Australian Elite-Level Male and Female Cricketers After Head Impact Protocol Modifications	Thomas Hill, John Orchard, and Alex Kountouris	Concussion Prevalence	Sports Health	2019	3	555	2015-2016 and 2016-2017	Descriptive Epidemiological Study	Australia	subsequent analysis of head impacts and concussions in elite-level male and female cricketers in Australia over 2 seasons
Cheerleading	Catastrophic High School and Collegiate Cheerleading Injuries in the United States: An Examination of the 2006-2007 Basket Toss Rule Change	Rebecca K Yau, Savannah G Dennis, Barry P Boden, Robert C Cantu, James A Lord, Kristen L Kucera	Concussion Prevalence	Sports Health	2019	6	Basket tossing injuries n=18, subjects= 54	2002- 2017	Case Series	USA	To describe catastrophic cheerleading injuries among high school and collegiate-level participants in the United States and to explore whether the 2006-2007 basket toss rule change was effective at reducing the number of catastrophic injuries
Football, soccer, volleyball, basketball, wrestling, baseball, softball	Concussions Among United States High School and Collegiate Athletes	Luke M Gessel, Sarah K Fields, Christy L Collins, Randall W Dick and R. Dawn Comstock	Concussion Prevalence	Journal of Athletic Training	2007	5	4431 injuries	2005-2006	Descriptive Epidemiological Study	USA	To investigate the epidemiology of concussions in a nationally representative sample of high school athletes and to compare rates of concussion among high school and collegiate athletes.
Hockey	A prospective study of physician-observed concussions during junior ice hockey: implications for incidence rates	Paul Sean Echlin, Charles H Tator, Michael D Cusimano, Robert C Cantu, Jack E Taunton, Ross E G Upshur, Craig R Hall, Andrew M Johnson, Lorie A Forwell, Elaine N	Concussion Prevalence	Neurosurg Focus	2010	10	67	2009-2010	Prospective cohort	Canada	The objective of this study was to measure the incidence of concussion (scaled relative to number of athlete exposures) and recurrent concussion within 2 teams of fourth-tier junior ice hockey players (16-21 years old) during 1

Volleyball	A Comparison of Women's Collegiate and Girls' High School Volleyball Injury Data Collected Prospectively Over a 4-Year Period	Jonathan C. Reeser, Andrew Gregory, Richard L. Berg, R. Dawn Comstock,	Concussion Prevalence	American Ortho Society of Sports Med	2015	4 729 High School Injuries/ 1380 College Injuries	2005-2009	Retrospective clinical review	USA	Analyze injury rates from collected data within the National Collegiate Athletic Association's Injury Surveillance System (NCAA ISS) and the High School Reporting Injuries Online (HS RIO).	
Basketball	Epidemiology of Secondary School Boys' and Girls' Basketball Injuries: National Athletic Treatment, Injury and Outcomes Network	Alex N. Allen, Erin B. Wasserman, Richele M. Williams, Janet E. Simon, Thomas P. Dompier, Zachary Y. Kerr, Alison R. Snyder Vaier, PhD	Concussion Prevalence	Journal of Athletic Training	2019	7	167	2011-2014	Descriptive Epidemiological Study	USA	To describe the epidemiology of time-loss (TL) and NTL injuries sustained by secondary school boys' and girls' basketball athletes.
football, cheerleading, wrestling, diving/swim, baseball, softball, boys' and girls' basketball, boys' and girls' tennis and boys' and girls' track, golf, crew, girls soccer	Epidemiology of Sport-Related Concussions in High School Athletes: National Athletic Treatment, Injury and Outcomes Network (NATION), 2011-2012 Through 2013-2014	Kathryn L. O'Connor, Melissa M. Baker, Sara L. Dalton, Thomas P. Dompier, Steven P. Broglio, and Zachary Y. Kerr	Concussion Prevalence	Journal of Athletic Training	2017	6	2004 injuries	2011-2014	Descriptive Epidemiological Study	USA	To describe the epidemiology of sport-related concussion (SRC) in 27 high school sports during the 2011-2012 through 2013-2014 academic years.
Basketball	Effect of Mouthguard's on dental injuries and concussions in college basketball	Cynthia R. Labella, Bryan W. Smith, Asger Sigurdsson	Concussion Prevalence	Official Journal of the American College of Sports Medicine	2001	3	70,936	1999-2000	Survey	USA	During the 1999 to 2000 basketball season, athletic trainers from 50 men's Division I college basketball programs used an Internet Web site to submit weekly reports of the. To determine the number of concussions that occurred on 3 collegiate Division I military academy football teams prior to and following recent changes in the NCAA concussion management policy.
Football	Reported Concussion Rates for Three Division I Football Programs: An Evaluation of the New NCAA Concussion Policy	Kelly G. Kilcoyne, Jonathan F. Dickens, Steven J. Svoboda, Brett D. Owens, Kenneth L. Cameron/Robert T. Sullivan, John-Paul Rue	Concussion Prevalence	American Ortho Society of Sports Med	2013	7	65 reported Concussions	2009-2011	Descriptive Epidemiological Study	USA	To determine the number of concussions that occurred on 3 collegiate Division I military academy football teams prior to and following recent changes in the NCAA concussion management policy.
boys' baseball, basketball, football, soccer, track, and wrestling; and girls' basketball, cheerleading, soccer, softball, track, and volleyball	Concussion Rates in U.S. Middle School Athletes, 2015-2016 School Year	Zachary Y. Kerr, Nelson Cortes, Amanda M. Caswell, Justin P. Ambegankar, Kaitlin Romm Hallsmith, Frederick Milbert, Shane V. Caswell	Concussion Prevalence	American Journal of Sports med	2017	7	2,622	2015-2016	Descriptive Epidemiological Study	USA	athletic exposure (AE) data in school-sanctioned games and practices in boys' baseball, basketball, football, soccer, track, and wrestling; and girls' basketball, cheerleading, soccer, softball, track, and volleyball.
Gymnastics/Cheer	Epidemiology of Cheerleading Stunt-Related Injuries in the United States	Brenda J. Shields, Soledad A. Fernandez, Gary A. Smith	Concussion Prevalence	Journal of Athletic Training	2009	3	338	2006-2007	Prospective injury surveillance study	USA	The numbers, types, and rates of cheerleading stunt-related injuries during a 1-year period (2006-2007) are reported. Data retrieved from RIO Reporting Information System (RIS) surveillance system.
Field Hockey	Head, Face, and Eye Injuries in Scholastic and Collegiate Lacrosse: A 4-Year Prospective Study	Andrew E. Lincoln, Richard Y. Hinton, Jon L. Almquist, Sean L. Lager, Randall W. Dick,	Concussion Prevalence	The American Journal of Sports Medicine	2007	5	1,156,573	2000-2003	Descriptive Epidemiological Study	USA	Injuries were analyzed in terms of (1) session type (practice or game), (2) nature of injury, (3) anatomical area injured, (4) mechanism of injury, (5) player activity at time of injury, and (6) most common activities for concussions.
Basketball	Epidemiology of Injuries in National Collegiate Athletic Association Men's Basketball: 2014-2015 Through 2018-2019	Sarah N. Morris, Avinash Chandran, Brandon B. Lempe, Adrian J. Boltz, Hannah J. Robison, Christy L. Collins	Concussion Prevalence	Journal of Athletic Training	2021	6	3481 injuries total	2014-2019	Descriptive Epidemiological Study	USA	Injury counts and rates per 1000 AEs were examined by event type (practice, competition), competition level (Division I, Division II, Division III, soccer, baseball).
Football, soccer, volleyball, basketball, wrestling, baseball, softball	National High School Athlete Concussion Rates From 2005-2006 to 2011-2012	Joseph A. Rosenthal, Randi E. Foraker, Christy L. Collins, R. Dawn Comstock	Concussion Prevalence	American Ortho Society of Sports Med	2014	5	2516 Concussions reported	2005-2012	Descriptive Epidemiological Study	USA	During the academic years from 2005-2006 through 2011-2012, all injuries (including concussions) and exposure data from high school athletes participating in 9 sports were prospectively gathered by the HS RIO injury surveillance system, which has been
football, cheerleading, wrestling, volleyball, baseball, softball, boys' and girls' basketball, boys' and girls' soccer, and boys' and girls' track	Epidemiology of concussions among United States high school athletes in 20 sports	Malika Marar, Natalia M. McIlvain, Sarah K. Fields, R. Dawn Comstock	Concussion Prevalence	American Journal of Sports Medicine	2012	4	14,635	2008-2010	Retrospective cohort study	USA	In the 20 sports studied over the course of the 2008-2010 school years
Hockey	Incidence of Sports-Related Concussion Among NCAA Women's Ice Hockey Athletes	Emily M. Brook, Emily Krostus, Caroline H. Hu, Marissa Giedman, Jamie E. Collins, Elizabeth G. Matzkin	Concussion Prevalence	Orthopedic Journal of Sports Medicine	2017	6	459	2014-2015	Descriptive Epidemiological Study	USA	To determine the incidence of sports-related concussion (SRC) in National Collegiate Athletic Association (NCAA) women's ice hockey athletes.
Hockey	Comparison of Concussion Rates Between NCAA Division I and Division III Men's and Women's Ice Hockey Players	John M. Rosene, Bryan Rakisins, Brie Silva, Tyler Woefel, Paul S. Visich, Thomas P. Dompier, Zachary Y. Kerr	Concussion Prevalence	American Journal of Sports Medicine	2017	7	415	2009-2015	Descriptive Epidemiological Study	USA	To compare the epidemiologic patterns of concussion in National Collegiate Athletic Association (NCAA) ice hockey by sex and division.
Volleyball	Descriptive Epidemiology of Injuries Sustained in National Collegiate Athletic Association Men's and Women's Volleyball, 2013-2014 to 2014-2015	Christine M. Baugh, Gil S. Weintraub, Andrew J. Gregory, Arstarque Djoko, Thomas P. Dompier, Zachary Y. Kerr	Concussion Prevalence	Sports Health	2018	6	18844	2014-2015	Descriptive Epidemiological Study	USA	To examine injury epidemiology in NCAA men's and women's volleyball athletes.
Softball	Descriptive Epidemiology of Collegiate Women's Softball Injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 Through 2003-2004	Stephen W. Marshall, Karrie L. Hamstra-Wright, Randall Dick, Katie A. Grove, and Julie Agel	Concussion Prevalence	Journal of Athletic Training	2007	5	5336 injuries	1988-1989-2003-2004	Descriptive Epidemiological Study	USA	The NCAA Injury Surveillance System has tracked injuries in all divisions of NCAA softball
Basketball	Descriptive Epidemiology of Collegiate Women's Basketball Injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 Through 2003-2004	Julie Agel, David E. Olson, Randall Dick, Elizabeth A. Arendt, Stephen W. Marshall, and Robby S. Sikka	Concussion Prevalence	Journal of Athletic Training	2007	7	3556 Injuries	1988-2004	Retrospective	USA	To review 16 years of National Collegiate Athletic Association (NCAA) injury surveillance data for women's basketball and to identify potential areas for injury prevention initiatives.
Field Hockey	The First Decade of Web-Based Sports Injury Surveillance: Descriptive Epidemiology of Injuries in US High School Girls' Field Hockey (2008-2009 Through 2013-2014) and National Collegiate Athletic Association Women's Field Hockey (2014-2015 Through 2018-2019)	Robert C. Lynam, Elizabeth C. Gardner, Jordan Paolucci, Dustin W. Currie, Sarah B. Knowles, Lauren A. Pierpoint, Erin B. Wasserman, Thomas P. Dompier, R. Dawn Comstock, P. Stephen W. Marshall, Zachary Y. Kerr, Avinash Chandran, Aliza K. Nedimyer, Alan Arakkal, Lauren A. Pierpoint, Scott L. Zuckerman	Concussion Prevalence	The American Journal of Sports Medicine	2015	1	150	2004-2009	Descriptive Epidemiological Study	USA	To describe the epidemiology of injuries sustained in high school girls' field hockey in the 2008-2009 through 2013-2014 academic years and collegiate women's field hockey in the 2004-2005 through 2013-2014 academic years using Web-based sports injury surveillance rates per 1000 AEs with 95% confidence intervals (CIs) practice vs game
boys' football, wrestling, soccer, basketball, and baseball and girls' volleyball, soccer, basketball, and softball	Concussion Incidence and Trends in 20 High School Sports	Zachary Y. Kerr, Avinash Chandran, Aliza K. Nedimyer, Alan Arakkal, Lauren A. Pierpoint, Scott L. Zuckerman	Concussion Prevalence	American Academy of Pediatrics	2019	6	6079 concussions	2013-2014 & 2017-2018	Descriptive Epidemiological Study	USA	Linear regression
Football	Incidence of Concussion During Practice and Games in Youth, High School, and Collegiate American Football Players	Thomas P. Dompier, Zachary Y. Kerr, Stephen W. Marshall, Brian Hainline, Erin M. Snook, Ross Hayden, Janet E. Simon	Concussion Prevalence	JAMA Pediatrics	2015	7	20354	2012-2013	Descriptive Epidemiological Study	USA	To examine and compare sport-related concussion outcomes (symptoms and return to play) in youth, high school, and collegiate football athletes.
Softball	Epidemiology of Injuries in National Collegiate Athletic Association Women's Softball: 2014-2015 Through 2018-2019	Kevin L. Veillard, Adrian J. Boltz, Hannah J. Robison, Sarah N. Morris, Christy L. Collins, Avinash Chandran	Concussion Prevalence	Journal of Athletic Training	2021	6	1511 injuries	2014-2019	Descriptive Epidemiological Study	USA	Exposure and injury data collected during competitive seasons in the NCAA Injury Surveillance Program during 2014-2015 through 2018-2019 (5 years) academic years were analyzed. Injury counts, rates, and
boys/girls—soccer and basketball, girls—field hockey and lacrosse, boys/girls—swim & dive, track & field, and cross country, boys—baseball	Concussion Incidence and Trends in 20 High School Sports	Zachary Y. Kerr, Avinash Chandran, Aliza K. Nedimyer, Alan Arakkal, Lauren A. Pierpoint, Scott L. Zuckerman	Concussion Prevalence	Pediatrics	2019	6	9542	2013-2018	Descriptive Epidemiological Study	USA	2013-2014- 2017/2018

Exposure Defined	Athletic Exposure Statistical Computation Reported	Statistical Analyses	Return to play time average (days)	Average concussions per athlete/ season	Incidence Rate Reporting	Stat Sig Reported
The injury rate was expressed per 1000 hours played	\sum all injuries / \sum all athlete-exposures	Regression Analysis of AE and Injury Rate	Not Reported	On average, there was a 6.1 concussion rate per 1000 hours	6.5% BEACH and 7.5% INDOOR	P < 0.05
Sum of concussive injuries per Season	-	T test on reported time loss following a concussion before and after implementation of concussion protocol	5 games missed	14.9 concussions per season	Not Reported	P < 0.05
Sum of concussive injuries	Percentage of head injuries	Generalized Linear Model	Not Reported	In riders who suffered a fall from horseback that resulted in concussion (n=461), 33.4% contacted a hard floor surface such as asphalt; 25.5% landed on grass or dirt surface and 20.8% landed on indoor riding arena floor surfacing. Of these 461 riders, 88% reported they were wearing a helmet	68.3% of head injuries occurred from horse handling, while mounting or dismounting only accounted for 1.5%	P < 0.05
Sum of concussive injuries	Percentage of head injuries	Measures of central tendency were utilized to evaluate response patterns	Over 51% (n = 19) took less than 5 days off, 30% (n = 11) took between 14 and 21 days and 8% (n = 3) took more than 30 days off	Not Reported	45% sustained a concussion	Not Reported
Sum of concussive injuries	Percentage of head injuries	Multivariable regression analysis	Not Reported	Not Reported	45.2 TBI came from equestrian, 2.8% mortality rate in ICU	$\alpha = 0.05$ yielding p < 0.01
Sum of concussive injuries	Percentage of head injuries	Nationals concussion estimated risk by (%), year were quantified through spearman correlation coefficients	Not Reported	correlation of $r = 0.97$. The total number of players experiencing a concussion compared with the total number of players experiencing LoC over the 11-year time period had a correlation	National estimates were calculated to be 49,185 concussions occurring in youth football players 5 to 13 years of age over an 11-year time frame	P < 0.05, r > 0.618
Sum of concussive injuries	Percentage of head injuries	Multivariate regression analysis	Eating disorder had an increased risk on concussion time-loss had an increased risk of time-loss injury (78% vs 65%; n=0.004)	Not Reported	42% reported concussion history	P < 0.05
Sum of concussive injuries	Percentage of head injuries	Multiple logistic regression was performed, and combined risk factors were calculated using inference trees	Not reported	127 reported concussions	24% sustained a concussion	P < 0.05
Sum of concussive injuries	Percentage of head injuries	Systematic Review	Not Reported	Not Reported	Salt Lake City Olympics 2002- 19.6% of all patients (bobsled, 10.1%; luge, 8.7%; skeleton, 0.7%) 2010 Olympics (19), 0.07 of all registered athletes reported concussions, which was twice as high as the concussion incidence in the 2008 Summer Olympic Games. 2012 Youth Games - 0.07	Not Reported
Sum of concussive injuries	Percentage of head injuries	Descriptive Statistics	Both the hospital stay and intensive care unit stays were longer for the snowboarders with an average stay of 20.4 days on the ward and 11.8 days in the ward	Head injury was 0.005 per 1000 for skiers, 0.004 per 1000 for snowboarders, concussion was in 60% of skiers and 21% in snowboarders	At the 2014 Sochi Olympics (21), 11 concussions 11-81 concussions per head injuries	Not Reported
Sum of concussive injuries	Percentage of head injuries	Regression Analysis of Odds Ratios	Not Reported	Concussions accounted for 39% of all injuries	Sledding concussions accounted for 55% of injuries while Skiing accounted for 39%, and snowboarding with the lowest reported percentage of 27%	Not Reported
Sum of concussive injuries	Percentage of head injuries	Logistic regression model using BIC	injuries leading to loss of time was 3.7 (95% CI 2.7 to 4.7) per 1000 player hours (men: 3.5, 95% CI 2.4 to 4.6; women: 4.1, 95% CI 2.1 to 6.1).	70.4% of the football players and 62.7% of the soccer players had experienced symptoms of a concussion during the previous year. Only 23.4% of the concussed football players and 19.8% of the concussed soccer players realized they had suffered a concussion. More than one concussion was experienced by 44.6% of the concussed football players. Concussions accounted for 17.7% of all injuries	head/neck injuries was 12.5 (95% CI 10.9 to 14.1) per 1000 player hours (men: 12.8, 95% CI 11.0 to 14.7; women: 11.5, 95% CI 8.4 to 14.6)	P < 0.05
Sum of concussive injuries	Percentage of head injuries	Regression Analysis of head impacts	Not Reported	Median number of reported symptoms on the day of injury for concussed players was 7 out of 24	84 reported female concussions and 16 reported male concussions	P < 0.05
risk of one head impact per every x amount of balls	Not Reported	Regression Analysis of head impacts	Median number of reported symptoms on the day of injury for concussed players was 7 out of 24	Concussion was diagnosed in 35 cases (18%, 95% CI 13-24%)	Risk of one head impact every 7,736 balls or approximately 13 days play	P < 0.002
Percentage of head injuries	-	Regression Analysis of History of Concussion	Not Reported	Men reporting concussion sustained an average of 2.20 \pm 0.12 concussions. Women reporting concussion sustained an average of 2.06 \pm 0.08 concussions.	In this group of 534, respondents reported sustaining an average of 2.14, SE (\pm) = 0.07 concussions	P < 0.05
per 'x' amount of resort visits	the number of head injuries divided by snow sports participation estimates derived from annual NSAA surveys	Regression Analysis of TBI visits	Not Reported	An estimated number of 78 538 (95% CI 66 350 to 90 727) snow sports-related head injuries among children and adolescents, 77.2% were traumatic brain injuries	The annual average incidence rate of TBI was 2.24 per 10 000 resort visits for children (4-12) compared with 3.13 per 10 000 visits for adolescents (13-17).	P < 0.05
Odds Ratio	-	Multivariate regression analysis and odds ratio	Not Reported	Not Reported	Snowboarders had a greater likelihood of concussion compared to skiers (estimated- β , 2.1; 95% CI, 1.48-2.85)	P < 0.05
Not Reported	calculate the relative risk of concussion at the different positions. It was assumed that there were four catchers, eight infielders, eight outfielders, and 20 pitchers on each of the 30 team's rosters	Independent sample T-tests, paired sample	31.3 days following concussions	Not Reported	Field players = 3 per 100,000 AE Catchers 9 per 100,000 AE	P < 0.05
Not Reported	\sum all injuries / \sum all athlete-exposures	Regression Analysis of AE	Not Reported	Not Reported	U16 boys- .63 per 1000hrs U18 boys- .69 per 1000 hrs U21 Men- 1.29 per 1000hrs 21 women - 3.03 per 1000hrs	Not Reported
Not Reported	Not Reported	Regression Analysis of Odds Ratios	Not Reported	Between 2014 and 2017, there were 332 athletes who participated in ice hockey; 47 sustained a concussion while enrolled in the CARE Consortium study	A prior concussion increased the odds of an incident concussion 2-fold (OR = 2.03; 95%CI: 1.04-4.08)	P < 0.05
Not Reported	Exposure = (Number of team-matches played * number of players per team) / duration of match / 60	Injury rate ratios (IRRs), χ^2 statistics	Not Reported	162 concussions	incidence rate ratio was 1.06 (confidence interval, 1.03-1.10)	P < 0.05
Not Reported	Not Reported	The incidence of concussion was reported in diverse ways, the two most common denominators being days of athletic exposure (DAEs) and player-year (PY)	Not Reported	alpine skiing (n = 10), freestyle skiing (cross, halfpipe, slopestyle, n = 2), snowboarding (halfpipe and slopestyle, n = 12), off-road motorcycle/motocross (n = 2), dinghy sailing (Europe and Laser classes, n = 1), sliding sports (bobsleigh, luge and skeleton, n = 2), skating sports (figure, short track and speed, n = 1)	overall incidence of concussion in children and adolescents aged <18 years involved in outdoor sports of 0.33/1000 DAEs	P < 0.05
Not Reported	Not Reported	Simple and multiple risk factor analyses were evaluated using Cox regression for observational Injury Risk Ratio	10 days on average	head and neck injury during match play compared with training had a 78-fold higher rate of concussions from 2001 to 2010	0.06 concussions per 1000 AE hours.	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
Injuries per 'x' amount of runs	Not Reported	The risk of sustaining any injury causing the loss of more than 1 day of practice was 0.04 per person per year, and 2% of all	Not Reported	10 reported concussions out of 407 injuries	2.5% sustained a concussion	Not Reported
If an athlete participated in any part of a game or practice, he or she was considered to have participated, and information was not collected on the degree of participation.	Not Reported	Generalized Poisson Regression Analysis of AE	Not Reported	overall rate of concussion was 17.15 (95% confidence interval: 13.30, 21.00) per 100,000 athlete-exposures	overall rate of concussion was 17.15 (95% confidence interval: 13.30, 21.00) per 100,000 athlete-exposures	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
Hours of game play	The average number of players per team per game on analysis of regular-season game participation via box scores that are publicly available. This average number over a season, multiplied by the number of team games at each professional level of baseball, was used as an estimate of game exposures	Regression Analysis of AE	1w 2d	Not Reported	0.42 per 1000 athletes	P < 0.05
Hours of game play	Video recordings of incidents where a player appeared to be hit in the head and the match was consequently interrupted by the referee were analysed and cross referenced with reports of acute time loss injuries from the team medical staff	Injury Rate Ratios, Poisson regression	Not Reported	incidence of 1.7 per 1000 player hours (concussion incidence 0.5 per 1000 player hours)	(22.0 per 1000 player hours in Norway and 14.8 in Iceland, p = 0.009)	P < 0.05
Hours of game play	Not Reported	Multivariate Regression	Not Reported	1.19/1000 hours of game play	females 1.22/1000 hours vs males 1.18/1000 hours	P < 0.05
Hours of game play	Not Reported	Not Reported	Not Reported	Professional MALE World Cup (6)- 51.0/1000 US (7)- 35.5 / 1000 Iceland (8)- 34.9/1000 UK (9) - 25.9/1000 Sweden (11)- 18/1000 Professional FEMALE Sweden (12)- 24/1000 Germany (13/14)- 23.3/1000	When concussions alone were considered, the incidence rate was 1.19/1000 player hours for men and 2.6/1000 player hours for women.	P < 0.05 95% CIs excluding 1.00 were considered statistically significant
Hours of game play	Match exposure was calculated by multiplying 1.5 h by 11 players and by the number of returned forms.	Reported % of injuries	Not Reported	Not Reported	Brazil(18%) - 5% concussed S. Africa- 10%-1%, Germ 2006- 9%-1% Japan 2002- 15%- 2% France 1998- 15%-1%	P < 0.05
Hours of game play	Match exposure per team was calculated using the following calculation, number of games x number of players on the field (11 players) x duration of the game in hours (1.5h per match)	Regression Analysis of injury Rate Risk (IRR)	06/07 - 5 (5) 07/08 - 17 (16) 08/09 - 12(10) 09/10 - 8 (6) 10/11 - 6(5) 11/12 - 8(9) 12/13 - 7(8) 13/14 - 4(3) 14/15 - 8(16) 15/16 - 10(18) 16/17 - 5(5) TOTAL- 8(10)	06/07- 0.41 [0.17-0.99] 07/08- 0.48 [0.22-1.07] 08/09- 0.32 [0.12-0.86] 09/10- 0.41 [0.17-0.98] 10/11- 0.58 [0.28-1.22] 11/12- 0.49 [0.22-1.10] 12/13- 0.57 [0.27-1.18] 13/14- 0.58 [0.28-1.22] 14/15- 0.49 [0.22-1.10] 15/16- 0.81 [0.44-1.51] 16/17- 0.74 [0.39-1.42] TOTAL- 0.53 [0.42-0.67]	Averaged time lost 10:10 days (median 6, range 1-47 days), head injuries IR of 1.77 (95% CI 1.56-2.01) per 1000 match hours	P < 0.05
Head injuries were classified as 'head/face' injuries and did not include neck or cervical spine injuries.	To calculate the exposure, we extracted data from the official FIS website for all WC competitions for each of the athletes interviewed	Regression Analysis of Injury Rate Ratio (100 athletes per season)	Not Reported	245 head/face injuries reported, nervous system injuries/concussions were the most common (81.6%) and 58 of these were severe (23.7%)	Highest in freestyle (5.7, 95% CI 4.5 to 6.8) and snowboard (5.0, 95% CI 4.0 to 6.0) compared with alpine skiing (3.5, 95% CI 2.7 to 4.4; RR 1.61, 95% CI 1.17 to 2.22 vs freestyle; RR 1.43, 95% CI 1.04 to 1.96 vs snowboard)	P < 0.05
head impacts/concussions per 1000 player days	Not Reported	Incidence calculation by player days	Not Reported	92 head impacts that resulted in 29 concussions in all men's teams (on 1178 team days of play) and 15 head impacts that resulted in 8 concussions in women's matches (on 367 team days of play)	The match incidence rates were 7.2 head impacts (2.3 concussions) per 1000 player days in elite male cricket and 3.7 head impacts (2.0 concussions) per 1000 player days in elite female cricket	Not Reported
catastrophic injuries were those resulting from participating in skills related to the sport, and were classified into categories of severity: serious (severe injuries without permanent functional disability), nonfatal (severe injuries with permanent functional disability), and fatal (injuries resulting in death)	Not Reported	Regression Analysis of catastrophic injuries	Not Reported	For basket tosses, injuries were most commonly sustained to the head (8/18; 47%)	cheerleaders sustained serious injuries (n = 27; 50%) during practice (n = 37; 89%) to the head (n = 28; 52%) and cervical spine (n = 17; 32%).	P < 0.05
Athlete-exposure (A-E) was defined as 1 athlete's participation in a practice or competition.	Not Reported	1000 AE. Statistical analyses included calculation of rate ratios (IRRs), proportion ratios (PRs), and χ^2 tests	More than 50% of athletes in every sport returned to play in 9 days or less	Concussions accounted for 8.9% of all injuries	Girls HS - 25/1000 Girls College - 0.45/1000 Boys HS - 18/1000 Boys College - 38/1000	All 95% confidence intervals (CIs) not containing 1.0 with P values of less than .05 were considered statistically significant.
Athlete exposures were defined within this sample as the sum of all games played, and were summed for all players.	Not Reported	The incidence of concussion was calculated as the number of observed or self-reported concussions occurring during observed games, divided by	Not Reported	incidence 21.5 concussions per 1000 athlete exposures)	A concussion was diagnosed by the physician in 19 (36.5%) of the 52 observed games	Not Reported

athlete exposure (AE) as 1 athlete's participation in 1 competition or practice without regard for the duration of participation.	Not Reported	Regression Analysis of AE	The average severity of game-related concussion was similar for both the NCAA ISS and HS RIO. 88.1% of affected collegiate athletes were absent for 3 or more days.	(NCAA ISS, 2.0 per 10,000 AEs; n = 69) (HS RIO, 0.6 per 10,000 AEs; n = 38)	Concussions were 3.4 times more likely to occur in a collegiate volleyball player than in a high school player (injury rate ratio, 3.4; 95% CI, 2.3-5.1)	P < 0.05
An athlete-exposure (AE) was defined as a single athlete participating in 1 secondary school-sanctioned practice or competition, regardless of duration, in which the athlete was exposed to the risk of injury.	\sum all injuries / \sum all athlete-exposures	Injury counts, rates, and rate ratios (IRRs) were reported with 95% confidence intervals (CIs).	Not Reported	Boys also experienced a lower concussion injury rate than girls for competitions (IRR = 0.48; 95% CI = 0.33, 0.69) but not for practices (IRR = 0.77; 95% CI = 0.52, 1.16).	Boys vs girls in competition (IRR = 0.48; 95% CI = 0.33, 0.69) Boys vs girls in practices (IRR = 0.77; 95% CI = 0.52, 1.16)	All mean differences with 95% CIs not containing 0.0 were considered statistically significant.
An athlete-exposure (AE) was defined as 1 athlete participating in 1 practice or competition. Only those athletes with playing time in a competition were included in competition exposure calculations.	Not Reported	Regression model on IRRs and IPRs were used to compare rates and proportions, respectively, in sex-comparable sports	Most athletes with SRCs reported symptom resolution within 7 days (40.7%; n = 814) or 8 to 14 days (21.7%; n = 23)	Reported Number of Concussions Foot = 1020, Wrest = 137, Gfield= 66, GGym= 8 Gvols= 74 Base= 18 Soft= 50 BBask= 92 Gask= 128 Bcrew = 0	Overall rate of concussion was 3.89 per 10000 AEs among all sports	All 95% confidence intervals (CIs) not containing 1.0 with P values of less than .05 were considered statistically significant.
An athlete exposure was defined as one athlete participating in one practice or contest where he/she was exposed to the possibility of injury.	Not Reported	Chi Test Of difference in concussion exposure from mouth guard users to nonmouth guard users	Not Reported	Not Reported	0.52 per 1000 athletes	P < 0.05
An athlete exposure was defined as 1 athlete participating in 1 practice or a game in which he was exposed to injury	\sum all injuries / \sum all athlete-exposures	Regression Analysis of Injury Rate Ratio	Not Reported	incidence rate across all 3 institutions was 2 times higher in 2010-2011 when compared with 2009-2010 (incidence rate ratio [IRR], 2.04; 95% CI, 1.20-3.55; P = 0.01)	Injury Risk 0.57 per 1000 athlete exposures in the 2009-2010 Injury Risk 1.16 per 1000 athlete exposures in the 2010-2011	P < 0.05
An athlete exposure (AE) was defined as one athlete participating in one school-sanctioned game or practice.	Not Reported	Regression Analysis of AE and Injury Rate Risk (IRR)	Not Reported	The overall concussion rate was higher in games than practices (1.15 vs 0.63/1,000 AEs, IRR=1.83, 95% CI=1.06, 3.15;	Game/Practice/Total/Game vs Practice 1000 AE's Total Male- 1.12 (0.43, 1.81) 0.80 (0.48, 1.12) 0.87 (0.58, 1.17) 1.40 (0.67, 2.92) Total Female- 1.18 (0.45, 1.91) 0.45 (0.20, 0.69)	All 95% confidence intervals (CIs) not containing 1.0.
An AE was defined as 1 cheerleader participating in 1 cheerleading event.	Not Reported	Multivariate logistic regression analysis's used, this involved the calculation of odds ratios (ORs) with 95% confidence intervals (CIs).	Not Reported	6% of reported injuries were concussions	0.04 injuries per 1000 AEs per practice and athletic events, 0.05 injuries per 1000 AEs	P ≤ .2
An AE was defined as 1 athlete participating in 1 practice or game in which the high school or college participant was exposed to the possibility of injury	Not Reported	Regression Analysis of AE	Not Reported	Concussions constituted a higher percentage of injuries among boys (73%) and men (85%) than among girls (40%) and women (41%).	High School Female/Male ratio = 0.77 Collegiate Female/Male ratio = 0.86	P < 0.05
An AE was defined as 1 athlete participating in 1 exposure event.	Not Reported	Regression Analysis of AE and Injury Rate Ratios (IRR)	Not Reported	3794 reported concussions over 5 years	5.05% sustained a concussion	95% CI excluding 1.00
An AE was defined as 1 athlete participating in 1 competition or practice.	Not Reported	Regression Analysis of AE	Not Reported	CONCUSSIONS/1000 exposures 2006 Football- .47(.41, .53) G.Soc- 0.36(.27, .47) B.Soc- .22 (.16, .31)	By 2011-2012, the overall rate of concussions had increased significantly from 0.23 (0.21, 0.25) to 0.51 (0.48, 0.55)	<0.05
An AE was 1 athlete participating in 1 athletic practice or competition.	Not Reported	Overall rates were calculated by dividing concussion incidence by AE using raw case counts, rate ratios (RRs) and injury proportion ratios (IPRs) with P values and 95% confidence.	40% returned in 3 days or less, 2% returned the same day	1288 concussions (65.6%) that occurred in competition and 647 (33.4%) that occurred during practice	concentration rate of 2.5 concussions per 10,000 AE. (IRR, 5.7; 95% CI, 5.2-6.3) girls - 1.6 per 10,000 AE boys - 3.1 per 10,000 AE	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
An AE is defined as 1 athlete competing in 1 practice or game over a given period of time	It was assumed there were approximately 6 AEs per week over a 20-week season for NCAA Division II and III players and a 24-week season for Division I players and multiplied the number of athletes participating while considering their school's division	Regression Analysis of AE	Not Reported	(n = 219, 47.7%) of respondents reported at least 1 diagnosed concussion over the duration of their entire organized ice hockey career. A total of 13.3% (n = 61) of respondents reported a diagnosed concussion during the 2014-2015 season.	The incidence rate was 1.18 (95% CI, 0.92-1.51) per 1000 athlete-exposures to a game or practice and 0.58 (95% CI, 0.45-0.74) per 1000 hours of ice time	P < 0.05
A reportable athlete-exposure (AE) was defined as 1 student-athlete participating in 1 NCAA-sanctioned practice or competition in which he or she was exposed to the possibility of athletic injury, regardless of the time associated with that participation.	Not Reported	Regression Analysis of AE and Injury Rate Ratios (IRR) and IPR	Not Reported	Division I men (0.83 per 1000 athlete-exposures [AEs]), followed by Division III women (0.78/1000 AEs), Division I women (0.65/1000 AEs), and Division III men (0.64/1000 AEs).	Division I men (0.83 per 1000 athlete-exposures [AEs]), Division III women (0.78/1000 AEs)	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
A reportable athlete-exposure (AE) was defined as 1 student-athlete participating in 1 NCAA-sanctioned practice or competition in which he or she was exposed to the possibility of athletic injury, regardless of the time associated with that participation	\sum all injuries / \sum all athlete-exposures	The injury rate was calculated as the number of injuries per 1000 AEs. Statistical analyses included calculation of injury rate ratios (IRRs)	Not Reported	concussions (men, 19.4%; women, 14.8%)	0.34 (0.07-0.61) men 1000 AE 0.39 (0.24-0.53) women	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
A reportable athlete-exposure (A-E) was defined as 1 student-athlete participating in 1 practice or competition in which he or she was exposed to the possibility of athletic injury, regardless of the time associated with that participation.	\sum all injuries / \sum all athlete-exposures	Regression Analysis of AE	Not Reported	Concussions accounted for 6% of injuries	0.25 game per 1000 AE 0.07 practice per 1000 AE	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
A reportable athlete-exposure (A-E) was defined as 1 student-athlete participating in 1 practice or competition in which he or she was exposed to the possibility of athletic injury, regardless of the time associated with that participation. Only participants with actual playing time were counted as having name exposures.	\sum all injuries / \sum all athlete-exposures	Regression Analysis of AE	Not Reported	Games = (6.5% of all injuries) (95%CI, 0.43, 0.56) Practices = (3.7%) of all injuries (95%CI, 0.13, 0.17)	Games - .50 per 1000 AE Practices- 0.15 per 1000 AE	The 95% CIs provide information about the precision of each rate and can be used to determine if 2 rates differ statistically from one another.
A reportable AE was defined as 1 student-athlete participating in 1 school-sanctioned practice or competition in which he or she was exposed to the possibility of athletic injury, regardless of the time associated with that participation.	Not Reported	Regression Analysis of AE and Injury Risk Ratios (IRR)	Nearly all of the players suffering a concussion were eventually able to return to play within the same season (92.8%). Of these, 77.3% returned in <10 days from the time of injury, while the other 22.7% returned in 11-30 days.	Concussions accounted for 42.8% of all head, face, and eye injuries reported (0.40 per 1000 AEs; 95% CI, 0.32-0.53).	high school girls reported 1.73/1000 AEs (95% CI = 1.62, 1.83) vs college which reported 5.36/1000 AEs (95% CI = 5.02, 5.69)	All mean differences with 95% CIs not containing 0.0 were considered statistically significant.
A reportable AE was defined as 1 athlete participating in 1 school-sanctioned practice or competition. Our injury of focus, concussion, was defined as (1) occurring as a result of participation in an organized practice or competition, (2) requiring medical attention by an AT or physician, and (3) being diagnosed as a concussion.	Not Reported	Regression Analysis of Injury Proportion Rates and Injury Rate Risk and AE	Not Reported	AE by 10,000 Boys Foot- 7.15 (6.73 to 7.61) Ice - 16.24 (10.34 to 25.49) Lax- 8.96 (6.84 to 11.72)	4.17 per 10000 AEs (95% CI: 4.09 to 4.26) overall	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
1 player participating in 1 game or practice	Not Reported	Descriptive analyses included the frequency and proportion of injuries. Comparative analyses included the calculation of risk or injury rate ratios (IRRs)	Not Reported	most concussions at the high school and college levels occurred in practices (57.7% and 57.6%, respectively). In addition, no concussions were reported in youth football players aged 5 to 7 years, even though this age group contributed more than 7000 AEs. 2223 reported concussions over 5 years	9.6% youth, 4.0% HS, 8.0% in college sustained a concussion	CI 1.00
1 AE was defined as 1 athlete participating in 1 exposure event)	Not Reported	Regression Analysis of AE	Not Reported	overall rate of 4.17 per 10 000 AEs (95% CI: 4.09 to 4.26)	5.5% sustained a concussion	IRRs with associated 95% CIs excluding 1.00 were considered statistically significant
Epidemiology of concussions in 20 high school sports during the 2013-2014 to 2017-2018 school years.	Not Reported	Regression Analysis of AE	-	overall rate of 4.17 per 10 000 AEs (95% CI: 4.09 to 4.26)	Boys' football had the highest overall concussion rate (10.40 per 10000 AEs), followed by girls' soccer (8.19 per 10000 AEs) and boys' ice hockey (7.69 per 10000 AEs)	AE's with associated 95% CIs excluding 1.00 were considered statistically significant

Stat Significance Reporting Bt Exper Group/Control (p =)	
Indoor volleyball athletes had significantly higher injury rates compared with beach volleyball players for concussion (7.5% vs 6.5%; P < .0001)	
2006 to 2010 was 1.6, significantly less than the 5.0 games missed from 2011 to 2014, following the institution of the NBA concussion protocol (P = .023)	
- The older the rider, the lower likelihood of suffering a head injury (OR=0.989, p=0.00005, 95% CI 0.985 to 0.993) - Riders involved in only a single incident type had higher likelihood of suffering a head injury (OR=1.292, p=0.011, 95% CI 1.061 to 1.573).	
Not Reported	
Loss of consciousness was nonsignificant in equestrian sports (p = .290) compared to roller blading, ski/snowboarding and aquatic based sports	
Over an 11-year period there was a statistically significant increase in 0.97. Additionally, The total number of players experiencing a concussion compared with the total number of players experiencing LoC over the 11-year time period had a correlation coefficient of r = 0.82	
Those with a concussion were more likely to have sought mental health treatment during college (32% vs. 23%; p=0.03).	
Four factors impacted concussion risk Older age (OR 1.03, 95% CI 1.01-1.06; p = 0.015) Females (OR 2.54, 95% CI 1.04-6.21; p = 0.04) having a diploma in horse riding vs not (OR 0.27, 95% CI 0.11-0.65; p = 0.004)	
Not reported	
Not reported	
Skiers and snowboarders were less likely to have head injuries than sledgers (odds ratio [OR] = 0.45; 95% CI, 0.21 to 0.98)	
Soccer players were found to have an 11.1 times greater chance of suffering a concussion during the 1997 season if they had sustained a previous recognized concussion while playing soccer (p < 0.05). Football players who had suffered a previous recognized concussion not occurring during football were 4.2 times as likely to sustain a concussion (p < 0.05)	
Not reported	
Impact to the back of the head or helmet was the strongest factor suggestive of concussion (PPV 40%, p=0.028)	
Not sig	
Goalie status (β = 0.58, t(639) = 3.829, p = 1.4 × 10 ⁻⁴) and maximum level (β = 0.41, t(639) = 8.25, p < 8.8 × 10 ⁻¹⁶) significantly predicted the number of concussions	
The incidence of TBI increased from 1996 to 2010 among adolescents (p<0.003).	
Snowboarders had a greater likelihood of concussion compared to skiers (estimated-β, 2.1; 95% CI, 1.48-2.85) after adjusting for helmet status and age. Overall, imputing missing values for helmets status had no effect on outcome for concussion.	
The incidence rate of concussions increased significantly among players in all positions after the implementation of the 7-day DL rule in 2011 (P<0.003)	
Not Reported	
History of at least one prior concussion was significantly associated with incident concussion (Z = 2.03; p = 0.043)	
It is shown that longer rehabilitation periods become more common over the years (P = 0.015)	
Apart from motocross, sailing showed the highest incident rate per 1000 DAEs at 3.73 (95% CI: 0.31 to 7.15)	
Risk Rate was found statistically significant (78.5; 95% CI, 24.4-252.5)	
Not Reported	
History of concussion was the only moderately strong risk factor for concussion (rate ratio = 2.28, 95% confidence interval: 1.24, 4.18).	
p=0.42	
When comparing the ratios of the head injuries to head incidents between the two countries, Iceland had rates 2.15 times higher than Norway (p < 0.0001).	
No statistically significant difference in females vs males in incidence risk P = .85, but females vs males in return-to-play was statistically significant (20 days vs 10 days, p = 0.02)	
females were 1.57 times more likely to experience poor outcomes (eg, severe disability, persistent vegetative state) than males and more likely to die post- trauma (95% CI 1.09-2.82).	
Not Reported	
p=0.004.	
Women had a higher injury incidence (5.8, 95% CI 4.8 to 6.9) versus men (5.3, 95% CI 3.2 to 4.6; RR 1.48, 95% CI 1.15 to 1.90) by 100 athletes	
Not Reported	
The overall basket toss injury rate (95% CI) was 0.75 (0.41-1.08) per 1,000,000 cheerleaders (0.76 [0.36-1.15] and 0.72 [0.09-1.36] for high school and college, respectively)	
- Girls had a higher rate of concussion (0.36 concussions per 1000 A-Es) than boys (0.22 concussions per 1000 A-Es) (RR = 1.68, 95% CI = 1.08, 2.60, P = .03)	
- Girls had a higher rate of concussion (0.21 concussions per 1000 A-Es) than boys (0.07 concussions per 1000 A-Es) (RR = 2.93, 95% CI = 1.94, 5.24, P < .01).	
- Concussions represented a significantly greater proportion of total injuries in softball players (6.5%, n = 3568) than in baseball players (2.9%, n = 1991) (PR = 1.91, 95% CI = 1.81, 2.01, P < .01)	
Not Reported	

Table

NCAA ISS, 8.0% of the total injuries recorded (n = 111); HS RIO, 3.8% of the total injuries recorded (n = 30); P < 0.001.
No statistical difference between sexes
Concussion rate was higher in competition than in practice (RR = 3.30; 95% CI = 3.02, 3.60). -Among sports in which both sexes participated (ie, baseball or softball, basketball, crew, cross-country, lacrosse, soccer, swimming and diving, tennis, indoor track and field, and outdoor track and field), the collective overall SFC rate was higher in girls than in boys (RR = 1.56; 95% CI = 1.34, 1.81) -Symptom-resolution time greater than 14 days was higher in girls than in boys (33.0% [n = 125] versus 24.4% [n = 78]; IPR = 1.34; 95% CI = 1.12, 1.60) There were no significant differences between mouthguard users and nonusers in rates of concussions (0.35 vs 0.55) or oral soft tissue injuries (0.69 vs 1.06)
Significant difference from 2010-2011 vs 2009-2010
- When considering sex-comparable sports only, the concussion rate was higher in girls than boys (0.66 vs 0.18/1,000 AEs, IRR=3.73, 95% CI=1.24, 11.23). - Concussion rates did not vary between girls and boys (0.61 vs 0.87/1,000 AEs, IRR=0.70, 95% CI=0.41, 1.19
collegiate teams were more likely to sustain a concussion or CHI than were cheerleaders on other types of teams (P = 0.02, OR = 3.10, 95% CI = 1.20, 8.06).
Sig in both collegiate and high school as both CI's crossed 1
Not Reported
5 of 9 showed a significant increase in concussion rates over time, including football (P = .002), boys basketball (P = .003), boys wrestling (P = .014), boys baseball (P = .036), and girls softball (P = .017)
girls had a higher concussion rate (1.7) than boys (1.0) (RR, 1.7; 95% CI, 1.4-2.0).
Diagnosed concussions over the 2014-2015 season were more likely to occur during the regular season (P < .001), in a game setting (P < .001).
only significant IRR was that the concussion rate was higher in Division I men than Division III men (IRR = 1.29; 95% CI, 1.02-1.65)
Not Reported
Concussion injury rate ratio was statically different in games vs practices (rate ratio = 3.6, 95% CI = 3.4, 3.8)
Concussion injury rate ratio was statically different in games vs practices (Rate Ratio= 3.3, 95% CI = 2.8, 4.0)
The injury rate from 2008-2009 through 2013-2014 was higher in college than in high school (3.25 versus 1.73/1000 AEs; IRR = 1.89; 95% CI = 1.63, 2.18).
the concussion rate was higher in competition than practice (10.37 vs 2.04 per 10000 AEs; IRR = 5.09; 95% CI: 4.89 to 5.31
sole difference found was that the college concussion rate (0.53 per 1000 AEs) was lower than that in high school (0.66 per 1000 AEs; IRR, 0.80; 95% CI, 0.67-0.96)
Not Reported
among sex-comparable sports, girls had larger proportions of concussions that were recurrent than boys did (9.3% vs 6.4%; injury proportion ratio = 1.44; 95% CI: 1.11 to 1.88). sex-comparable sports, concussion rates were higher in girls than in boys (3.95 vs 1.51 per 10,000 AEs; injury rate ratio =