Table of Contents Table 1. Post-COVID Condition Diagnosis Codes
Table 2. Summary of the Use of the Post-COVID Condition Diagnosis Code in Medicare Fee-for-Service
Table 3. Summary of Time Gaps Between Post-COVID Condition and COVID-19 Diagnoses
Table 4. Concurrent <i>ICD-10-CM</i> Diagnosis Summary – COVID-19 Cases with and without Long COVID among Community-Dwelling and Nursing Home Beneficiaries Aged ≥65 Year
Figure 1. Weekly Summary of Medical Encounters with Diagnosis Code B94.8 or U09.9

eTable 1. Post-COVID Condition Diagnosis Codes

Diagnosis Code	Diagnosis Code Description
U09.9	Post COVID-19 condition, unspecified
B94.8	Sequelae of other specified infectious and parasitic diseases

eTable 2. Summary of the Use of the Post-COVID Condition Diagnosis Code in Medicare Fee-for-Service

Category	Medicare FFS		Community Dwelling		Nursing Home		ESRD	
	No.	0/0 a	No.	%	No.	%	No.	%
Total Beneficiaries	31,847,927		28,804,470		2,743,507		398,572	
Post- COVID Condition	193,691	0.6%	163,931	0.6%	26,410	1.0%	4,376	1.1%
Post- COVID Condition with Prior COVID-19 diagnosis	161,007	83.1%	132,746	81.0%	25,178	95.3%	4,078	93.2%

Abbreviations: FFS, Fee-for-Service; ESRD, End-Stage Renal Disease

^a The percent is calculated using the number in each row as the numerator and the number in the previous row as the denominator.

eTable 3. Summary of Time Gaps Between Post-COVID Condition and COVID-19 Diagnoses

Category Total Beneficiaries	Community	y Dwelling	Nursing Home	
	No.	%	No.	0/0
Total Beneficiaries	142,255		13,384	
All post-COVID condition diagnoses <28 days after first COVID-19 diagnosis	36,429	25.6%	3,004	22.4%
Post-COVID condition diagnoses <28 days and ≥28 days after first COVID-19 diagnosis	15,836	11.1%	1,810	13.5%
First post-COVID condition diagnosis ≥28 days after first COVID-19 diagnosis	89,990	63.3%	8,570	64.0%

eTable 4. Concurrent *ICD-10-CM* Diagnosis Summary – COVID-19 Cases with and without Long COVID among Community-Dwelling and Nursing Home Beneficiaries Aged ≥65 Year

Diagnosis Code	Diagnosis Code Description	COVID-19 Only ^a	Long COVID ^b	
		Claims, %	Claims, %	Rate Ratio ^c
Community-Dw	elling Beneficiaries	n = 13,663,828	n = 261,417	
R053	Chronic cough	0.0%	1.9%	39.5
Z430	Encounter for attention to tracheostomy	0.0%	0.9%	36.8
R5382	Chronic fatigue, unspecified	0.1%	2.1%	28.9
G7289	Other specified myopathies	0.0%	0.7%	26.1
Z8616	Personal history of COVID-19	0.2%	6.0%	24.2
R0609	Other forms of dyspnea	0.2%	3.6%	22.6
Z431	Encounter for attention to gastrostomy	0.1%	1.1%	19.3
G7281	Critical illness myopathy	0.2%	4.1%	17.1
J8410	Pulmonary fibrosis, unspecified	0.3%	3.9%	15.4
Z8701	Personal history of pneumonia (recurrent)	0.2%	3.4%	14.6
J849	Interstitial pulmonary disease, unspecified	0.2%	2.1%	11.0
J9611	Chronic respiratory failure with hypoxia	0.5%	4.2%	8.9
M5450	Low back pain, unspecified	0.2%	1.2%	8.1
R4189	Other symptoms and signs involving cognitive functions and awareness	0.1%	1.1%	7.6
Z7409	Other reduced mobility	0.2%	1.4%	6.1
Nursing Home E	Beneficiaries	n = 4,378,517	n = 26,411	
Z430	Encounter for attention to tracheostomy	0.0%	1.1%	31.5
G7281	Critical illness myopathy	0.1%	2.2%	19.8
Z8616	Personal history of COVID-19	0.6%	10.0%	16.7
Z8701	Personal history of pneumonia (recurrent)	0.2%	2.2%	10.9
J8410	Pulmonary fibrosis, unspecified	0.2%	1.6%	10.2
Z8619	Personal history of other infectious and parasitic diseases	0.3%	2.7%	8.3
Z20822	Contact with and (suspected) exposure to COVID-19	0.5%	3.7%	7.2
Z431	Encounter for attention to gastrostomy	0.3%	1.7%	5.9
M5450	Low back pain, unspecified	0.2%	1.0%	5.8
F32A	Depression, unspecified	0.4%	2.0%	5.7
Z9911	Dependence on respirator [ventilator] status	0.3%	1.3%	4.8
J9611	Chronic respiratory failure with hypoxia	0.7%	2.9%	4.5
G928	Other toxic encephalopathy	0.1%	0.3%	4.3
J9622	Acute and chronic respiratory failure with hypercapnia	0.3%	1.2%	4.1
Z9981	Dependence on supplemental oxygen	1.2%	4.9%	4.1

 $^{^{\}rm a}$ COVID-19 only refers to claims for individuals with a COVID-19 diagnosis without a subsequent post-COVID condition diagnosis.

^b Long COVID refers to claims for individuals with a post-COVID condition diagnosis ≥28 days after their first COVID-19 diagnosis.

^c The rate ratio was ordered in descending order among the top 100 most frequent diagnoses on Long COVID and COVID-19 only claims.



