

SUPPLEMENTARY MATERIAL

A Pilot Randomized Trial of Combined Cognitive-Behavioural Therapy and Exercise Training Versus Exercise Training Alone for the Management of Chronic Insomnia in Obstructive Sleep Apnea

Amanda Cammalleri^{1*}, Aurore A. Perrault^{1,2*}, Alexandra Hillcoat¹, Emily Carrese-Chacra³, Lukia Tarelli^{1,2,3}, Rahul Patel¹, Marc Baltzan^{4,7}, Florian Chouchou^{1,5}, Thien Thanh Dang-Vu^{1,2,6}, Jean-Philippe Gouin^{2,3,6}, Veronique Pepin^{1,6,7**}

1. *Department of Health, Kinesiology, and Applied Physiology, Concordia University, Montreal, QC, Canada*
2. *Centre de Recherche de l'Institut Universitaire de Gériatrie de Montréal, CIUSSS Centre-Sud-de-l'Île-de-Montréal, QC, Canada*
3. *Department of Psychology & Centre for Clinical Research in Health, Concordia University, Montreal, QC, Canada*
4. *Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, QC, Canada*
5. *EA4075 IRISSE - Département STAPS, Université de La Réunion, France*
6. *PERFORM center, Concordia University, Montreal, QC, Canada*
7. *Axe maladies chroniques, Centre de recherche du CIUSSS du Nord-de-l'Île-de-Montréal, QC, Canada*

** These authors contributed equally*

*** Corresponding author*

Supplemental Results




Figure S1 – Change in insomnia severity

Individual changes (black line) and mean (green and orange bars) ISI score at baseline (T1), 8-weeks post-randomization (T2) and 16-weeks post-randomization (T3) for the EX group (left panel; N=8) and CBTi-EX group (right panel; N=6)

Table S1 – Change in cardiorespiratory fitness

EX Group				CBTi-EX Group			
Participant Number	T1	T2	T3	Participant Number	T1	T2	T3
E1	20.4	21.8	NA ¹	C11	21.6	20.8	NA ⁴
E2	23.3	29.5	25.3	C12	43.2	40.5	44
E3	14.6	18.8	NA ¹	C13	19	16.4	16.9
E4	13.2	20.9	20.1	C14	14.7	14.8	17
E5	25.5	27.3	28.4	C15	17.6	16.3	14.8
E6	28.7	26.6	28	C16	19.3	17.6	19.7
E7	27.2	29.7	34.6	C17	13.5	14.6	16.9
E8	13.3	14.3	NA ³	C18	25.2	25.2	NA ⁵
E9	11	NA ²	NA ²	C19	23.6	32.2 ⁶	27.4
E10	18.6	NA ²	24.7	<i>Mean</i>	<i>21.97</i>	<i>22.04</i>	<i>22.39</i>
<i>Mean</i>	<i>19.58</i>	<i>23.6125</i>	<i>26.85</i>	<i>SD</i>	<i>8.83</i>	<i>9</i>	<i>10.38</i>
<i>SD</i>	<i>6.41</i>	<i>5.54</i>	<i>4.82</i>				

Individual changes and mean (± SD) for VO₂ peak (ml/kg/min) at baseline (T1), 8-weeks post-randomization (T2) and 16-weeks post-randomization (T3) for the EX group (left panel) and CBTi-EX group (right panel)

NA¹- Could not come in for CPET for medical reasons

NA²- Could not come in for CPET due to COVID quarantine

NA³- Could not come in for CPET due to personal reasons

NA⁴- Did not exercise from T2- T3 due to lack of accessibility. Was not included in any analyses.

NA⁵- Could not come in for CPET due to COVID quarantine

⁶-CPET conducted outside the time window prescribed by the protocol (9 months instead of 8-weeks) due to COVID restrictions

Table S2 – Change in objective sleep (PSG-related measures).

PSG measures	T1	T2	T3	Group*Time effect	Group effect	Time effect
<i>Sleep Efficiency (%)</i>						
EX	75.5 ± 11.5	76.1 ± 11.8	68.0 ± 5.2	S(2,26) =2.8	S(1,13) =3.7	S(2,26) =3.2
CBTi-EX	82.2 ± 14.8	78.4 ± 11.8	80.7 ± 11.1	p=.24	p=.054	p=.19
<i>Total Sleep Time (min)</i>						
EX	357.1 ± 73.6	343.0 ± 50.2	316.3 ± 46.6	F(2,26) =0.52	F(1,13) =0.65	F(2,26) =0.67
CBTi-EX	365.0 ± 76.5	344.6 ± 62.7	358.1 ± 61.4	p=.6	p=.43	p=.51
<i>Sleep Onset Latency (min)</i>						
EX	13.6 ± 15.9	19.0 ± 23.4	16.9 ± 12.4	S(2,26) =0.75	S(1,13) =1.21	S(2,26) =1.89
CBTi-EX	14.0 ± 6.3	24.3 ± 19.5	27.3 ± 34.5	p=.68	p=.27	p=.38
<i>Wake After Sleep Onset (min)</i>						
EX	86.6 ± 54.9	84.7 ± 53.3	108.7 ± 62.3	F(2,26) =0.34	F(1,13) =4.8	F(2,26) =0.36
CBTi-EX	61.8 ± 59.2	51.9 ± 29.4	56.5 ± 17.6	p=.71	p=.05	p=.7
<i>Sleep Fragmentation Index (SFI)</i>						
EX	12.3 ± 6.9	13.4 ± 5.5	12.8 ± 6.8	F(2,26) =1.2	F(1,23) =0.3	F(2,26) =0.6
CBTi-EX	11.1 ± 2.8	10.4 ± 3.0	12.3 ± 3.6	p=.3	p=.55	p=.54
<i>Mean SpO2</i>						
EX	90.4 ± 14.8	95.3 ± 1.1	94.2 ± 3.2	F(2,26) =0.38	F(1,13) =0.37	F(2,26) =0.51
CBTi-EX	95 ± 1.1	95.4 ± 1.2	94.6 ± 0.6	p=.56	p=.55	p=.5
<i>SpO2 <90 (%)</i>						
EX	5.7 ± 15.3	0.52 ± 0.68	11.3 ± 29.9	F(2,26) =0.27	F(1,13) =0.24	F(2,26) =0.44
CBTi-EX	1.99 ± 3.9	2.2 ± 4.4	4.34 ± 4.7	p=.66	p=.63	p=.55

Mean (±SD) PSG-extracted measures at baseline (T1), 8-weeks post-randomization (T2) and 16-weeks post-randomization (T3); CBTi-EX N=7, EX N=8

Table S3 – Change in self-reported sleep quality and sleepiness.

Questionnaires	T1	T2	T3	Group*Time effect	Group effect	Time effect
<i>PSQI</i>						
EX	10.8 ± 3.3	10.5 ± 3.9	7.7 ± 2.7*	F(2,22) =0.13	F(1,11) =2.8	F(2,22) =0.13
CBTi-EX	8.5 ± 3.1	8.8 ± 3.3	5 ± 1.6*	p=.87	p=.11	p=.003*
<i>ESS</i>						
EX	10.5 ± 4.6	7.75 ± 4.3	8.5 ± 5.8	F(2,24) =1.36	F(1,12) =0.55	F(2,24) =1.84
CBTi-EX	11.33 ± 5	11.5 ± 5.9	9.83 ± 6.2	p=.27	p=.47	p=.18

Mean (\pm SD) scores at baseline (T1), 8-weeks post-randomization (T2) and 16-weeks post-randomization (T3); *PSQI* CBTi-EX N=6, EX N=7; *ESS* CBTi-EX N=6, EX N=8
PSQI, Pittsburgh Sleep Quality Index; *ESS*, Epworth Sleepiness Scale