

Factors associated with peri-neonatal mortality in Sao Tome & Principe: a prospective cohort study

Alexandra Vasconcelos (A Vasconcelos)^{1*}, Swasilanne Sousa (S Sousa)², Nelson Bandeira (N Bandeira)², Marta Alves (M Alves)³, Ana Luísa Papoila (AL Papoila)³, Filomena Pereira (F Pereira)¹, Maria Céu Machado (MC Machado)⁴

* Correspondence: alexandravasc@gmail.com (AV)
(AV) ORCID: 0000-0001-8042-0846

Author affiliations

1. Unidade de Clínica Tropical - Global Health and Tropical Medicine (GHTM), Instituto de Higiene e Medicina Tropical (IHMT), Universidade NOVA de Lisboa. Lisboa - Portugal
2. Maternidade, Hospital Dr. Ayres de Menezes, São Tomé, República Democrática de São Tomé e Príncipe
3. CEAUL, NOVA Medical School/Faculdade de Ciências Médicas, Universidade NOVA de Lisboa, Lisboa - Portugal
4. Professora Catedrática Jubilada da Faculdade de Medicina de Lisboa, Universidade de Lisboa. Lisboa – Portugal

Abstract

Background

Neonatal mortality reduction is a global goal but its factors in high-burden nations vary and are poorly understood. This study was conducted to identify the factors affecting perinatal neonatal mortality in Sao Tome & Principe (STP), one of the smallest African countries.

Methods

Institution-based prospective cohort study conducted at Hospital Dr. Ayres Menezes. Maternal-neonate dyads enrolled were followed up after the 28th day of life (n=194) for identification of neonatal death-outcome (n=22) and alive-outcome groups (n=172). Data were collected from pregnancy cards, hospital records and face-to-face interviews. After the 28th day of birth, a phone call was made to evaluate the newborn's health status. A logistic regression model was used to identify the relevant factors associated with mortality, a p value <0.05 was considered statistically significant.

Results

The mean gestational age of the death-outcome and alive-outcome groups was 36 (SD=4.8) and 39 (SD=1.4) weeks, respectively. Death-outcome group (n=22) included sixteen stillbirths, four early and two late neonatal deaths. High-risk pregnancy score [cOR 2.91, 95%CI:1.18-7.22], meconium-stained fluid [cOR 4.38, 95%CI:1.74-10.98], prolonged rupture of membranes [cOR 4.84, 95%CI:1.47-15.93], transfer from another unit [cOR 6.08, 95%CI:1.95-18.90], and instrumental vaginal delivery [cOR 8.90, 95%CI:1.68-47.21], were factors significantly associated with mortality. The odds of experiencing death were higher for newborns with infectious risk, IUGR, resuscitation maneuvers, fetal distress at birth, birth asphyxia, and unit care admission. Female newborn [cOR 0.37, 95%CI:0.14-1.00] and birth

weight of more than 2500 g [cOR 0.017, 95%CI:0.002-0.162] were found to be protective factors. In the multivariable model, meconium-stained fluid was significantly associated with death outcome.

Conclusion

Factors such as having a high-risk pregnancy score, meconium-stained amniotic fluid, prolonged rupture of membranes, being transferred from another unit, and an instrumental-assisted vaginal delivery increased 4- to 9-fold the risk of stillbirth and neonatal death. Of the factors associated with peri-neonatal mortality in this study, avoiding health-worker-related factors associated with delays in prompt intrapartum care is a key strategy to implement in Sao Tome & Principe.

Keywords: neonatal mortality, perinatal mortality, stillbirth, neonatal death, Sao Tome & Principe

71 Introduction

72

73 The first 28 days of life—the neonatal period—are the most vulnerable time for a child’s
 74 survival, and a significant number die before birth never having the chance to take their first
 75 breath. Perinatal mortality includes stillbirths and early neonatal deaths (ENND), indicating
 76 the death of a live newborn before the age of seven completed days [1]. Late neonatal deaths
 77 (LNNDs) are those that occur after 7 days to 28 completed days of birth [2].

78 Globally, perinatal mortality accounts for three-fourths of deaths during the neonatal period
 79 [3,4]. More than half of the cases of stillbirths occur when pregnant women are in labor, and
 80 these deaths are directly related to the lack of skilled care at this critical time [5,6]. On the
 81 other hand, the largest contributors to neonatal mortality (ENND plus LNND) are
 82 complications of preterm birth, birth asphyxia, infection, and congenital malformations
 83 although they can differ depending on the country context [7].

84 Only in the last two decades, mainly after Lawn et al published article titled “4 million
 85 neonatal deaths: When? Where? Why?” [8], attention started to be given to the neonatal
 86 period in developing countries although stillbirths are still invisible and missing from the
 87 Sustainable Development Goals (SDG) agenda [5,9,10]. It urges to highlight that stillbirths
 88 and newborn deaths account for twice as many deaths as malaria and human
 89 immunodeficiency virus (HIV) infection combined but have received much less awareness
 90 and funding in these resource-constrained countries [8,9].

91 Understanding the causes of stillbirths (fetal deaths) is also complex, as there are many
 92 promoting and interacting factors [11]. In most low –to medium - income countries (LMICs),
 93 it is difficult to determine the exact reason for the stillbirth; therefore, the cause of death is
 94 often classified as “unexplained” [5]. Researchers report that different risk factors for fetal
 95 death, such as maternal factors (advanced maternal age, high pre-pregnancy body mass index,

96 smoking, low socioeconomic status), obstetric history (grand multiparity, previous stillbirth),
 97 antepartum factors (fewer than four antenatal visits, fetal growth restriction, maternal anemia,
 98 maternal fever and infections, antepartum hemorrhage, hypertension), and intrapartum factors
 99 (preterm birth, extremes of neonatal birth weight, cesarean delivery, operative vaginal
 100 delivery, and assisted breech delivery), are all factors that have been reported in various
 101 studies as causes of stillbirths [6,11-15].

102 Sao Tome & Principe (STP) is an LMIC with low HIV/AIDS prevalence and a malaria
 103 preelimination phase [16,17]. There is an established antenatal care (ANC) service, with a
 104 98% rate of women seen at least once by a skilled health provider during pregnancy and a
 105 95.4% rate of deliveries occurring in health units [17]. In STP, peri-neonatal mortality is
 106 considered a public health problem in the country, given that neonatal deaths accounts for
 107 approximately 43% of all under5 deaths [18]. At the time this study was initiated, there was
 108 an annual rate of 22 stillbirths and 22 newborn deaths per 1000 livebirths [18].

109 We are aware that the concept of knowing what works in terms of reducing perinatal and
 110 neonatal mortality is complicated by a huge diversity of country contexts and of determinants
 111 of maternal and neonatal health [19-22]. However, according to Lawn and other authors,
 112 identifying and addressing avoidable causes of neonatal death is possible even in poorly
 113 functioning health systems, justifying our current study [8,9]. Answering - why, when, and
 114 where - newborns die in Sao Tome & Principe will enable the design of appropriate planning
 115 to prevent this major public health problem since, for appropriate prevention of fetal and
 116 newborn mortality, data pertaining to its determinants are important.

117 To the best of our knowledge, this is the first study to assess peri-neonatal mortality in STP,
 118 and it was undertaken within the context of a broader project on neonatal adverse birth
 119 outcomes (ABOs) in this LMIC [23-25].

120 Therefore, we conducted a prospective cohort study to identify the most important factors
121 associated with mortality among newborns delivered at HAM.

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

Materials and methods

Study design

An institution-based prospective cohort study was conducted at Hospital Dr. Ayres de Menezes (HAM) for mother-neonate dyads followed up until the 28th day after delivery (neonatal period).

Setting

The archipelago of Sao Tome & Principe is one of the smallest sub-Saharan African (SSA) countries, with approximately 200.000 inhabitants and a total land surface of approximately 1,001 km² on two islands (Sao Tome and a smaller island named Principe) [16,17]. The rural districts have ANC services and maternity units with basic obstetric services in Lembá, Lobata, Caué, Me-Zochi, and Principe Island, except for the district of Cantagalo. There is only one hospital in the country, Hospital Dr. Ayres de Menezes (HAM), with the only maternity in the country with Comprehensive Emergency Obstetric and Neonatal Care (EmONC) capable of providing blood transfusions and performing cesarean sections [26]. As a tertiary healthcare facility, HAM receives the most complicated cases from facilities in lower levels of care. There is no health insurance policy in the country or any private maternity units. In STP, there are approximately 3.2 doctors per 10 000 inhabitants and 27 nurses per 10 000 inhabitants, and at HAM, there are three to four obstetricians and one to two general doctors who provide care to the neonates [26]. There are no neonatologists in the country. The midwives in the labor ward are responsible for the initial resuscitation of normal deliveries, and doctors from the pediatric department are called to the labor ward to attend babies in distress.

The HAM maternity unit has a facility-based clinical care unit for ill newborn babies, but there is no neonatal or child intensive care unit in the country. The Neonatal Care Unit (NCU) receives high-risk babies delivered within the institution and referrals from other health facilities or from home with a total capacity to admit six babies. This unit, like others in SSA settings, is basic and able to manage simple neonatal complications such as hypothermia, feeding problems and sepsis suspicion. Although NCU was rebuilt in 2016, there is still a lack of continuous positive airway pressure therapy, surfactant therapy and enteric feeding for assisting sick babies.

Study population and follow-up

All mother-neonate dyads admitted to the HAM maternity unit for childbirth constituted the source population whereas the study populations were selected neonates delivered in the HAM maternity unit during the study period. Recruitment of participants (mother-newborn dyad) occurred from July 2016 to November 2018. During the study period, 4540 deliveries were recorded, corresponding to 450 cesarean deliveries and 3740 normal vaginal births.

The inclusion criteria for participants were as follows: 1) all neonates delivered at HAM with a gestational age of 28 weeks or more and 2) newborns who were born outside the hospital but were later admitted at HAM. A total of 537 newborns were initially enrolled.

The exclusion criteria included the following: 1) all neonates delivered at HAM with a gestational age of less than 28 weeks, 2) newborns whose mothers had no antenatal pregnancy card, 3) newborns whose mothers had cognitive impairment. The newborn was also excluded if his health status was unknown at his twenty-eight days of life.


Consenting participants in the sample were followed up (mother-newborn dyad) throughout their stays until hospital discharge. The survival status of neonates after discharge was ascertained by making a follow-up mobile phone call at the end of the neonatal period. Those

194 who could not be reached by phone after four attempts in different weeks were taken as
 195 nonrespondents, and the mother-newborn dyad was excluded. A flowchart of participation in
 196 the study is shown in Fig 1.

197

198 **Figure 1** Flowchart of participation in the study

199



200

201

202 Sampling method

203 This is a substudy undertaken to estimate adverse birth outcomes (ABOs) and related risk
 204 factors in the country.

205 The sample size initially calculated for the broader project, using Raosoft®
 206 (<http://www.raosoft.com/samplesize.html>), was based on the formula for sample size and
 207 margin of error from the software, reaching the minimum sample size of $S_{\alpha} = 355$, with 355
 208 (95%) and 579 (99%) interval confidence. The original study included 537 mother-neonate
 209 dyads enrolled based on the following assumptions: two-sided 95% confidence level, power
 210 of 80% to detect an odds ratio of at least two for ABO. Since the sample size was not
 211 calculated for present outcomes, to assess peri-neonatal mortality, a power analysis was

performed, varying from 77% to 87% for outcomes such as gestational age (GA<37) for this study. In this study, 194 mother-neonate dyads were included. Participants were selected through random sampling. Each morning, from the pile of mothers' medical folders, every second interval folder was selected and then carried on requesting contentment for enrollment. To guarantee a sample with few biases and effects by means of confounding variables, the study was conducted in different months (two weeks every two other months), avoiding seasonal interference (rain season and malaria period). Women were interviewed only after the delivery, although the invitation and consent were obtained during her admission to the maternity unit before birth (live birth or stillbirth).

Selection of groups: death and alive neonatal outcome

A total of 194 mother-neonate dyads were included and were divided into two groups according to the newborn status (alive versus dead) at the end of the neonatal period. Newborns who died before the 28th day of life (death-outcome group with stillbirths included; n=22) were compared with newborns alive at the 28th day of life (alive-outcome group; n=172) concerning maternal, antepartum, and intrapartum characteristics.

Study Variables

The dependent variables for the study were death (stillbirth and neonatal - early and late - death) and live newborns.

The definition of stillbirth used was the WHO/ICD (for international comparison and reporting) as a baby born without any signs of life at or after 28 weeks of gestation or at least 1000 g in birth weight. [9,27]. Stillbirths were also described according to whether they were an intrapartum or antepartum stillbirth. Intrapartum stillbirth was defined as a dead-born fetus where intrauterine death occurred after the onset of labor and before birth (fresh stillbirth)

[9]. Antepartum stillbirth is a dead born fetus where intrauterine death occurs before the onset of labor (macerated stillbirth) [9].

Neonatal mortality was defined as deaths in the first 28 days of life among live births. Death within the first 24 h (<24 h) or first day of life was recorded in units of completed minutes or hours of life [2]. Deaths occurring ≥ 24 h of life were recorded in days, from day 1 to 27 completed days [2]. The neonatal death definition used in this study was the one from the World Health Organization (WHO) as “deaths among live births during the first 28 completed days of life” [1], which were further subdivided into early neonatal deaths (ENND) as deaths between 0 and 7 completed days of birth and late neonatal deaths (LNND) as deaths after 7 days to 28 completed days of birth [2,28].

For neonates who died at home after discharge, probable causes of death were assigned by the principal investigator (pediatrician) using the International Statistical Classification of Diseases and Related Health Problems ICD-11 coding. For neonates who died in the HAM, the cause of death was assigned by physicians, who work in the hospital, and confirmed by the principal investigator.

The independent variables tested in this study included factors grouped into five categories: 1) newborns’ maternal sociodemographic factors (age, educational status, occupation, marital status, partner’s education, and residence); 2) preconception factors (previous contraceptive utilization), plus current obstetric condition (gravidity, parity, previous abortion and stillbirth, previous cesarean section and preceding birth interval); 3) ANC service (number of visits, gestational age at first ANC visit, obstetric ultrasound, number of fetuses in the ultrasound) plus antepartum factors as positive ANC screenings (high-pregnancy risk score, maternal anemia as hemoglobin concentration < 11 g/dl, bacteriuria, hyperglycemia, Rh incompatibility, intestinal parasitic infection (IPI), malaria, HIV, syphilis, hepatitis B virus

and sickle cell); 4) health facility-related factors (being transferred from another unit, who assisted the delivery and partograph use) plus intrapartum factors as mode of delivery and complications (fetal malpresentation [29], umbilical cord complication [30], prolonged rupture of membranes [31,32], meconium-stained amniotic fluid [33], postpartum hemorrhage as bleeding >500 mL, preeclampsia defined as hypertension $\geq 140/90$ mmHg and proteinuria in dipsticks in women who were normotensive at ANC, and obstructed labor [34]); 5) newborns characteristics as (gestational age, sex, birth weight) and complications (intrauterine growth restriction (IUGR), congenital anomalies, infectious risk, neonatal resuscitation, fetal distress at birth, birth asphyxia and admission at NCU) [35-37].

Data collection

Data on antepartum, intrapartum, and postpartum characteristics of participants were gathered and collected from ANC pregnancy cards, obstetric maternal and newborn clinical records. For antepartum data, relevant details of the perinatal history and antenatal period were collected systematically from the ANC pregnancy card. Intrapartum data were collected from labor follow-up sheets, delivery summaries and maternal medical records. Postpartum data were abstracted from newborn birth charts and/or newborn medical records if admitted to the NCU.

Maternal sociodemographic characteristics were supplemented with a structured administered questionnaire through a face-to-face interview of the mothers 12-24 hours after delivery, similar to other studies from LMICs [38].

Data quality alive-outcome group

The questionnaires were administered in Portuguese, the country national language. The questionnaire was pretested at HAM one month before data collection in 23 mothers, and

modification was made based on the pretest result, mainly adjusting terminology for more culturally friendly terms. Women's consent to participate in the study was obtained at the time of admission at HAM, but the interview was held after a woman was stabilized and ready to be discharged. Continuous follow-up and supervision of data collection were made by the supervisors. The collected data were checked daily for completeness. The principal investigator (pediatrician) executed and was responsible for all main activities as follows: 1) obtaining consent and enrollment of the participants, 2) data collection from antenatal cards plus maternal clinical and newborns' records, 3) newborns' clinical observation (for diagnosis confirmation), 4) face-to-face interviews, 5) administering all phone interviews and 6) data collection entry into the database.

Data management

The anonymity and safety of the participants were ensured. Data were secured in a confidential and private location. Participants were referred to by identification numbers, and the informed consent forms were kept separate from the questionnaires. Both could only be linked by a coding sheet available only to the principal investigator.

Data analysis

Data were entered into the QuickTapSurvey app (©2010-2021 Formstack), and the dataset was exported to Excel for cleaning and further analysis using the Statistical Package for the Social Sciences for Windows, version 25.0 (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp.). All data were checked for completeness and accuracy by the principal investigator and a qualified biostatistician.

Descriptive statistics, namely, frequencies and percentages were estimated. A univariable analysis was also performed. In this study, the neonatal death-outcome group was coded as 1, and the neonatal alive-outcome group was coded as 0. The proportion of missing data ranged from 0.8 to 10% across variables, and missing values higher than 10% were described in the analysis.

Univariable logistic regression analysis was used to estimate the unadjusted effect of each independent variable on the dependent variable. Variables that had a p value less than 0.25 were considered in the multivariable model. A p value < 0.05 was considered statistically significant in multivariable logistic regression analysis.

Ethics approval and consent to participate

The Ministry of Health of Sao Tome & Principe and the main board of the Hospital Dr. Ayres de Menezes are dedicated ethics oversight bodies, and both approved this study. At the time of the study protocol was submitted (year 2016), there were no ethics committees in STP. All methods were performed in accordance with the relevant guidelines and regulations in practice. Written informed consent was obtained from all participants (or their parent or legal guardian in the case of adolescent under 16 or for illiterate pregnant women) after the purpose of the research was explained orally by the principal investigator. Participants' or their legal representatives' also consented to have the results of this research work published. Participation in the survey was voluntary, as participants could decline to participate at any time during the study.

Results

A total of 194 newborns were followed up during their first 28 days of life. In this study, the newborn's mean gestational age (GA) was 38.86 weeks with a standard deviation (SD) of 2.26 (minimum 28 - maximum 42 weeks). The death-outcome group (n=22) had a mean GA and birth weight of 36 (SD=4.83) and 2515 (SD=997) g, respectively, while their counterparts (172 in the alive-outcome group) had 39 (SD=1.41) weeks and 3209 (SD=507) g, respectively. The mean maternal age was 27.14 years, with a SD of 6.86 (minimum 15 - maximum 43) years old. The mean maternal age for the death-outcome group and alive-outcome group was 30.73 (SD=7.45) and 26.68 (SD=6.66) years, respectively.

The death-outcome group under-study included 16 stillbirths (72.7%), four (18%) ENND and two (9%) LNND. Stillbirths were 69% intrapartum stillbirths (fresh stillbirths), and 31% were antepartum stillbirths (macerated stillbirths). Stillbirth characteristics are further described in Additional file 1, and the early and late neonatal deaths (ENND and LNND) are described in Additional file 2.

The maternal characteristics as well as antepartum, intrapartum, and postpartum factors for the total of the participants and for the death-outcome group versus the alive-outcome group are described in Tables 1 and 2.

359 There were no maternal deaths in this study, with a total of 2 maternal near-misses occurring
360 in the stillbirths' mothers' death-outcome group numbers 5 and 11 that needed hysterectomy
361 intervention due to atonic uterus with major obstetric hemorrhage.

362

363 **Table 1** Univariable logistic regression analysis of peri-neonatal mortality among newborns admitted
364 at HAM, Sao Tome & Principe (n= 194; death-outcome group :22 and alive-outcome group: 172)

Variables	Categories	Total n=194	Death-outcome group n=22 n (%)	Alive- outcome group n=172 n (%)	cOR	(95%) CI	p value
Sociodemographic characteristics							
Mother's age	14-19	28 (14.4)	3 (13.6)	25 (14.5)	1		
	20-34	131 (67.5)	11 (50.0)	120 (69.8)	0.764	0.199-2.939	0.695
	≥35	35 (18)	8 (36.4)	27 (15.7)	2.469	0.588-10.36	0.217
Mother's education	none+primary	106 (54.6)	19 (86.4)	87 (50.6)	1		
	secondary	88 (45.4)	3 (13.6)	88 (45.4)	0.162	0.046-0.566	0.004
Mother's occupation	unemployed	139 (72.0)	16 (72.7)	123 (71.9)	1		
	employed	54 (28)	6 (27.3)	48 (28.1)	0.961	0.355-2.601	0.961
Marital status	union/married	124 (63.9)	14 (63.6)	110 (64)	0.986	0.392-2.482	0.977
	single	70 (36.1)	8 (36.4)	62 (36)	1		
Baby's father education	none+ primary	63 (41.7)	9 (81.8)	54 (38.6)	1		
	secondary	88 (58.3)	2 (18.2)	86 (61.4)	0.14	0.029-0.67	0.014
Residence ¹	urban	92 (48.2)	9 (45)	83 (48.5)	0.867	0.342-2.200	0.765
	rural	99 (51.8)	11 (55)	88 (51.5)	1		
Preconceptional							
Contraception previous use	yes	38 (24.8)	3 (20)	35 (25.4)	0.736	0.196-2.760	0.649
	no	115 (75.2)	12 (80)	103 (74.6)	1		

365

Obstetric history							
Gravidity	1	42 (21.6)	4 (18.2)	38 (22.1)	1		
	2-5	100 (51.5)	8 (36.4)	92 (53.5)	0.826	0.235-2.907	0.766
	≥5	52 (26.8)	10 (45.5)	42 (24.4)	2.262	0.655-7.815	0.197
Parity	0	53 (27.3)	5 (22.7)	48 (27.9)	1		
	1-4	121 (62.4)	12 (54.5)	109 (63.4)	1.057	0.353-3.166	0.921
	≥5	20 (10.3)	5 (22.7)	15 (8.7)	3.200	0.814-12.574	0.096
Previous abortion	yes	59 (30.4)	6 (27.3)	53 (30.8)	0.842	0.312-2.272	0.734
	no	135 (69.6)	16 (72.7)	119 (69.2)	1		
Previous stillbirth	yes	17 (8.8)	4 (18.2)	13 (7.6)	2.718	0.801-9.225	0.109
	no	177 (91.2)	18 (81.8)	159 (92.4)	1		
Poor birth spacing ²	yes	39 (20.1)	6 (27.3)	33 (19.2)	1.580	0.574-4.346	0.376
	no	155 (79.9)	16 (72.7)	139 (80.8)	1		

Antenatal care							
GA at first ANC visit	≤12	98 (60.5)	6 (27.3)	92 (53.5)	1		
	>12	58 (35.8)	9 (40.9)	55 (32)	2.509	0.847-7.430	0.097
Number of ANC visits	1-4	19 (10.1)	2 (10)	17 (10.1)	1		
	5-7	84 (44.4)	16 (80)	68 (40.2)	2.000	0.419-9.547	0.385
	≥ 8	86 (45.5)	2 (10)	84 (49.7)	0.202	0.027-1.538	0.123
Obstetric ultrasound	0	73 (38.4)	7 (35)	66 (38.8)	1		
	1	87 (45.8)	12 (60)	75 (44.1)	1.509	0.561-4.056	0.415
	2	30 (15.8)	1 (5)	29 (17.1)	0.349	0.041-2.976	0.336
Twin pregnancy	yes	14 (7.2)	0	14 (8.1)	-	-	-
	no	180 (92.8)	22 (100)	158 (91.9)	1		

366

367

Antenatal care screenings							
High-pregnancy risk ³	yes	70 (36.1)	13 (59.1)	57 (33.1)	2.914	1.176-7.220	0.021
	no	124 (63.9)	9 (40.9)	115 (66.9)	1		
Maternal disease during pregnancy ⁴	yes	123 (63.4)	11 (50)	112 (65.1)	0.622	0.244-1.587	0.321
	no	66 (34)	9 (40.9)	57 (33.1)	1		
	missing	5 (2.6)	2 (9.1)	3 (1.7)	4.222	0.618-28.867	0.142
Maternal anaemia ⁵	yes	59 (30.4)	6 (27.3)	53 (30.8)	1.736	0.533-5.655	0.360
	no	98 (50.5)	6 (27.3)	92 (53.5)	1		
	not tested	37 (19.1)	10 (45.5)	27 (15.7)	5.679	1.892-17.048	0.002
Bacteriuria	yes	62 (32)	5 (22.7)	57 (33.1)	0.593	0.208-1.690	0.328
	no	132 (68)	17 (77.3)	115 (66.9)	1		
Hyperglycaemia ⁶	yes	8 (4.1)	1 (4.5)	7 (4.1)	1.122	0.132-9.578	0.916
	no	186 (95.9)	21 (95.5)	165 (95.9)	1		
Rh incompatibility	yes	7 (3.6)	0	7 (4.1)	-	-	-
	no	187 (96.4)	22 (100)	165 (95.9)			
IPIs	yes	98 (50.5)	8 (36.4)	90 (52.3)	0.521	0.208-1.305	0.164
	no	96 (49.5)	14 (63.6)	82 (47.7)			
Malaria	yes	1 (0.5)	0	1 (0.6)	-	-	-
	no	193 (99.5)	22 (100)	171 (99.4)			
HIV	yes	2 (1.0)	1 (5.0)	1 (0.6)	1		
	no	192 (99)	19 (95)	163 (99.4)	0.123	0.007-2.037	0.143
HsAg positive	yes	6 (3.1)	2 (9.1)	4 (2.3)	1		
	no	113 (58.2)	10 (45.5)	103 (59.9)	0.194	0.032-1.195	0.077
Sickle cell positivity ⁷	yes	13 (6.8)	1 (4.5)	12 (7.0)	0.631	0.078-5.102	0.666
	no	181 (93.2)	21 (95.5)	160 (93)	1		

368

Health facility-related factors							
Baby delivered at HAM	yes	188 (96.9)	21 (95.5)	167 (97.1)	1.590	0.177-14.275	0.679
	no	6 (3.1)	1 (4.5)	5 (2.9)	1		
Transferred from another unit*	yes	16 (8.2)	6 (27.3)	10 (5.8)	6.075	1.953-18.900	0.002
	no	178 (91.8)	16 (72.7)	162 (94.2)	1		
Delivery assisted by	obstetrician	36 (18.6)	5 (22.7)	31 (18)	1		
	midwife	153 (78.9)	16 (72.7)	137 (79.7)	0.724	0.247-2.126	0.557
	home labor	5 (2.6)	1 (4.5)	4 (2.3)	1.550	0.143-16.852	0.719

Partograph use	yes	76 (39.2)	9 (40.9)	67 (39)	1.085	0.440-2.678	0.860
	no	118 (60.8)	13 (59.1)	105 (61)	1		
Fetal malpresentation ⁸	yes	2 (1)	0	2 (1.2)	-	-	-
	no	192 (99)	22 (100)	170 (98.8)			
PROM ⁹	yes	15 (8.1)	5 (23.8)	10 (6.1)	4.844	1.473-15.930	0.009
	no	171 (91.9)	16 (76.2)	155 (93.9)	1		
Pre/Eclampsia ¹⁰	yes	17 (8.8)	2 (9.1)	15 (8.7)	1.047	0.223-4.917	0.954
	no	177 (91.2)	20 (90.9)	157 (91.3)	1		
Obstructed labour ¹¹	yes	22 (11.3)	4 (18.2)	18 (10.5)	1.901	0.579-6.239	0.289
	no	172 (88.7)	18 (81.8)	154 (89.5)	1		
Postpartum haemorrhage ¹²	yes	2 (1)	0	2 (1.2)	-	-	-
	no	192 (99)	22 (100)	170 (98.8)			
Normal Vaginal delivery	yes	155 (79.9)	15 (68.2)	140 (81.4)	0.490	0.185-1.300	0.152
	no	39 (20.1)	7 (31.8)	32 (18.6)			
Caesarean section	yes	33 (17)	4 (18.2)	29 (16.9)	1.096	0.345-3.477	0.877
	no	161 (83)	18 (81.8)	143 (83.1)			
Previous caesarean	yes	10 (5.2)	2 (9.1)	8 (4.7)	2.050	0.407-10.333	0.384
	no	184 (94.8)	20 (90.9)	164 (95.3)			
Instrumental vaginal delivery ¹³	yes	6 (3.1)	3 (13.6)	3 (1.7)	8.895	1.676-47.208	0.010
	no	188 (96.9)	19 (86.4)	169 (98.3)			
Meconium-stained amniotic fluid	yes	43 (22.2)	11 (50)	32 (18.6)	4.375	1.744-10.975	0.002
	no	151 (77.8)	11 (50)	140 (81.4)			
Umbilical cord complication	yes	8 (4.1)	1 (4.5)	7 (4.1)	1.122	0.132-9.578	0.916
	no	186 (95.9)	21 (95.5)	165 (95.9)			

369

Newborn's characteristics							
Gestational Age ¹⁴	28 - 31	3 (1.5)	3 (13.6)	0	1		
	32 - 36	8 (4.1)	4 (18.2)	4 (2.3)	-	-	-
	37 - 41	176 (90.7)	15 (68.2)	161 (93.6)	-	-	-
	≥42	7 (3.6)	0	7 (4.1)	-	-	-
Sex	feminine	92 (47.4)	6 (27.3)	86 (50)	0.375	0.140-1.004	0.051
	masculine	102 (52.6)	16 (72.7)	86 (50)			
Birth weight ¹⁵	<1500 g	5 (2.6)	4 (18.2)	1 (0.6)	1		
	1500-2499 g	18 (9.3)	7 (31.8)	11 (6.4)	0.159	0.015-1.732	0.131
	2500g-3999 g	161 (83)	10 (45.5)	151 (87.8)	0.017	0.002-0.162	<0.001
	≥ 4000 g	10 (5.2)	1 (4.5)	9 (5.2)	0.028	0.001-0.564	0.020
IUGR ¹⁶	yes	9 (4.6)	6 (27.3)	3 (1.7)	21.125	4.82-92.586	<0.001
	no	185 (95.4)	16 (72.7)	169 (98.3)	1		
Infectious risk ¹⁷	yes	50 (25.8)	14 (63.6)	36 (20.9)	6.611	2.574-16.978	<0.001
	no	144 (74.2)	8 (36.4)	136 (79.1)			
Congenital malformation	yes	4 (2.1)	4 (18.2)	0	-	-	-
	no	190 (97.9)	18 (81.8)	172 (100)			

370

371 This is the Table 1 legend.

Abbreviations: ANC – antenatal care; CI – confidence interval; cOR: crude odds ratio; GA – gestational age; HAM: Hospital Dr. Ayres de Menezes; IPI – intestinal parasitic infection; IUGR – intrauterine growth restriction; PROM – prolonged rupture of membranes.

¹Urban residence for women living in the capital city (Água Grande) and rural areas in all other districts (Mé-Zochi, Cantagalo, Lobata, Lembá, Caué and Príncipe Island).

²Poor birth spacing birth intervals of less than 2 years [39]

³High-pregnancy risk is registered in the ANC pregnancy card if the current pregnancy is defined as one or more of the following: 1) pregnant women age less than 15 years old or greater than 35, 2) grand multipara for women with six or more labors, 3) previous history of a stillbirth or early neonatal death, 6) previous caesarean section, and 7) previous hemorrhagic complication.

⁴Maternal disease was operationally defined as one or more of the following conditions during the current pregnancy: pre/eclampsia, gestational diabetes, malaria, bacteriuria, anemia, and intestinal parasitic infection.

⁵anaemia during pregnancy as a hemoglobin concentration <11 g/dl

⁶glycaemia >105 mg/dL,

⁷Through a sickle cell solubility test, which involves treating a thin blood film with sodium dithionate under hypoxic conditions and observing for sickling under a light microscope, that is the screening technique available in STP and performed to pregnant women with anemia or clinical suspicion. A positive result can suggest either sickle cell anemia or the sickle cell trait [40].

⁸Foetal malpresentation was determined if the presenting fetal part was noncephalic (e.g. breech, transverse, oblique) [29].

⁹Prolonged rupture of membrane (PROM) was defined as a rupture of membrane lasting longer than 18 hours before labor began [31,32].

¹⁰Preeclampsia (hypertension $\geq 140/90$ mmHg and proteinuria in dipsticks in women who were normotensive at ANC).

¹¹Obstructed labor was operationally defined as the sum of all cesarean sections due to mechanical problems or fetal distress and all instrumental delivery [34].

¹²Postpartum hemorrhage was defined as >500 mL bleeding.

¹³IVD in STP is only performed by vacuum.

¹⁴Gestational age was estimated from the date of onset of the last normal menstrual period or through ultrasound dating of pregnancy. Prematurity was defined as a delivery before 37 complete weeks of gestation from the date of onset of the last normal menstrual period and subcategorized as very preterm (28 to 31 weeks) and moderate to late preterm (32 to 37 weeks).

¹⁵Low birth weight was defined as a newborn weight less than 2500 grams (up to and including 2499 g) at birth regardless of gestational age. Low birth weight was further categorized into very low birth weight (VLBW, <1500 g) [41]. Low birth weight is a result of preterm birth, intrauterine growth restriction or both [41].

Large birth weight when ≥ 4000 g (macrosomia) irrespective of gestational age.

¹⁶There is still no consensual definition for intrauterine growth restriction (IUGR) or fetal growth restriction in Africa [42]. In this study, it was defined as the term LBW (i.e., birthweight <2500 g and gestational age ≥ 37 weeks of gestation) due to low in utero measurements through obstetric ultrasounds in the country [42].

¹⁷Infectious risk was operationally defined as the sum of all the following risk factors: 1) maternal fever (axillary temperature >37.9 C) at the time of delivery, 2) prolonged rupture of membrane (≥ 18 hours), and/or 3) foul-smelling amniotic fluid [43].

Factors associated with perinatal and neonatal mortality

Binary logistic regression analysis was performed to assess the association of peri-neonatal mortality with several characteristics (Table 1 and Table 2). Crude analysis showed that meconium-stained amniotic fluid, prolonged rupture of membranes, transfer from another unit, and instrumental vaginal delivery were significantly associated with peri-neonatal mortality.

Sociodemographic factors

A maternal secondary education level (cOR 0.162, 95% CI 0.46–0.566, $p=0.004$) and a baby's father secondary education level (cOR 0.140, 95% CI 0.029–0.67, $p=0.014$) were found to be protective factors for peri-neonatal mortality.

Antepartum factors

The odds of peri-neonatal mortality were three times higher among mothers classified as having a high-pregnancy risk (cOR 2.91, 95% CI 1.18–7.22, $p=0.021$). Mothers without hemoglobin test during the ANC follow-up had a higher risk of having a peri-neonatal death (cOR 5.68, 95% CI 1.89–17.05, $p=0.002$) than those without anemia.

429

430 Health facility-related factors

431 The odds of peri-neonatal mortality were six times higher among mothers transferred from
432 another unit compared to those directly admitted at HAM maternity (cOR 6.08, 95% CI 1.95–
433 18.90, p=0.002).

434

435 Intrapartum factors

436 PROM as well as meconium-stained amniotic fluid were other intrapartum factors, with the
437 odds of mortality being almost five times higher for newborns whose mothers had a PROM
438 (cOR 4.84, 95% CI 1.47–15.93, p=0.009), and four times higher for those with a meconium-
439 stained amniotic fluid (cOR 4.38, 95% CI 1.74–10.98, p=0.002).

440 For the mode of delivery, having an instrumental assisted delivery was associated with an
441 eightfold higher risk of perinatal and neonatal mortality (cOR 8.90, 95% CI 1.68–47.21,
442 p=0.010).

443

444

445 Newborns' factors

446 Newborns with intrauterine growth restriction had a twenty-one-fold higher risk of mortality
447 (cOR 21.13, 95% CI 4.82–92.59, p<0.001), and newborns with an infectious risk had almost
448 sevenfold higher odds of dying (cOR 6.61, 95% CI 2.57–16.98, p<0.001). Regarding newborn
449 characteristics, female neonates (cOR 0.38, 95% CI 0.14–1.00, p=0.051) and birth weight
450 greater than 2500 g and lower than 3999 g (cOR 0.017, 95% CI 0.002–0.162, p<0.001), and
451 greater than 4000 g (cOR 0.028, 95% CI 0.001–0.564, p=0.020) were protective factors
452 against peri-neonatal mortality.

Postpartum characteristics were only assessed for a total of six death-outcome group since sixteen stillbirths were not further included for analysis. Performance of neonatal resuscitation, fetal distress at birth (APGAR score at first-minute inferior to seven), birth asphyxia, and admission to the neonatal care unit were all related to an increased risk for neonatal mortality (Table 2).

Table 2 Univariable logistic regression analysis of neonatal mortality among newborns who were live births n= 178 [death-outcome group: 6 and alive-outcome group: 172]

Variables	Categories	Total n=178 ^{III}	Death-outcome group n=6 ^L n (%)	Alive-outcome group n=172 n (%)	cOR	(95%) CI	p value
Neonatal resuscitation	yes	7 (3.9)	2 (33.3)	5 (2.9)	16.700	2.457-113.495	0.004
	no	171 (96.1)	4 (66.7)	167 (97.1)	1		
Fetal distress at birth*	yes	26 (14.6)	5 (83.3)	21 (12.2)	35.952	4.004-322.859	0.001
	no	152 (85.4)	1 (16.7)	151 (87.8)	1		
Birth asphyxia**	yes	10 (5.6)	4 (66.7)	6 (3.5)	55.333	8.421-363.600	<0.001
	no	168 (94.4)	2 (33.3)	166 (96.5)	1		
Admission at NCU	yes	15 (8.4)	3 (50)	12 (7)	13.333	2.425-73.310	0.003
	no	163 (91.6)	3 (50)	160 (93)	1		

This is the Table 2 legend.

Abbreviations: CI – confidence interval; cOR: crude odds ratio; NCU – neonatal care unit.

^{III} stillbirths excluded

*Fetal distress was defined as a low Apgar score <7 at the first minute of life (score from 0 to <7) [24,44].

**Birth asphyxia was defined as a low Apgar score <7 at the fifth minute of life (score from 0 to <7) [24,44].

Multivariable model

473 Considering the small size of the outcome group (n=22), only one significant variable
474 remained in the multivariable model. Newborns whose mothers had meconium-stained
475 amniotic fluid were at a higher risk of peri-neonatal death.

476

477

478

Discussion

In this prospective cohort study, the main aim was to define risk factors, the most vulnerable period for neonatal death - partum, early or late neonatal period - and to identify where they die - in the maternity, at home or in the pediatric ward - as we were able to follow them up until their 28th day of life. In this study, we found that 90% died in HAM maternity before the 7th day of life, that is, during the perinatal period, with mainly being stillborn (73%). The magnitude of stillbirths observed in this study (3%) is in line with the studies conducted in Nigeria (4.8%) [45] and Tanzania (3.5%) [46] and lower compared with studies from Ethiopia (6.7%) [47]. Most stillbirths occurred during intrapartum, as also identified in LMICs similar to STP, and most could have been prevented [48,49].

In this study, intrapartum characteristics, such as pregnant women with meconium-stained amniotic fluid, prolonged rupture of membranes, and instrumental vaginal delivery, were the main factors significantly associated with peri-neonatal death. Health facility-related factors, as pregnant women transferred from another unit were also at a significant risk.

High-pregnancy risk score notification was the only antepartum factor significantly associated with peri-neonatal mortality. For newborns, experiencing fetal distress at birth, needing resuscitation maneuvers, having birth asphyxia and admission to the neonatal unit were identified as high-risk factors for death. Being a female newborn and having a birth weight greater than 2500 grams were found to be protective factors.

Odds of experiencing death were identified as four times higher among newborns from mothers with meconium-stained amniotic fluid. This finding is consistent with studies from Ethiopia [6,50] and Yemen [12] that also reported higher rates of fetal death due to intrauterine passage of meconium into amniotic fluid and a fivefold increase in perinatal mortality compared with low-risk patients with clear amniotic fluid [51]. Meconium-stained

amniotic fluid is associated with fetal distress since the fetus, in response, inhales the meconium, which in turn leads to airway obstruction, surfactant dysfunction and pneumonitis, which leads to loss of the fetus [6].

PROM was also a significant factor associated with neonatal mortality in this study, showing fivefold higher odds for the death-outcome group. This association was also identified in other studies from low-resource constrained countries reporting one-third of stillbirths occurring during labor as a result of prolonged labor or obstructed labor not attended to promptly [35,36,38,52,53]. The reason for this might be because PROM is a risk factor for early-onset neonatal sepsis as well as a high risk of fetal distress, respiratory distress syndrome, intraventricular hemorrhage, and death [31,32,54]. In this study we found an overall prevalence of PROM of 8.1%, which is in accordance with the global incidence that ranges from approximately 5% to 10% of all deliveries in the world [54,55].

Lack of emergency obstetric care (EmOC) is well-known to increase the risk of neonatal mortality, as laboring mothers with complications cannot immediately receive appropriate health services, such as access to a cesarean section [36]. In this study, mothers who needed to be transferred from another health unit for delivery at HAM had a sixfold higher risk of having a neonatal death outcome than mothers directly admitted at HAM. Half the mothers who were transferred from the death-outcome group had to do a 60 -kilometer journey that took approximately two to three hours until reaching HAM maternity in the capital city. Distance from a health facility with EmOC and the considerable travel times are well-known barriers and influential factors of birth adverse outcomes also found in other studies in sub-Saharan Africa [56,57].

In this study, the overall rate of 3.1% of instrumental assisted vaginal birth deliveries was found to be lower compared to 17% of births by cesarean section and 79.9% of normal vaginal deliveries. Instrumental-assisted deliveries are an effective intervention for deliveries complicated by prolonged labor or fetal distress but are also related to adverse outcomes since fetal distress or cephalopelvic disproportion are more frequently the motive to use this technique. Therefore, the ninefold higher risk of death outcome found in this study, similar to other researchers reporting [13,58,59], relates to late appropriate intrapartum care intervention with subsequently higher rates of fetal distress and risk of an intrapartum stillbirth.

Antenatal care health services in STP follow a high-risk stratification for each pregnancy and, in this study, this notification of a high-risk pregnancy in the pregnant women antenatal pregnancy card was significantly associated with a threefold higher risk for fetal and neonatal death than alive-outcome group. High-pregnancy risk was operationally defined as one or more of the following according to STP national practice, namely, age as inferior to 15 years old or superior to 35, grand multipara, previous history of a stillbirth or early neonatal death, previous cesarean section, and previous hemorrhagic complication. This is consistent with other studies that reported higher rates of perinatal and neonatal mortality with extreme maternal ages and previous adverse obstetric history (grand multipara, previous stillbirth, or early neonatal death) [1,12]. However, in this study, we could not find a statistically significant difference between groups when each of these variables were independently analyzed.

Regarding the newborns' characteristics, the overall sex ratio at birth was 52.6% for males and 47.4% for females. The death rate was higher in male neonates (72%) than in the female death-outcome group (27.3%), with female sex being a protective factor. Sex variations are

frequently reported, with a highly consistent pattern of excess male mortality across different populations and income groups [60]. Male infants are more vulnerable to fetal and early neonatal death, with most studies reporting no gender difference in mortality after 7 days of age [60]. This sex difference may be explained by a sex-specific difference in the growth and function of male and female placentae, making boys more vulnerable to different adverse outcomes [60,62].

Birth weights of more than 2500 g were protective factors for death outcome in this study. This finding is comparable with different research on other African countries that associates a birth weight of less than 2500 g (LBW) with higher rates of stillbirth and neonatal death [1]. Additionally, the odds of experiencing death were 21 times higher for babies with IUGR or fetal growth restriction defined, in this study, as a term LBW (birthweight <2500 g and gestational age ≥ 37 weeks) [42]. Low birth weight is a worldwide recognized important multifaceted public health problem since LBW infants are 20 times more likely to develop complications and die than normal weight infants [6,63].

We found that the odds of death were six times higher for newborns with maternal infectious risk. This finding is in agreement with previous studies conducted in SSA, as neonatal infectious risk is linked to amniotic fluid contamination that can cause intrapartum fetal infection or early postpartum neonatal infection with sepsis complicated with septic shock and multiple organ dysfunction, in which both are the most common causes of death in the perinatal period [35,43]. Infection is an important cause of stillbirth and neonatal death in LMICs, although there is a lack of overall information regarding the organisms involved, the types of transmission, and the mechanisms of death in these constrained countries [64]. In Sao Tome & Principe, malaria and syphilis are no longer an infection-burden and cause of

579 stillbirth or neonatal death; therefore, other bacterial and viral maternal infections should be
580 linked to stillbirth and ENND [64]. In STP, there are no means to establish a neonatal
581 infection through blood cultures; that is, neonatal infection is based only on clinical signs
582 with no bacteriological documentation. Additionally, nothing is known about the rate of
583 vertical transmission of Streptococcus Group B (GBS) from colonized mothers at birth in the
584 country. Therefore, there is a current gap in the knowledge, capability of making diagnosis
585 and prevention and treatment of neonatal infections in STP. For instance, a study from
586 Ethiopia [64] detected that the rate of vertical transmission of GBS from colonized mothers at
587 birth was as high as 45.02% due to term PROM, PROM ≥ 18 hours before delivery and
588 mothers having fever during labor. Detecting the risk for vertical transmission of GBS and
589 implementing prevention methods such as adequate intrapartum administration of antibiotics
590 are recognized as easy and affordable practices for reducing mortality due to infectious
591 causes and should be implemented in the country along with culture techniques [43].

592 All four congenital major malformations detected in this study resulted in perinatal mortality,
593 although due to the low number of newborns enrolled it was not possible to establish a
594 statistically significant difference. Congenital malformations are known to be associated with
595 poor outcomes for newborns, and 10–20% of stillbirths are attributed to intrinsic fetal
596 anomalies [53]. Studies from Zimbabwe [65], Ethiopia [1] and Yemen [12] found that
597 congenital anomalies among stillbirths were associated with a 5-fold [65], 34-fold [1] and 40-
598 fold [12] higher risk than those among live births. Another important consideration regarding
599 congenital anomalies is the type of malformation, since some, such as neural tube defects, are
600 preventable and can be reduced by 30 to 50% by folic acid supplementation in pregnant
601 women and have been estimated to cause 29% of deaths related to congenital anomalies in
602 LMICs [4]. In this study, three out of the four major anomalies were neural tube defects,

highlighting the need to enhance proper supplementation to preconceptional women in the country.

In this study, we were not able to find a higher risk associated with newborn gestational age. The high rate of stillbirths found in our study as well as the low number of preterm babies enrolled (5.6%) explain the lack of association. Stillbirths are among the most common pregnancy-related adverse outcomes worldwide, but they differ between low- and high-income countries. In high-income countries, most stillbirths occur early in the preterm period, whereas in low resource constrained countries, most occur in term or in late preterm births, as found in our study [66]. Additionally, the neonatal mortality is higher in babies born between 37 and 38 weeks of gestation than in those born between 39 and weeks [4].

From stillbirths in this study, 69% occurred during the intrapartum period, in accordance with other studies from LMICs [5]. These intrapartum stillbirths mean that intrauterine death occurred after the onset of labor and before birth (fresh stillbirth). We can also guess that some of them could be early neonatal deaths, as there are known barriers and difficulties in these contexts to establish whether a fetus or motionless newborn is living or dead after its delivery. For instance, information on some Apgar scores in these death-outcome group was not recorded, perhaps because of lack of time, especially when the neonates had to be rushed to receive resuscitation maneuvers. Studies from SSA highlight that it is very frequent that some depressed but living fetuses with a possible heartbeat do not receive resuscitation maneuvers and are prompt classified as stillbirths [1,66]. This is supported by a systematic review of sixteen hospitals and community-based perinatal mortality studies [67,68].

In this study, newborns who needed resuscitation maneuvers were at a 16-fold higher risk of dying. Additionally, the odds of neonatal death from fetal distress at birth and birth asphyxia were 36 and 55 times higher, respectively. These results are in line with studies conducted in

Cameroon [69] and Ethiopia [36] as well as published literature that suggests that lower APGAR scores are associated with severe multiorgan damage resulting in brain damage, lung dysfunction, cardiomyopathy, renal failure, hepatic failure, necrotizing enterocolitis and consequently death [70]. Previous studies [23,24] that compared outcomes between adolescent pregnant girls and older counterparts identified that adverse outcomes imputable to adolescent births were fetal distress (OR 1.94, 95% CI 1.18-3.18) and performance of neonatal resuscitation maneuvers (OR 2.4, 95% CI 1.07-5.38), highlighting the risks surrounding this period among deliveries at HAM maternity unit.

Perinatal asphyxia can be caused by factors grouped according to whether they are before birth (antepartum risk factors), during birth (intrapartum risk factors), or after birth (postpartum or fetal risk factors) [44,70]. Nonetheless, the single most important predictor is, undoubtedly, the quality of intrapartum care during labor and delivery.

The most important aspect of this study for public health is that it identifies potential characteristics that predispose newborns to life-threatening conditions, which is critical to address the underlying causes and provide prompt interventions by various stakeholders in the healthcare system [71].

This study allows us to perceive how to answer our question “when, where and why do newborns die in STP?” as “when”: mostly during labor, “where”: *in uterus* and “why”: mainly due to fetal distress and intrapartum-related deaths probably due to low quality and constraints of care during labor and delivery. Ending preventable stillbirths and neonatal deaths does not necessarily require new or innovative interventions. Most modifiable risk factors identified in this study can be addressed and prevented with timely, quality care during childbirth, including ongoing intrapartum monitoring and opportune intervention in case of complications [72]. Some interventions, such as computerized cardiotocography to monitor a baby’s well-being in the womb by measuring contractions, are estimated to reduce

the rate of infant deaths around the time of birth by 80% when compared with traditional cardiotocography [73].

In summary, the findings of this study will be useful to health policymakers and program developers in implementing appropriate interventions to achieve the newborn health post-2015 Sustainable Development Goals of no more than 12 neonatal deaths per 1000 live births in Sao Tome & Principe by 2030 [74].

Strengths and Limitations

In this study, the researcher retrieved maternal and neonatal data directly from ANC cards and maternity registers to limit recall bias. The selection of the death-outcome group and alive-outcome group was based on the records of maternal and neonatal registers; therefore, it is less likely that this study has misclassification biases both in the exposure and death-outcome group □ alive-outcome group categories [75].

Regarding the limitations, this is a relatively small study aiming to identify factors associated with peri-neonatal mortality in Sao Tome & Principe with a cohort of 194 newborns that were followed up until 28th days of age, 22 died and 172 survived. Thus, this study results cannot be generalized.

The newborns lost to follow-up were mainly due to mothers who provided the wrong mobile number or who did not answer the phone call at a significantly high rate; hence, the real number of early and late neonatal deaths can be much higher than the rate we have found.

Current knowledge gaps, including those associated with local burden, bacterial etiology, and eventual difficulty of nurses and mothers in perceiving subtle clinical signs associated with neonatal infections prevented us from having a clear picture of the situation in this study [43].

Data on maternal weight gain during pregnancy, nutritional status, BMI, maternal intrapartum antibiotic prophylaxis, antenatal corticosteroid administration, and total time of labor were not available.

Another limitation is that some of the variables mentioned in the univariable model had wide confidence intervals and high odds ratios due to the low number of death-outcome group enrolled in this study. This was a barrier to conducting a multivariable model, a limitation also reported in other studies [37]. Due to the small number of neonatal death-outcome group enrolled and these wide confidence intervals, the multivariable logistic regression model could only identify one independent determinant of mortality outcome.

Nonetheless, the current study can assist Sao Tome & Principe policy makers and stakeholders in designing new policies for the country to improve maternal and neonatal health outcomes.

Conclusions

Complications such as a high-risk pregnancy score, meconium-stained amniotic fluid, prolonged rupture of membranes, being transferred from another unit, and an instrumental-assisted vaginal delivery increased the risk of stillbirth and neonatal death between 4- and 9-fold, and 90% of all these deaths occurred in the perinatal period.

Newborns with an infectious risk, intrauterine growth restriction, fetal distress at birth, who needed resuscitation maneuvers, birth asphyxia, and those admitted to the neonatal unit had a 3- to 55-fold higher risk for dying than the alive-outcome group. Female newborn and birth weight of more than 2500 g were found to be protective factors.

Implementing an efficient referral system that directs high-risk pregnancies to the EmOC facility, equipment (continuous fetal heart monitoring capability in all district maternity wards) and personnel technical skills will improve peri-neonatal outcomes and survival in Sao Tome & Principe.

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718 **Declarations**

719

720 **Ethics approval and consent to participate**

721 The Ministry of Health of Sao Tome & Principe and the main board of the Hospital Dr. Ayres
 722 de Menezes are dedicated ethics oversight bodies, and both approved this study. At the time
 723 of the study protocol was submitted (year 2016), there were no ethics committees in STP. All
 724 methods were performed in accordance with the relevant guidelines and regulations in
 725 practice. Written informed consent was obtained from all participants (or their parent or legal
 726 guardian in the case of adolescent under 16 or for illiterate pregnant women) after the
 727 purpose of the research was explained orally by the principal investigator. Participants' or
 728 their legal representatives' also consented to have the results of this research work published.
 729 Participation in the survey was voluntary, as participants could decline to participate at any
 730 time during the study.

731

732 **Consent for publication**

733 Not applicable.

734

735 **Availability of data and materials**

The datasets used and/or analyzed during the current study are all available within the manuscript itself.

Competing interests

No.

Funding

AV was supported by the Fundação para a Ciência e Tecnologia (FCT) (<https://www.fct.pt/index.phtml.pt/>), grant number SFRH/BD/117037/2016. The funder had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Abbreviations

ANC – antenatal care; CI – confidence interval; cOR: crude odds ratio; GA – gestational age; HAM: Hospital Dr. Ayres de Menezes; IPI – intestinal parasitic infection; IUGR – intrauterine growth restriction; PROM – prolonged rupture of membranes; NCU – Neonatal Care Unit

Authors' contributions

AV, MCM and FP carried out the conception and design of the study. AV was responsible for field activities, data collection, and wrote the manuscript. MCM and FP critically evaluated and made progressive suggestions throughout the study and revised the manuscript. MA and ALP performed statistical analysis and reviewed the manuscript. SS and NB were involved in the study design at the country level. All the authors read and approved the final draft of the manuscript.

Acknowledgments

A special remark for the late Professor João Luís Baptista MD PhD - AV research cosupervisor - a great man who was a thinker and a fighter for Africa's improvement of public health. We are indebted to all the women who participated in the study. The authors would like to thank Elizabeth Carvalho and the 1) medical team and nurses of Hospital Dr. Ayres de Menezes Maternity unit for their support, especially to the chief-nurse Paulina Oliveira; and 2) Ana Sequeira, Rita Coelho, Ana Margalha, Ana Castro, Alexandra Coelho,

and Inês Gomes for field support. We would like to acknowledge Instituto Camões, I.P. for logistic assistance in STP.

Authors' Information

AV is a Portuguese pediatrician with 14 years of short- and long- missions in Sao Tome & Principe, with a postgraduation in Tropical Medicine from the London Hospital of Tropical Medicine through the first East-African Course in Uganda and Tanzania and is about to finish her PhD thesis on neonatal mortality and morbidity in STP.

References

1. Getiye Y, Fantahun M. Factors associated with perinatal mortality among public health deliveries in Addis Ababa, Ethiopia, an unmatched case control study. BMC pregnancy and childbirth. 2017;17(1):1-7.
2. Pathirana J, Muñoz FM, Abbing-Karahagopian V, Bhat N, Harris T, Kapoor A, Keene DL, Mangili A, Padula MA, Pande SL, Pool V. Neonatal death: Case definition & guidelines for data collection, analysis, and presentation of immunization safety data. Vaccine. 2016;34(49):6027-37.
3. WHO. Every new born: an action plan to end preventable deaths. February 2014. Available at: [https://cdn.who.int/media/docs/default-source/mca-documents/advisory-groups/quality-of-care/every-new-born-action-plan-\(enap\).pdf?sfvrsn=4d7b389_2](https://cdn.who.int/media/docs/default-source/mca-documents/advisory-groups/quality-of-care/every-new-born-action-plan-(enap).pdf?sfvrsn=4d7b389_2)
4. Lehtonen L, Gimeno A, Parra-Llorca A, Vento M. Early neonatal death: A challenge worldwide. Semin Fetal Neonatal Med. 2017;22(3):153-160. doi: 10.1016/j.siny.2017.02.006. Epub 2017 Feb 24. PMID: 28238633.
5. Hug L, You D, Blencowe H, Mishra A, Wang Z, Fix MJ, Wakefield J, Moran AC, Gaigbe-Togbe V, Suzuki E, Blau DM. Global, regional, and national estimates and trends in stillbirths from 2000 to 2019: a systematic assessment. The Lancet. 2021 Aug 28;398(10302):772-85. [https://doi.org/10.1016/S0140-6736\(21\)01112-0](https://doi.org/10.1016/S0140-6736(21)01112-0)
6. Dagne HM, Melku AT, Abdi AA. Determinants of stillbirth among deliveries attended in Bale Zone Hospitals, Oromia Regional State, Southeast Ethiopia: a case-control study. International Journal of Women's Health. 2021;13:51.
7. Lawn JE, Kerber K, Enweronu-Laryea C, Massee Bateman O. Newborn survival in low resource settings—are we delivering?. BJOG: An International Journal of Obstetrics & Gynaecology. 2009;116:49-59.
8. Lawn JE, Cousens S, Zupan J, Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: when? Where? Why?. The lancet. 2005 Mar 5;365(9462):891-900.
9. Lawn JE, Blencowe H, Pattinson R, Cousens S, Kumar R, Ibiebele I, et al. Stillbirths: Where? When? Why? How to make the data count? Lancet. 2011;377(9775):1448–63.
10. Blencowe H, Cousens S. Addressing the challenge of neonatal mortality. Tropical Medicine & International Health. 2013;18(3):303-12.
11. Lavin T, Allanson ER, Nedkoff L, Preen DB, Pattinson RC. Applying the international classification of diseases to perinatal mortality data, South Africa. Bulletin of the World Health Organization. 2018 Dec 1;96(12):806.

812 12. Obad MA, Taher R, Qayad M, Khader YS. Risk factors of stillbirth in Yemen. J Neonatal
813 Perinatal Med. 2018;11(2):131–136. doi:10.3233/ NPM-18174618.

814 13. Adane, et al. Adverse birth outcomes among deliveries in Gonder university hospital,
815 Northwest Ethiopia. BMC Pregnancy Childbirth. 2014;14:90
816 <http://www.biomedcentral.com/1471-2393/14/90>

817 14. Chi BH, Wang L, Read JS, Taha TE, Sinkala M, Brown ER, Valentine M, Martinson F,
818 Goldenberg RL: Predictors of stillbirth in sub-saharan Africa. Obstet Gynecol. 2007, 110 (5):
819 989-997. 10.1097/01.AOG.0000281667.35113.a5.

820 15. Stringer EM, Vwalika B, Killam WP, Giganti MJ, Mbewe R, Chi BH, Chintu N, Rouse
821 D, Goldenberg RL, Stringer JS: Determinants of stillbirth in Zambia. Obstet Gynecol. 2011,
822 117 (5): 1151-1159. 10.1097/AOG.0b013e3182167627.

823 16. Sao Tome and Principe WHO statistical profile. WHO Libr. 2015. [https://www.who.int/](https://www.who.int/gho/countries/stp.pdf)
824 [gho/ countries/stp. pdf](https://www.who.int/gho/countries/stp.pdf).

825 17. INE e UNICEF. 2020. Inquérito de Indicadores Múltiplos 2019, Relatório final. São
826 Tomé, São Tomé e Príncipe: Instituto Nacional de Estatística e Fundo das Nações Unidas
827 para a Infância. Available at: [https://mics-surveys-](https://mics-surveys-prod.s3.amazonaws.com/MICS6/West%20and%20Central%20Africa/Sao%20Tome%20and%20Principe/2019/Survey%20findings/Sao%20Tome%20e%20Principe%202019%20MICS%20Survey%20Findings%20Report_Portuguese.pdf)
828 [prod.s3.amazonaws.com/MICS6/West%20and%20Central%20Africa/Sao%20Tome%20and](https://mics-surveys-prod.s3.amazonaws.com/MICS6/West%20and%20Central%20Africa/Sao%20Tome%20and%20Principe/2019/Survey%20findings/Sao%20Tome%20e%20Principe%202019%20MICS%20Survey%20Findings%20Report_Portuguese.pdf)
829 [%20Principe/2019/Survey%20findings/Sao%20Tome%20e%20Principe%202019%20MICS](https://mics-surveys-prod.s3.amazonaws.com/MICS6/West%20and%20Central%20Africa/Sao%20Tome%20and%20Principe/2019/Survey%20findings/Sao%20Tome%20e%20Principe%202019%20MICS%20Survey%20Findings%20Report_Portuguese.pdf)
830 [%20Survey%20Findings%20Report_Portuguese.pdf](https://mics-surveys-prod.s3.amazonaws.com/MICS6/West%20and%20Central%20Africa/Sao%20Tome%20and%20Principe/2019/Survey%20findings/Sao%20Tome%20e%20Principe%202019%20MICS%20Survey%20Findings%20Report_Portuguese.pdf)

831 18. UNICEF IN de E (INE) e. Inquérito de Indicadores Múltiplos, São Tome e Principe,
832 MICS-STP, 2014, Principais resultados. SaoTome, SaoTome e Principe, INE e UNICEF.
833 2015; Instituto Nacional de Estatística, 2016. Inquérito aos Indicadores Múltiplos 2014 de
834 São Tomé e Príncipe, Relatório Final. São Tomé, São Tomé e Príncipe.
835 [http:// ms. gov. st/ wp- conte nt/ uploads/ 2019/ 01/ MICS- Final- Report- STP_ Portu gu%](http://ms.gov.st/wp-content/uploads/2019/01/MICS-Final-Report-STP_Portugu%C3%AAs.Pdf)
836 [C3% AAs. Pdf](http://ms.gov.st/wp-content/uploads/2019/01/MICS-Final-Report-STP_Portugu%C3%AAs.Pdf)

837 19. Bhutta Z, Das JK, Bahl R, Lawn JE, Salam R, Paul VK, et al. Can available interventions
838 end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? Lancet.
839 2014;384(9940):347–70.

840 20. Beyond2015, Gcap, Ifp. Civil Society Demands for the Post-2015 Agenda from 39
841 Countries. 2013;

842 21. Dickson KE, Simen-Kapeu A, Kinney M V., Huicho L, Vesel L, Lackritz E, et al. Every
843 Newborn: Health-systems bottlenecks and strategies to accelerate scale-up in countries.
844 Lancet. 2014;384(9941):438–54.

- 845 22. Nair N, Tripathy P, Prost A, Costello A, Osrin D. Improving newborn survival in low-
846 income countries: Community-based approaches and lessons from South Asia. *PLoS Med.*
847 2010;7(4).
- 848 23. Vasconcelos A, Bandeira N, Sousa S, Pereira F, Machado MD. Adolescent pregnancy in
849 Sao Tome and Principe: a cross-sectional hospital-based study. *BMC Pregnancy and*
850 *Childbirth.* 2022 Dec;22(1):1-5.
- 851 24. Vasconcelos A, Bandeira N, Sousa S, Machado MC, Pereira F. Adolescent pregnancy in
852 Sao Tome and Principe: are there different obstetric and perinatal outcomes?. *BMC*
853 *Pregnancy and Childbirth.* 2022 Dec;22(1):1-1.
- 854 25. Vasconcelos A, Sousa S, Bandeira N, Baptista JL, Machado MD, Pereira F. PO 8592
855 Why, when and where do newborns not only get sick but also die in são tomé and príncipe?
856 A case-control study.
- 857 26. Ministério da Saúde_ República Democrática de Sao Tome e Principe_Plano Nacional do
858 Desenvolvimento da Saúde 2017-2021
- 859 27. Tavares Da Silva F, Gonik B, McMillan M, Keech C, Dellicour S, Bhange S, Tila M,
860 Harper DM, Woods C, Kawai AT, Kochhar S, Munoz FM; Brighton Collaboration Stillbirth
861 Working Group. Stillbirth: Case definition and guidelines for data collection, analysis, and
862 presentation of maternal immunization safety data. *Vaccine.* 2016;34(49):6057-6068. doi:
863 10.1016/j.vaccine.2016.03.044. Epub 2016 Jul 16. PMID: 27431422; PMCID: PMC5139804.
- 864 28. Kikuchi K, Ansah EK, Okawa S, Enuameh Y, Yasuoka J, Nanishi K, et al. Effective
865 linkages of continuum of care for improving neonatal, perinatal, and maternal mortality: a
866 systematic review and metaanalysis. *PloS one.* 2015; 10(9):e0139288. doi:
867 10.1371/journal.pone.0139288 PMID: 26422685
- 868 29. Mulowooza J, Santos N, Isabirye N et al. Midwife-performed checklist and ultrasound to
869 identify obstetric conditions at labour triage in Uganda: A quasi-experimental study.
870 *Midwifery.* 2021;96:102949.
- 871 30. Weissmann-Brenner A, Meyer R, Domniz N, Levin G, Hendin N, Yoeli-Ullman R,
872 Mazaki-Tovi S, Weissbach T, Kassif E. The perils of true knot of the umbilical cord:
873 Antepartum, intrapartum and postpartum complications and clinical implications. *Archives of*
874 *Gynecology and Obstetrics.* 2022;305(3):573-9.
- 875 31. Chiabi A, Takou V, Mah E, Nguefack S, Siyou H, Takou V, et al. Risk factors for
876 neonatal mortality at the Yaounde Gynaeco-Obstetric and Pediatric Hospital. *Cameroon.*
877 *Iranian Journal of Pediatrics.* 2014;24(4):393–400.

878 32. Tewabe T, Mohammed S, Tilahun Y et al. Clinical outcome and risk factors of neonatal
879 sepsis among neonates in Felege Hiwot referral Hospital, Bahir Dar, Amhara Regional State,
880 North West Ethiopia 2016: a retrospective chart review. BMC Res Notes. 2017;10, 265.
881 <https://doi.org/10.1186/s13104-017-2573-1>

882 33. Avagliano L, Massa V, Bulfamante G. Meconium-stained amniotic fluid and histologic
883 signs of fetal distress in stillbirths. European Journal of Obstetrics & Gynecology and
884 Reproductive Biology. 2021;266:55-62.

885 34. Shaikh S, Shaikh AH, Shaikh SA, Isran B. Frequency of obstructed labor in teenage
886 pregnancy. Nepal Journal of Obstetrics and Gynaecology. 2012;7(1):37-40.

887 35. Alemu AY, Belay GM, Berhanu M, Minuye B. Determinants of neonatal mortality at
888 neonatal intensive care unit in Northeast Ethiopia: unmatched case-control study. Tropical
889 Medicine and Health. 2020 Dec;48(1):1-0.

890 36. Yego F, D'Este C, Byles J, Nyongesa P, Williams JS. A case-control study of risk factors
891 for fetal and early neonatal deaths in a tertiary hospital in Kenya. BMC pregnancy and
892 childbirth. 2014 Dec;14(1):1-9.

893 37. Demisse AG, Alemu F, Gizaw MA, Tigabu Z. Patterns of admission and factors
894 associated with neonatal mortality among neonates admitted to the neonatal intensive care
895 unit of University of Gondar Hospital, Northwest Ethiopia. Pediatric health, medicine and
896 therapeutics. 2017;8:57.

897 38. Abadiga M, Mosisa G, Tsegaye R, et al. Determinants of adverse birth outcomes among
898 women delivered in public hospitals of Ethiopia, 2020. Archives of Public Health.
899 2022;80(1):1-7. <https://doi.org/10.1186/s13690-021-00776-0>

900 39. Pimentel J, Ansari U, Omer K, et al. Factors associated with short birth interval in low-
901 and middle-income countries: a systematic review. BMC Pregnancy Childbirth 2020;20,156.
902 <https://doi.org/10.1186/s12884-020-2852-z>

903 40. Esoh K, Wonkam-Tingang E, Wonkam A. Sickle cell disease in sub-Saharan Africa:
904 transferable strategies for prevention and care. The Lancet Haematology. 2021

905 41. Cutland CL, Lackritz EM, Mallett-Moore T, Bardaji A, Chandrasekaran R, Lahariya C,
906 Nisar MI, Tapia MD, Pathirana J, Kochhar S, Muñoz FM. Low birth weight: Case definition
907 & guidelines for data collection, analysis, and presentation of maternal immunization safety
908 data. Vaccine. 2017;35(48Part A):6492.

909 42. Accrombessi M, Zeitlin J, Massougboji A, Cot M, Briand V. What do we know about
910 risk factors for fetal growth restriction in Africa at the time of sustainable development goals?
911 A scoping review. Paediatric and perinatal epidemiology. 2018;32(2):184-96.

912 43. Huynh B, Kermorvant-Duchemin E, Herindrainy P et al. Bacterial Infections in Neonates,
913 Madagascar, 2012–2014. *Emerging Infectious Diseases*. 2018; 24(4), 710-717.
914 <https://doi.org/10.3201/eid2404.161977>

915 44. Wosenu L, Worku AG, Teshome DF, Gelagay AA. Determinants of birth asphyxia
916 among live birth newborns in University of Gondar referral hospital, northwest Ethiopia: A
917 case-control study. *PLoS ONE*. 2018;13(9): e0203763.

918 45. Mbachu, I, Achigbu, K, Odinaka, K, et al. Tracking stillbirths by referral pattern and
919 causes in a rural tertiary hospital in Southern Nigeria. *Niger Postgrad Med J* 2018; 25(2): 87–
920 93.

921 46. Chuwa, FS, Mwanamsangu, AH, Brown, BG, et al. Maternal and fetal risk factors for
922 stillbirth in Northern Tanzania: a registry-based retrospective cohort study. *PLoS ONE* 2017;
923 12(8): e0182250.

924 47. Mulatu T, Debella A, Feto T, Dessie Y. Determinants of stillbirth among women who
925 gave birth at Hiwot Fana Specialized University Hospital, Eastern Ethiopia: A facility-based
926 cross-sectional study. *SAGE Open Medicine*. 2022 Feb;10:20503121221076370.

927 48. Chaibva, et al. Adverse pregnancy outcomes, ‘stillbirths and early neonatal deaths’ in
928 Mutare district, Zimbabwe. *BMC Pregnancy Childbirth*. 2019;19:86.
929 <https://doi.org/10.1186/s12884-019-2229-3>

930 49. Lawn JE, Blencowe H, Waiswa P, et al, for The Lancet Ending Preventable Stillbirths
931 Series study group with The Lancet Stillbirth Epidemiology investigator group. Stillbirths:
932 rates, risk factors, and acceleration towards 2030. *Lancet* 2016;
933 [https://doi.org/10.1016/S0140-6736\(15\)00837-5](https://doi.org/10.1016/S0140-6736(15)00837-5)

934 50. Mamo SA, Teshome GS, Tesfaye T, Goshu AT. Perinatal asphyxia and associated factors
935 among neonates admitted to a specialized public hospital in South Central Ethiopia: A
936 retrospective cross-sectional study. *PloS one*. 2022;17(1):e0262619.

937 51. Desai D, Maitra N, Patel P. Fetal heart rate patterns in patients with thick meconium
938 staining of amniotic fluid and its association with perinatal outcome. *Int J Reprod Contracept*
939 *Obstet Gynecol*. 2017;6(3):1030–5

940 52. Asiki, et al. Adverse pregnancy outcomes in rural Uganda (1996–2013): trends and
941 associated factors from serial cross-sectional surveys. *BMC Pregnancy Childbirth*.
942 2015;15:279. <https://doi.org/10.1186/s12884-015-0708-8>

943 53. Lawn J, Shibuya K, Stein C. No cry at birth: global estimates of intrapartum stillbirths
944 and intrapartum-related neonatal deaths. *Bull World Health Organ*. 2005;83:409–17

945 54. Assefa, N., Berhe, H., Girma, F. et al. Risk factors of premature rupture of membranes in
946 public hospitals at Mekele city, Tigray, a case control study. BMC Pregnancy Childbirth.
947 2018; 18, 386 <https://doi.org/10.1186/s12884-018-2016-6>

948 55. Alam MM, Saleem AF, Shaikh AS, Munir O, Qadir M. Neonatal sepsis following
949 prolonged rupture of membranes in a tertiary care hospital in Karachi, Pakistan. The Journal
950 of Infection in Developing Countries. 2014;8(01):067-73.

951 56. Geleto, A., Chojenta, C., Musa, A. et al. Barriers to access and utilization of emergency
952 obstetric care at health facilities in sub-Saharan Africa: a systematic review of literature. Syst
953 Rev. 2018; 7, 183. <https://doi.org/10.1186/s13643-018-0842-2>

954 57. Habte A, Wondimu M. Determinants of maternal near miss among women admitted to
955 maternity wards of tertiary hospitals in Southern Ethiopia, 2020: A hospital-based case-
956 control study. PloS one, 2021. 16(5): p. e0251826.
957 <https://doi.org/10.1371/journal.pone.0251826>

958 58. Thakur N, Sunny AK, Gurung R, Basnet O, Litorp H, Ashish KC. Rate and Neonatal
959 Outcomes Among Instrument Assisted Vaginal Birth in 12 Public Hospitals in Nepal.
960 Preprint. DOI: <https://doi.org/10.21203/rs.3.rs-36775/v1>

961 59. Hailemichael HT, Debelew GT, Alema HB, Weldu MG, Misgina KH. Determinants of
962 adverse birth outcome in Tigray region, North Ethiopia: Hospital-based case-control study.
963 BMC pediatrics. 2020;20(1):1-9. <https://doi.org/10.1186/s12887-019-1835-6>

964 60. Mondal D, Galloway TS, Bailey TC, Mathews F. Elevated risk of stillbirth in males:
965 systematic review and meta-analysis of more than 30 million births. BMC medicine.
966 2014;12(1):1-1.

967 61. Aghai ZH, Goudar SS, Patel A, Saleem S, Dhaded SM, Kavi A, Lalakia P, Naqvi F,
968 Hibberd PL, McClure EM, Nolen TL. Gender variations in neonatal and early infant
969 mortality in India and Pakistan: a secondary analysis from the Global Network Maternal
970 Newborn Health Registry. Reproductive Health. 2020;17(3):1-1.
971 <https://doi.org/10.1186/s12978-020-01028-0>

972 62. Pongou R. Why is infant mortality higher in boys than in girls? A new hypothesis based
973 on preconception environment and evidence from a large sample of twins. Demography.
974 2013;50(2):421-44.

975 63. KC A, Basel PL, Singh S. Low birth weight and its associated risk factors: Health
976 facility-based case-control study. PloS one. 2020;15(6):e0234907.

977 64. Yadeta TA, Worku A, Egata G, Seyoum B, Marami D, Berhane Y. Vertical transmission
978 of group B Streptococcus and associated factors among pregnant women: a cross-sectional
979 study, Eastern Ethiopia. *Infection and drug resistance*. 2018;11:397.

980 65. El T, Gombe N, Shambira G, Chadambuka A, Mufuta T, Zizhou S. Determinants of
981 perinatal mortality in Marondera district, Mashonal and East Province of Zimbabwe.2009.
982 *Pan Afr Med J*. 2011;8(7).

983 66. Goldenberg RL, Harrison MS, McClure EM. Stillbirths: the hidden birth asphyxia—US
984 and Global perspectives. *Clinics in perinatology*. 2016;43(3):439-53.

985 67. Gutman A, Harty T, O'Donoghue K, Greene R, Leitao S. Perinatal mortality audits and
986 reporting of perinatal deaths: systematic review of outcomes and barriers. *Journal of Perinatal*
987 *Medicine*. 2022;26. <https://doi.org/10.1515/jpm-2021-0363>

988 68. Berhan Y, Berhan A. Perinatal mortality trends in Ethiopia. *Ethiopian journal of health*
989 *sciences*. 2014;24:29-40.

990 69. Chiabi A, Takou V, Mah E, Nguefack S, Siyou H, Takou V, et al. Risk factors for
991 neonatal mortality at the Yaounde Gynaeco-Obstetric and Pediatric Hospital. Cameroon.
992 *Iranian Journal of Pediatrics*. 2014;24(4):393–400.

993 70. American College of Obstetricians and Gynecologists. The Apgar score (Committee
994 Opinion No. 644). *Obstet Gynecol*. 2015;126(4):52–5.

995 71. Habte A, Lukas K, Melis T, Tamene A, Sahle T, Hailu M, et al. (2022) Determinants of
996 neonatal near miss among neonates admitted to public hospitals in Southern Ethiopia, 2021:
997 A case-control study. *PLoS ONE* 17(5): e0268041.
998 <https://doi.org/10.1371/journal.pone.0268041>

999 72. Hug L, You D, Blencowe H, Mishra A, Wang Z, Fix MJ, Wakefield J, Moran AC,
1000 Gaigbe-Togbe V, Suzuki E, Blau DM. Global, regional, and national estimates and trends in
1001 stillbirths from 2000 to 2019: a systematic assessment. *The Lancet*. 2021 Aug
1002 28;398(10302):772-85. [https://doi.org/10.1016/S0140-6736\(21\)01112-0](https://doi.org/10.1016/S0140-6736(21)01112-0)

1003 73. Ota, E, da Silva Lopes, K, Middleton, P, Flenady, V, Wariki, WM, Rahman, MO, et al..
1004 Antenatal interventions for preventing stillbirth, fetal loss and perinatal death: an overview of
1005 Cochrane systematic reviews. *Cochrane Database Syst Rev* 2020;12:CD009599.
1006 <https://doi.org/10.1002/14651858.CD009599.pub2>.

1007 74. Paulson KR, Kamath AM, Alam T, Bienhoff K, Abady GG, Abbas J, Abbasi-Kangevari
1008 M, Abbastabar H, Abd-Allah F, Abd-Elsalam SM, Abdoli A. Global, regional, and national
1009 progress towards Sustainable Development Goal 3.2 for neonatal and child health: all-cause

1010 and cause-specific mortality findings from the Global Burden of Disease Study 2019. The
 1011 Lancet. 2021;398(10303):870-905. [https://doi.org/10.1016/S0140-6736\(21\)01207-1](https://doi.org/10.1016/S0140-6736(21)01207-1)
 1012 75. KC A, Basel PL, Singh S. Low birth weight and its associated risk factors: Health
 1013 facility-based case-control study. PloS one. 2020;15(6):e0234907.
 1014 <https://doi.org/10.1371/journal.pone.0234907>

