

Study or Subgroup	logIRR	SE	Weight	Incidence Rate Ratio IV, Random, 95% CI	Incidence Rate Ratio IV, Random, 95% CI
-------------------	--------	----	--------	---	---

year = 2021

klompmaker	-0.0269	0.0083	8.2%	0.97 [0.96; 0.99]
russette	-0.0862	0.0284	7.4%	0.92 [0.87; 0.97]
russette	-0.1079	0.0298	7.3%	0.90 [0.85; 0.95]
russette	-0.2291	0.0357	7.0%	0.80 [0.74; 0.85]
Total (95% CI)			29.8%	0.90 [0.78; 1.03]

Heterogeneity: $\text{Tau}^2 = 0.0063$; $\text{Chi}^2 = 37.96$, $\text{df} = 3$ ($P < 0.01$); $I^2 = 92\%$

year = 2023

sikarwar	-0.0737	0.0573	5.6%	0.93 [0.83; 1.04]
sikarwar	-0.1701	0.0604	5.4%	0.84 [0.75; 0.95]
sikarwar	-0.2147	0.0613	5.3%	0.81 [0.71; 0.91]
sikarwar	-0.2765	0.0636	5.2%	0.76 [0.67; 0.86]
Total (95% CI)			21.5%	0.83 [0.73; 0.96]

Heterogeneity: $\text{Tau}^2 = 0.0038$; $\text{Chi}^2 = 6.1$, $\text{df} = 3$ ($P = 0.11$); $I^2 = 51\%$

year = 2022

yang	0.0069	0.0065	8.2%	1.01 [0.99; 1.02]
yang	-0.0132	0.0008	8.2%	0.99 [0.99; 1.00]
yang	0.0103	0.0039	8.2%	1.01 [1.01; 1.03]
yang	0.0245	0.0232	7.6%	1.02 [0.95; 1.04]
yang	-0.0353	0.0083	8.2%	0.97 [0.95; 0.98]
yang	-0.0419	0.0071	8.2%	0.96 [0.95; 0.97]
Total (95% CI)			48.7%	0.99 [0.96; 1.02]

Heterogeneity: $\text{Tau}^2 = 0.0005$; $\text{Chi}^2 = 70.53$, $\text{df} = 5$ ($P < 0.01$); $I^2 = 93\%$

Total (95% CI) **100.0%** **0.93 [0.88; 0.98]**

Prediction interval **[0.77; 1.12]**

Heterogeneity: $\text{Tau}^2 = 0.0069$; $\text{Chi}^2 = 163.02$, $\text{df} = 13$ ($P < 0.01$); $I^2 = 92\%$

Test for subgroup differences: $\text{Chi}^2 = 19.10$, $\text{df} = 2$ ($P < 0.01$)

