

Metabolic syndrome and hemorrhagic stroke among symptomatic CCMs in the Mexican Hispanic Population

Alok K. Dwivedi¹, David Jang¹, Ofek Belkin¹, Justin Aickareth¹, Mellisa Renteria¹,
Majd Hawwar¹, Croft, Jacob¹, Ammar M Kalas², Marc Zuckerman², Jun Zhang^{1*}

Departments of ¹Molecular and translational Medicine and ²Internal Medicine

Texas Tech University Health Sciences Center, El Paso, TX, USA

Supplemental Materials

*All correspondence:

Jun Zhang, Sc.D., Ph.D.

Molecular and translational Medicine

Texas Tech University Health Sciences Center El Paso

5001 El Paso Dr., El Paso, TX 79905

Tel: (915) 215-4197

Email: jun.zhang2000@gmail.com

Legends

Suppl Table 1. Variables of interest for data collection of CCM patients. All clinical data from the patients admitted to TTUHSCEP/UMC presenting with confirmed or assumed cerebral cavernous malformations (CCMs) will be collected based on variables of interest.

Suppl. Table 2. An interaction between age and metabolic syndrome (MetS) on hemorrhagic stroke among Mexican -Hispanic CCM patients

Suppl. Table 3. Distribution of MetS, epilepsy, and hemorrhagic stroke by age groups among Hispanic CCM patients

Suppl Table 1. Variables of interest for data collection of CCM patients. All clinical data from the patients admitted to TTUHSCEP/UMC presenting with confirmed or assumed cerebral cavernous malformations (CCMs) and MetS parameters will be collected based on variables of interest listed below. (Note: not all risk factors were used in this study).

Age years (EX: 32)

Date of first-ever ICH (if occurred) (year)

Date of the first-ever seizure (if occurred) (years)

Time point of any treatment (neurosurgery, radiosurgery) (if occurred) (year)

Confirmed familial disease (MRI pattern plus one affected relative or pos. genotyping) (yes/no)

Assumed familial disease (MRI pattern without an affected relative/genotyping) (yes/no)

Time point of the last follow-up at the hospital/institution (Year)

Confirmed mutation (CCM1/2/3) (if available)

Functional state (mRS) at last FU (if available)

Stroke Type (Ischemic 1 Hemorrhagic 2)

Place of Birth (Mexico, El Paso, UNK)

Admit Date (Year only)

Hypertension (1, yes; 2, no)

Hypertension numbers

Systolic BP at presentation

Diastolic BP at presentation

Mean Arterial Pressure number

Tobacco abuse (1, yes; 2, no)

Tobacco Info (packs/day)

Diabetes(1, yes; 2, no)

Diabetes Info (I/II; glucose Value)

Heart Disease (1, yes; 2, no)

Suppl. Table 2

Parameters	OR	95%CI		p-value
MetS	0.01	0.00	1.43	0.071
Age-years	1.00	0.98	1.02	0.904
MetS and age	1.09	1.01	1.18	0.029
MeTS	0.26	0.03	2.26	0.223
Age \geq 50	0.89	0.45	1.73	0.722
MetS and Age \geq 50	18.98	1.66	216.70	0.018

MetS: metabolic syndrome; OR: odds ratio; CI: confidence interval

Suppl. Table 3

Parameters	Age \leq 35	Age (35-50)	Age \geq 50	p-value
Sample size (N)	33	44	107	
Metabolic syndrome				0.38
No	31 (93.9%)	37 (86.0%)	86 (83.5%)	
Yes	2 (6.1%)	6 (14.0%)	17 (16.5%)	
Epilepsy				0.77
No	24 (72.7%)	35 (79.5%)	79 (74.5%)	
Yes	9 (27.3%)	9 (20.5%)	27 (25.5%)	
Stroke				0.60
No	23 (69.7%)	29 (65.9%)	64 (60.4%)	
Yes	10 (30.3%)	15 (34.1%)	42 (39.6%)	
Hemorrhagic Stroke	OR, p-value	OR, p-value	OR, p-value	
MetS-Yes	0.41, 0.575	0.44, 0.405	4.67, 0.006	
Epilepsy-Yes	0.65, 0.606	1.01, 0.987	2.37, 0.054	

OR: odds ratio.