

Gene Symbol (n=55)	Gene MIM #	Protein	Related Disease	Phenotype MIM #	Mode of Inheritance
<i>ADCY10</i>¹	605205	Adenylate cyclase 10	Hypercalciuria, absorptive, susceptibility to	143870	AD
<i>AGXT</i>²	604285	Alanine--glyoxylate and serine-- pyruvate aminotransferase	Hyperoxaluria, primary, type I	259900	AR
<i>ALDOB</i>³	612724	Aldolase, fructose-bisphosphate B	Fructose intolerance, hereditary	229600	AR
<i>ALPL</i>⁴	171760	Alkaline phosphatase	Hypophosphatasia, adult	146300	AD, AR
			Hypophosphatasia, child	241510	AR
			Hypophosphatasia infantile	241500	AR
<i>APRT</i>⁵	102600	Adenine phosphoribosyltransferase	Adenine phosphoribosyltransferase deficiency, Urolithiasis (DHA stones), renal failure	614723	AR
<i>ATP6V0A4</i>⁶	605239	ATPase H+ transporting V0 subunit a4	Distal renal tubular acidosis 3, with or without progressive sensorineural hearing loss	602722	AR
<i>ATP6V1B1</i>⁷	192132	ATPase H+ transporting V1 subunit B1	Distal renal tubular acidosis 2 with progressive sensorineural hearing loss	267300	AR
<i>ATP7B</i>⁸	606882	ATPase copper transporting beta	Wilson Disease	277900	AR
<i>BSND</i>⁹	606412	Barttin CLCNK type accessory subunit beta	Bartter syndrome, type 4a, neonatal, with sensorineural deafness	602522	AR
<i>CA2</i>¹⁰	611492	Carbonic anhydrase 2	Osteopetrosis, autosomal recessive 3, with renal tubular acidosis	259730	AR
<i>CASR</i>¹¹	601199	Calcium sensing receptor	Hyperparathyroidism, neonatal	239200	AD, AR
			Hypocalcemia, autosomal dominant	601198	AD
			Hypocalcemia, autosomal dominant, with Bartter syndrome	601198	AD
			Hypocalciuric hypercalcemia, type I	145980	AD
			Epilepsy idiopathic generalized, susceptibility to, 8	612899	N/A
<i>CLCN5</i>¹²	300008	Chloride voltage-gated channel 5	Dent Disease 1	300009	XLR
			Hypophosphatemic rickets	300554	XLR
			Nephrolithiasis, type I	310486	XLR
			Proteinuria, low molecular weight, with hypercalciuric nephrocalcinosis	308990	XLR
<i>CLCNKA</i>¹³	602024	Chloride voltage-gated channel Ka	Bartter syndrome, type 4b, digenic	613090	DR
<i>CLCNKB</i>¹⁴	602023	Chloride voltage-gated channel Kb	Bartter syndrome, type 4b, digenic	613090	DR

			Bartter syndrome, type 3	607364	AR
CLDN16 ¹⁵	603959	Claudin 16	Hypomagnesemia 3, renal	248250	AR
CLDN19 ¹⁶	610036	Claudin 19	Hypomagnesemia 5, renal, with ocular involvement	248190	AR
CTNS ¹⁷	606272	Cystinosis, lysosomal cystine transporter	Cytinosis, typical/atypical nephropathic	219800	AR
			Cystinosis, late-onset juvenile or adolescent nephropathic	219900	AR
CYP24A1 ¹⁸	126065	Cytochrome P450 family 24 subfamily A member 1	Hypercalcemia, infantile, 1	143880	AR
FAH ¹⁹	613871	Fumarylacetoacetate hydrolase	Tyrosinemia, type I	276700	AR
FAM20A ²⁰	611062	FAM20A golgi associated secretory pathway pseudokinase	Amelogenesis imperfecta, type IG (enamel-renal syndrome)	204690	AR
FOXI1 ²¹	601093	Forkhead box I1	Enlarged vestibular aqueduct	600791	AR
G6PC ²²	613742	Glucose-6-phosphatase catalytic subunit	Glycogen storage disease Ia	232200	AR
GNA11 ²³	139313	G protein subunit alpha 11	Hypocalcemia, autosomal dominant 2	615361	AD
			Hypocalciuric hypercalcemia, type II	145981	AD
GPHN ²⁴	603930	Gephyrin	Molybdenum cofactor deficiency C	615501	AR
GRHRP ²⁵	604269	Glyoxylate and hydroxypyruvate reductase	Hyperoxaluria, primary, type II	260000	AR
HNF4A ²⁶	600281	Hepatocyte nuclear factor 4 alpha	Fanconi renotubular syndrome 4, with maturity-onset diabetes of the young	616026	AD
HOGA1 ²⁷	613597	4-hydroxy-2-oxoglutarate aldolase 1	Hyperoxaluria, primary, type III	613616	AR
HPRT1 ²⁸	308000	Hypoxanthine phosphoribosyltransferase 1	Kelley-Seegmiller syndrome, partial HPRT deficiency, HPRT related gout	300323	XLR
			Lesch-Nyhan syndrome	300322	XLR
KCNJ1 ²⁹	600359	Potassium inwardly rectifying channel subfamily J member 1	Bartter syndrome, type 2	241200	AR
KCNJ10 ³⁰	602208	Potassium inwardly rectifying channel subfamily J member 10	SESAME syndrome	612780	AR
KL ³¹	604824	Klotho	Tumoral calcinosis, hyperphosphatemic, familial, 3	617994	AR
LRP2 ³²	600073	LDL receptor related protein 2	Donnai-Barrow syndrome	222448	AR
MAGED2 ³³	300470	MAGE family member D2	Bartter syndrome, type 5, antenatal, transient	300971	XLR
MGP ³⁴	154870	Matrix Gla protein	Keutel syndrome	245150	AR
MOCOS ³⁵	613274	Molybdenum cofactor sulfuryase	Xanthinuria, type II	603592	AR

<i>MOCS1</i> ³⁶	603707	Molybdenum cofactor synthesis 1	Molybdenum cofactor deficiency A	252150	AR
<i>MOCS2</i> ³⁶	603708	Molybdenum cofactor synthesis 2	Molybdenum cofactor deficiency B	252160	AR
<i>OCRL</i> ³⁷	300535	OCRL inositol polyphosphate-5-phosphatase	Dent disease 2	300555	XLR
			Lowe syndrome	309000	XLR
<i>ORAI1</i> ³⁸	610277	ORAI calcium release-activated calcium modulator 1	Immunodeficiency 9	612782	AR
			Myopathy, tubular aggregate, 2	615883	AD
<i>PRPS1</i> ³⁹	311850	Phosphoribosyl pyrophosphate synthetase 1	Phosphoribosylpyrophosphate synthetase superactivity	300661	XLR
<i>SLC12A1</i> ²⁹	600839	Solute carrier family 12 member 1	Barter syndrome, type 1	601678	AR
<i>SLC22A12</i> ⁴⁰	607096	Solute carrier family 22 member 12	Hypouricemia, renal (RHUC1)	220150	AR, AD
<i>SLC26A1</i> ⁴¹	610130	Solute carrier family 26 member 1	Nephrolithiasis, calcium oxalate	167030	AR
<i>SLC2A2</i> ⁴²	138160	Solute carrier family 2 member 2	Fanconi-Bickel Syndrome	227810	AR
<i>SLC2A9</i> ⁴³	606142	Solute carrier family 2 member 9	Renal hypouricemia, RHUC2	612076	AR, AD
<i>SLC34A1</i> ⁴⁴	182309	Solute carrier family 34 member 1	Hypercalcemia, infantile, 2	616963	AR
			Nephrolithiasis/osteoporosis, hypophosphatemic, 1	612286	AD
			Fanconi renotubular syndrome 2	613388	AR
			Hypercalcemia, infantile, 2	616963	AR
<i>SLC34A3</i> ⁴⁵	609826	Solute carrier family 34 member 3	Hypophosphatemic rickets with hypercalciuria	241530	AR
<i>SLC3A1</i> ⁴⁶	104614	Solute carrier family 3 member 1	Cystinuria	220100	AD, AR
<i>SLC4A1</i> ⁴⁷	109270	Solute carrier family 4 member 1	Distal renal tubular acidosis 1	179800	AD
			Distal renal tubular acidosis 4 with hemolytic anemia	611590	AR
<i>SLC5A1</i> ⁴⁸	82380	Solute carrier family 5 member 1	Glucose/galactose malabsorption	606824	AR
<i>SLC7A9</i> ⁴⁹	604144	Solute carrier family 7 member 9	Cystinuria	220100	AD, AR
<i>SLC9A3R1</i> ⁵⁰	604990	SLC9A3 regulator 1	Hypophosphatemic nephrolithiasis/osteoporosis 2, (NPHLOP2)	612287	AD
<i>VDR</i> ⁵¹	601769	Vitamin D receptor	Rickets, vitamin D-resistant, type IIA; Idiopathic hypercalcuria	277440	AR
<i>WDR72</i> ⁵²	613214	WD repeat domain 72	Amelogenesis imperfecta, type IIA3	613211	AR
<i>XDH</i> ⁵³	607633	Xanthine dehydrogenase	Xanthinuria, type I	278300	AR

Supplemental Table 1. List of genes known to cause monogenic kidney stone disease along with their corresponding MIM number, protein, related disorder, and mode of inheritance.

1. Reed BY, Gitomer WL, Heller HJ, et al. Identification and characterization of a gene with base substitutions associated with the absorptive hypercalciuria phenotype and low spinal bone density. *J Clin Endocrinol Metab.* 2002;87(4):1476-1485. doi:10.1210/jcem.87.4.8300
2. Purdue PE, Allsop J, Isaya G, Rosenberg LE, Danpure CJ. Mistargeting of peroxisomal L-alanine:glyoxylate aminotransferase to mitochondria in primary hyperoxaluria patients depends upon activation of a cryptic mitochondrial targeting sequence by a point mutation. *Proc Natl Acad Sci U S A.* 1991;88(23):10900-10904. doi:10.1073/pnas.88.23.10900
3. Paoletta G, Santamaria R, Buono P, Salvatore F. Mapping of a restriction fragment length polymorphism within the human aldolase B gene. *Hum Genet.* 1987;77(2):115-117. doi:10.1007/BF00272375
4. Whyte MP, Rettinger SD, Vrabel LA. Infantile hypophosphatasia: enzymatic defect explored with alkaline phosphatase-deficient skin fibroblasts in culture. *Calcif Tissue Int.* 1987;40(5):244-252. doi:10.1007/BF02555256
5. Kelley WN, Levy RI, Rosenbloom FM, Henderson JF, Seegmiller JE. Adenine phosphoribosyltransferase deficiency: a previously undescribed genetic defect in man. *J Clin Invest.* 1968;47(10):2281-2289. doi:10.1172/JCI105913
6. Smith AN, Skaug J, Choate KA, et al. Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. *Nat Genet.* 2000;26(1):71-75. doi:10.1038/79208
7. Karet FE, Finberg KE, Nelson RD, et al. Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat Genet.* 1999;21(1):84-90. doi:10.1038/5022
8. Gromadzka G, Schmidt HHJ, Genschel J, et al. Frameshift and nonsense mutations in the gene for ATPase7B are associated with severe impairment of copper metabolism and with an early clinical manifestation of Wilson's disease. *Clin Genet.* 2005;68(6):524-532. doi:10.1111/j.1399-0004.2005.00528.x
9. Brennan TM, Landau D, Shalev H, et al. Linkage of infantile Bartter syndrome with sensorineural deafness to chromosome 1p. *Am J Hum Genet.* 1998;62(2):355-361. doi:10.1086/301708
10. Sly WS, Hewett-Emmett D, Whyte MP, Yu YS, Tashian RE. Carbonic anhydrase II deficiency identified as the primary defect in the autosomal recessive syndrome of osteopetrosis with renal tubular acidosis and cerebral calcification. *Proc Natl Acad Sci U S A.* 1983;80(9):2752-2756. doi:10.1073/pnas.80.9.2752
11. Pearce SH, Williamson C, Kifor O, et al. A familial syndrome of hypocalcemia with hypercalciuria due to mutations in the calcium-sensing receptor. *N Engl J Med.* 1996;335(15):1115-1122. doi:10.1056/NEJM199610103351505
12. Lloyd SE, Pearce SHS, Fisher SE, et al. A common molecular basis for three inherited kidney stone diseases. *Nature.* 1996;379(6564):445-449. doi:10.1038/379445a0

13. Schlingmann KP, Konrad M, Jeck N, et al. Salt wasting and deafness resulting from mutations in two chloride channels. *N Engl J Med*. 2004;350(13):1314-1319. doi:10.1056/NEJMoa032843
14. Simon DB, Bindra RS, Mansfield TA, et al. Mutations in the chloride channel gene, CLCNKB, cause Bartter's syndrome type III. *Nat Genet*. 1997;17(2):171-178. doi:10.1038/ng1097-171
15. Simon DB, Lu Y, Choate KA, et al. Paracellin-1, a renal tight junction protein required for paracellular Mg²⁺ resorption. *Science*. 1999;285(5424):103-106. doi:10.1126/science.285.5424.103
16. Konrad M, Schaller A, Seelow D, et al. Mutations in the tight-junction gene claudin 19 (CLDN19) are associated with renal magnesium wasting, renal failure, and severe ocular involvement. *Am J Hum Genet*. 2006;79(5):949-957. doi:10.1086/508617
17. Town M, Jean G, Cherqui S, et al. A novel gene encoding an integral membrane protein is mutated in nephropathic cystinosis. *Nat Genet*. 1998;18(4):319-324. doi:10.1038/ng0498-319
18. Schlingmann KP, Kaufmann M, Weber S, et al. Mutations in CYP24A1 and idiopathic infantile hypercalcemia. *N Engl J Med*. 2011;365(5):410-421. doi:10.1056/NEJMoa1103864
19. Aponte JL, Segal GA, Hauser LJ, et al. Point mutations in the murine fumarylacetoacetate hydrolase gene: Animal models for the human genetic disorder hereditary tyrosinemia type 1. *Proc Natl Acad Sci U S A*. 2001;98(2):641-645.
20. Jaureguiberry G, De la Dure-Molla M, Parry D, et al. Nephrocalcinosis (enamel renal syndrome) caused by autosomal recessive FAM20A mutations. *Nephron Physiol*. 2012;122(1-2):1-6. doi:10.1159/000349989
21. Yang T, Vidarsson H, Rodrigo-Blomqvist S, Rosengren SS, Enerback S, Smith RJH. Transcriptional control of SLC26A4 is involved in Pendred syndrome and nonsyndromic enlargement of vestibular aqueduct (DFNB4). *Am J Hum Genet*. 2007;80(6):1055-1063. doi:10.1086/518314
22. Seydewitz HH, Matern D. Molecular genetic analysis of 40 patients with glycogen storage disease type Ia: 100% mutation detection rate and 5 novel mutations. *Hum Mutat*. 2000;15(1):115-116. doi:10.1002/(SICI)1098-1004(200001)15:1<115::AID-HUMU23>3.0.CO;2-W
23. Li D, Opas EE, Tuluc F, et al. Autosomal dominant hypoparathyroidism caused by germline mutation in GNA11: phenotypic and molecular characterization. *J Clin Endocrinol Metab*. 2014;99(9):E1774-1783. doi:10.1210/jc.2014-1029
24. Reiss J, Gross-Hardt S, Christensen E, Schmidt P, Mendel RR, Schwarz G. A mutation in the gene for the neurotransmitter receptor-clustering protein gephyrin causes a novel form of molybdenum cofactor deficiency. *Am J Hum Genet*. 2001;68(1):208-213. doi:10.1086/316941
25. Williams HE, Smith LH. L-glyceric aciduria. A new genetic variant of primary hyperoxaluria. *N Engl J Med*. 1968;278(5):233-238. doi:10.1056/NEJM196802012780502
26. Hamilton AJ, Bingham C, McDonald TJ, et al. The HNF4A R76W mutation causes atypical dominant Fanconi syndrome in addition to a β cell phenotype. *J Med Genet*. 2014;51(3):165-169. doi:10.1136/jmedgenet-2013-102066

27. Belostotsky R, Seboun E, Idelson GH, et al. Mutations in DHPSL are responsible for primary hyperoxaluria type III. *Am J Hum Genet.* 2010;87(3):392-399. doi:10.1016/j.ajhg.2010.07.023
28. Davidson BL, Tarlé SA, Van Antwerp M, et al. Identification of 17 independent mutations responsible for human hypoxanthine-guanine phosphoribosyltransferase (HPRT) deficiency. *Am J Hum Genet.* 1991;48(5):951-958.
29. Simon DB, Karet FE, Rodriguez-Soriano J, et al. Genetic heterogeneity of Bartter's syndrome revealed by mutations in the K⁺ channel, ROMK. *Nat Genet.* 1996;14(2):152-156. doi:10.1038/ng1096-152
30. Bockenhauer D, Feather S, Stanescu HC, et al. Epilepsy, ataxia, sensorineural deafness, tubulopathy, and KCNJ10 mutations. *N Engl J Med.* 2009;360(19):1960-1970. doi:10.1056/NEJMoa0810276
31. Ichikawa S, Imel EA, Kreiter ML, et al. A homozygous missense mutation in human KLOTTHO causes severe tumoral calcinosis. *J Clin Invest.* 2007;117(9):2684-2691. doi:10.1172/JCI31330
32. Kantarci S, Al-Gazali L, Hill RS, et al. Mutations in LRP2, which encodes the multiligand receptor megalin, cause Donnai-Barrow and facio-oculo-acoustico-renal syndromes. *Nat Genet.* 2007;39(8):957-959. doi:10.1038/ng2063
33. Laghmani K, Beck BB, Yang SS, et al. Polyhydramnios, Transient Antenatal Bartter's Syndrome, and MAGED2 Mutations. *N Engl J Med.* 2016;374(19):1853-1863. doi:10.1056/NEJMoa1507629
34. Munroe PB, Olgunturk RO, Fryns JP, et al. Mutations in the gene encoding the human matrix Gla protein cause Keutel syndrome. *Nat Genet.* 1999;21(1):142-144. doi:10.1038/5102
35. Ichida K, Matsumura T, Sakuma R, Hosoya T, Nishino T. Mutation of human molybdenum cofactor sulfurase gene is responsible for classical xanthinuria type II. *Biochem Biophys Res Commun.* 2001;282(5):1194-1200. doi:10.1006/bbrc.2001.4719
36. Shalata A, Mandel H, Reiss J, et al. Localization of a gene for molybdenum cofactor deficiency, on the short arm of chromosome 6, by homozygosity mapping. *Am J Hum Genet.* 1998;63(1):148-154. doi:10.1086/301916
37. Attree O, Olivos IM, Okabe I, et al. The Lowe's oculocerebrorenal syndrome gene encodes a protein highly homologous to inositol polyphosphate-5-phosphatase. *Nature.* 1992;358(6383):239-242. doi:10.1038/358239a0
38. Feske S, Gwack Y, Prakriya M, et al. A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature.* 2006;441(7090):179-185. doi:10.1038/nature04702
39. Becker MA, Losman MJ, Wilson J, Simmonds HA. Superactivity of human phosphoribosyl pyrophosphate synthetase due to altered regulation by nucleotide inhibitors and inorganic phosphate. *Biochim Biophys Acta.* 1986;882(2):168-176. doi:10.1016/0304-4165(86)90151-0
40. Enomoto A, Kimura H, Chairoungdua A, et al. Molecular identification of a renal urate anion exchanger that regulates blood urate levels. *Nature.* 2002;417(6887):447-452. doi:10.1038/nature742

41. Gee HY, Jun I, Braun DA, et al. Mutations in SLC26A1 Cause Nephrolithiasis. *Am J Hum Genet.* 2016;98(6):1228-1234. doi:10.1016/j.ajhg.2016.03.026
42. Akagi M, Inui K, Nakajima S, et al. Mutation analysis of two Japanese patients with Fanconi-Bickel syndrome. *J Hum Genet.* 2000;45(1):60-62. doi:10.1007/s100380050013
43. Matsuo H, Chiba T, Nagamori S, et al. Mutations in glucose transporter 9 gene SLC2A9 cause renal hypouricemia. *Am J Hum Genet.* 2008;83(6):744-751. doi:10.1016/j.ajhg.2008.11.001
44. Prié D, Beck L, Friedlander G, Silve C. Sodium-phosphate cotransporters, nephrolithiasis and bone demineralization. *Curr Opin Nephrol Hypertens.* 2004;13(6):675-681. doi:10.1097/00041552-200411000-00015
45. Lorenz-Depiereux B, Benet-Pages A, Eckstein G, et al. Hereditary hypophosphatemic rickets with hypercalciuria is caused by mutations in the sodium-phosphate cotransporter gene SLC34A3. *Am J Hum Genet.* 2006;78(2):193-201. doi:10.1086/499410
46. Calonge MJ, Gasparini P, Chillarón J, et al. Cystinuria caused by mutations in rBAT, a gene involved in the transport of cystine. *Nat Genet.* 1994;6(4):420-425. doi:10.1038/ng0494-420
47. Bruce LJ, Cope DL, Jones GK, et al. Familial distal renal tubular acidosis is associated with mutations in the red cell anion exchanger (Band 3, AE1) gene. *J Clin Invest.* 1997;100(7):1693-1707. doi:10.1172/JCI119694
48. Turk E, Zabel B, Mundlos S, Dyer J, Wright EM. Glucose/galactose malabsorption caused by a defect in the Na⁺/glucose cotransporter. *Nature.* 1991;350(6316):354-356. doi:10.1038/350354a0
49. Feliubadaló L, Font M, Purroy J, et al. Non-type I cystinuria caused by mutations in SLC7A9, encoding a subunit (bo,+AT) of rBAT. *Nat Genet.* 1999;23(1):52-57. doi:10.1038/12652
50. Karim Z, Gérard B, Bakouh N, et al. NHERF1 mutations and responsiveness of renal parathyroid hormone. *N Engl J Med.* 2008;359(11):1128-1135. doi:10.1056/NEJMoa0802836
51. Scott P, Ouimet D, Valiquette L, et al. Suggestive evidence for a susceptibility gene near the vitamin D receptor locus in idiopathic calcium stone formation. *J Am Soc Nephrol.* 1999;10(5):1007-1013. doi:10.1681/ASN.V1051007
52. El-Sayed W, Parry DA, Shore RC, et al. Mutations in the beta propeller WDR72 cause autosomal-recessive hypomaturation amelogenesis imperfecta. *Am J Hum Genet.* 2009;85(5):699-705. doi:10.1016/j.ajhg.2009.09.014
53. Dent CE, Philpot GR. Xanthinuria, an inborn error (or deviation) of metabolism. *Lancet.* 1954;266(6804):182-185. doi:10.1016/s0140-6736(54)91257-x