

Supplement - Table 2 - Type of healthcare provider initially contacted for low back pain - Full Cohort

		% of Episodes			RR and 95% CI Compared to Pre	
		Pre-COVID	Early COVID	Late COVID	Early	Late
Commercial Insurance (CI)						
Episodes		150761	158123	124958		
Non-Rx	DC	23.6%	24.6%	24.1%	1.04 (1.03, 1.05)	1.02 (1.01, 1.03)
	PT	3.7%	3.1%	3.4%	0.86 (0.83, 0.89)	0.94 (0.90, 0.97)
	LAc	0.3%	0.2%	0.2%	0.66 (0.57, 0.77)	0.72 (0.61, 0.84)
Primary Care	PCP	23.6%	24.7%	21.7%	1.05 (1.03, 1.06)	0.92 (0.91, 0.93)
	Nurse	6.7%	6.4%	5.9%	0.96 (0.93, 0.98)	0.89 (0.87, 0.92)
	PA	5.0%	4.6%	4.7%	0.92 (0.89, 0.95)	0.95 (0.92, 0.98)
	DO	0.2%	0.2%	0.2%	0.98 (0.84, 1.15)	0.98 (0.83, 1.16)
Physician Specialist	OS	7.2%	7.4%	8.2%	1.03 (1.01, 1.06)	1.14 (1.11, 1.17)
	PM	4.4%	3.2%	2.4%	0.72 (0.69, 0.75)	0.54 (0.52, 0.56)
	PMR	3.8%	3.2%	3.0%	0.84 (0.81, 0.87)	0.80 (0.77, 0.83)
	Neuro	0.8%	0.7%	0.6%	0.88 (0.81, 0.95)	0.70 (0.64, 0.76)
	Rheum	0.8%	0.8%	0.6%	0.94 (0.87, 1.02)	0.76 (0.69, 0.83)
	NS	0.4%	0.4%	0.4%	1.01 (0.91, 1.12)	0.94 (0.83, 1.05)
	Oth	1.9%	2.2%	1.9%	1.13 (1.07, 1.19)	1.00 (0.95, 1.05)
Emergency Medicine/ Urgent Care	Hosp	8.4%	8.3%	10.2%	1.00 (0.97, 1.02)	1.22 (1.19, 1.25)
	Rad	6.8%	7.6%	9.5%	1.11 (1.08, 1.14)	1.40 (1.36, 1.43)
	EM	1.8%	1.9%	2.2%	1.03 (0.98, 1.09)	1.17 (1.11, 1.23)
	UC	0.6%	0.6%	0.7%	1.02 (0.93, 1.12)	1.14 (1.04, 1.26)
Medicare Advantage (MA)						
Episodes		530583	493573	368842		
Non-Rx	DC	7.7%	7.4%	8.4%	0.97 (0.95, 0.98)	1.09 (1.08, 1.11)
	PT	2.7%	2.3%	3.0%	0.87 (0.85, 0.89)	1.12 (1.09, 1.15)
	LAc	0.1%	0.1%	0.1%	0.96 (0.83, 1.12)	1.69 (1.47, 1.95)
Primary Care	PCP	36.3%	39.7%	34.2%	1.09 (1.09, 1.10)	0.94 (0.94, 0.95)
	Nurse	7.8%	7.8%	6.8%	1.00 (0.99, 1.01)	0.87 (0.85, 0.88)
	PA	4.9%	4.7%	4.5%	0.94 (0.92, 0.96)	0.92 (0.90, 0.93)
	DO	0.1%	0.1%	0.1%	0.99 (0.89, 1.11)	1.04 (0.92, 1.17)
Physician Specialist	OS	4.8%	5.0%	5.9%	1.05 (1.03, 1.07)	1.24 (1.22, 1.27)
	PM	8.6%	5.7%	3.8%	0.66 (0.65, 0.67)	0.44 (0.43, 0.45)
	PMR	3.5%	2.7%	2.4%	0.76 (0.74, 0.77)	0.67 (0.65, 0.69)
	Neuro	0.9%	0.9%	0.6%	0.96 (0.92, 1.00)	0.71 (0.68, 0.75)
	Rheum	0.9%	0.9%	0.8%	0.98 (0.94, 1.02)	0.81 (0.77, 0.84)
	NS	0.4%	0.4%	0.3%	0.89 (0.84, 0.95)	0.80 (0.75, 0.86)
	Oth	2.1%	2.3%	2.1%	1.13 (1.11, 1.16)	1.03 (1.00, 1.06)
Emergency Medicine/ Urgent Care	Hosp	10.6%	10.9%	14.6%	1.03 (1.02, 1.04)	1.38 (1.37, 1.40)
	Rad	7.3%	7.9%	10.6%	1.08 (1.07, 1.10)	1.46 (1.44, 1.48)
	EM	1.1%	1.1%	1.4%	0.96 (0.92, 0.99)	1.27 (1.22, 1.32)
	UC	0.3%	0.3%	0.4%	1.01 (0.94, 1.09)	1.31 (1.21, 1.41)

Pre=Pre-COVID period, Early=first 12 months post-COVID, Late=13-24 months post-COVID

PCP=primary care physician, PA=physician assistant, DO=doctor of osteopathy, DC=doctor of chiropractic, PT=physical therapist, LAc=licensed acupuncturist, OS=orthopedic surgeon, PMR=physical medicine and rehabilitation, PM=pain management, Neuro=neurologist, Rheum=rheumatologist, NS=neurosurgeon, MD-Oth=other physician specialists, Hosp=hospital, Rad=radiologist, EM=emergency medicine, UC=urgent care

For %, cells with red text indicate the type of HCP initially contacted was not significantly different from pre-COVID period (Fisher's Exact p > .05)

For %, cells with black text indicate the type of HCP initially contacted was significantly different from pre-COVID period (Fisher's Exact p > .05)

For risk ratio, cells in red indicate the confidence interval includes 1 and likelihood of HCP being initially contacted is not different than the pre-COVID reference