

Supplement - Table 3 - Type of healthcare services provided for low back pain - Full Cohort

		% of Episodes			RR and 95% CI Compared to Pre	
		Pre-COVID	Early COVID	Late COVID	Early	Late
Commercial Insurance (CI)						
Episodes - At Least 1 Service		150705	158069	124913		
First Line	Manipulation - Chiropractic	27.6%	28.2%	26.9%	1.02 (1.01, 1.03)	0.98 (0.96, 0.99)
	Active Care	26.5%	25.5%	25.9%	0.96 (0.95, 0.97)	0.98 (0.97, 0.99)
	Manual Therapy	17.8%	16.4%	16.3%	0.92 (0.91, 0.94)	0.92 (0.90, 0.93)
	Passive Therapy	15.8%	14.8%	13.5%	0.94 (0.92, 0.95)	0.86 (0.84, 0.87)
	Acupuncture	0.8%	0.6%	0.6%	0.70 (0.65, 0.77)	0.71 (0.65, 0.77)
	Manipulation - Osteopathic	0.8%	0.7%	0.7%	0.94 (0.87, 1.02)	0.84 (0.77, 0.92)
Second Line	Imaging - Radiography	36.8%	37.4%	41.1%	1.02 (1.01, 1.03)	1.12 (1.11, 1.13)
	Rx - NSAID	31.9%	30.5%	27.8%	0.96 (0.95, 0.97)	0.87 (0.86, 0.88)
	Rx - Muscle Relaxant	24.7%	23.7%	21.5%	0.96 (0.95, 0.97)	0.87 (0.86, 0.88)
	Imaging - MRI	17.1%	17.5%	18.1%	1.02 (1.01, 1.04)	1.06 (1.04, 1.08)
	Rx Oral Steroid	13.7%	12.1%	10.5%	0.88 (0.86, 0.89)	0.77 (0.75, 0.78)
	Rx Gabapentin	4.2%	3.6%	2.9%	0.86 (0.83, 0.89)	0.69 (0.66, 0.72)
	Rx - Other	15.8%	14.1%	11.2%	0.89 (0.88, 0.91)	0.71 (0.69, 0.72)
Third Line	Rx Opioid	21.0%	17.1%	13.3%	0.81 (0.80, 0.82)	0.63 (0.62, 0.65)
	Spinal Injection	13.7%	12.4%	10.8%	0.90 (0.89, 0.92)	0.78 (0.77, 0.80)
	Imaging - CT	3.4%	3.3%	3.4%	0.97 (0.94, 1.01)	1.00 (0.96, 1.04)
	Surgery	2.7%	2.5%	1.8%	0.93 (0.89, 0.98)	0.67 (0.64, 0.71)
Medicare Advantage (MA)						
Episodes - At Least 1 Service		530269	493163	368380		
First Line	Manipulation - Chiropractic	11.1%	11.0%	11.5%	1.00 (0.98, 1.01)	1.04 (1.03, 1.05)
	Active Care	16.2%	16.1%	17.7%	0.99 (0.98, 1.00)	1.10 (1.09, 1.11)
	Manual Therapy	8.2%	7.9%	8.6%	0.98 (0.96, 0.99)	1.05 (1.04, 1.07)
	Passive Therapy	3.1%	2.8%	2.7%	0.90 (0.88, 0.92)	0.87 (0.84, 0.89)
	Acupuncture	0.4%	0.4%	0.5%	1.15 (1.08, 1.22)	1.42 (1.33, 1.51)
	Manipulation - Osteopathic	0.4%	0.4%	0.4%	0.95 (0.89, 1.01)	0.87 (0.81, 0.93)
Second Line	Imaging - Radiography	33.7%	34.4%	38.5%	1.02 (1.01, 1.02)	1.14 (1.14, 1.15)
	Rx - NSAID	32.4%	30.4%	25.9%	0.94 (0.93, 0.94)	0.80 (0.79, 0.81)
	Rx - Muscle Relaxant	21.5%	19.6%	16.6%	0.91 (0.90, 0.92)	0.77 (0.77, 0.78)
	Imaging - MRI	21.1%	22.2%	23.6%	1.05 (1.04, 1.06)	1.12 (1.11, 1.13)
	Rx Oral Steroid	10.4%	8.9%	7.7%	0.86 (0.85, 0.87)	0.74 (0.73, 0.75)
	Rx Gabapentin	5.0%	4.4%	3.0%	0.87 (0.86, 0.89)	0.60 (0.58, 0.61)
	Rx - Other	26.9%	26.8%	20.3%	1.00 (0.99, 1.00)	0.76 (0.75, 0.76)
Third Line	Rx Opioid	40.5%	32.6%	24.1%	0.80 (0.80, 0.81)	0.59 (0.59, 0.60)
	Spinal Injection	21.6%	19.7%	17.1%	0.91 (0.90, 0.92)	0.79 (0.79, 0.80)
	Imaging - CT	7.9%	8.0%	8.1%	1.01 (1.00, 1.03)	1.03 (1.02, 1.05)
	Surgery	2.9%	2.7%	2.0%	0.93 (0.91, 0.95)	0.67 (0.66, 0.69)

Pre=Pre-COVID period, Early=first 12 months post-COVID, Late=13-24 months post-COVID

For %, cells with red text indicate the type of HCP initially contacted was not significantly different from pre-COVID period (Fisher's Exact p > .05)

For %, cells with black text indicate the type of HCP initially contacted was significantly different from pre-COVID period (Fisher's Exact p > .05)

For risk ratio, cells in red indicate the confidence interval includes 1 and likelihood of HCP being initially contacted is not different than the pre-COVID reference