| 1  | Title   |
|----|---|
|    |   |
| 2  | • Machine Learning Classifier Models for Predicting Sarcopenia in the Elderly Based |
| 3  | on Physical Factors   |
| 4  |   |
| 5  | Name of Author (Academic degree)  |
| 6  | • Emmett Kim, PhD, PT (ORCID: 0000-0002-2385-6843)                                  |
| 7  |   |
| 8  | Affiliation   |
| 9  | DataAid Research Team   |
| 10 |   |
| 11 | Corresponding Author  |
| 12 | • Emmett Kim, PhD, PT   |
| 13 |   |
| 14 | E-mail  |
| 15 | • <u>emetk2310@gmail.com</u>  |
| 16 |   |
| 17 | Telephone   |
| 18 | <ul> <li>+82-2-670-3379</li> </ul>  |
| 19 |   |

#### ABSTRACT

21

20

22 Background: As the elderly population gradually increases, musculoskeletal disorders such as 23 sarcopenia are increasing. Diagnostic techniques such as X-ray, CT, and MRI imaging are used to 24 predict and diagnose sarcopenia, and methods using machine learning are gradually increasing. 25 **Purpose:** The purpose of this study was to create a model that can predict sarcopenia using physical 26 characteristics and activity-related variables without medical diagnostic equipment such as imaging 27 equipment for the elderly aged 60 years or older. 28 Method: A sarcopenia prediction model was constructed using public data obtained from the Korea 29 National Health and Nutrition Examination Survey. Models were built using the multi-layer 30 perceptron, XGBoost, LightGBM, and RandomForest algorithms, and the feature importance of the 31 model with the highest accuracy was analyzed through evaluation metrics. 32 *Result*: The sarcopenia prediction model built with the LightGBM algorithm showed the highest test 33 accuracy at 0.852. In constructing the LightGBM model, physical characteristics variables such as 34 BMI showed high importance, and activity-related variables were also used in constructing the model. 35 *Conclusion*: The sarcopenia prediction model composed only of physical characteristics and activity-36 related factors showed excellent performance, and the use of this model will help predict sarcopenia 37 in the elderly living in communities with insufficient medical resources or difficult access to medical 38 facilities. 39 40 Keywords: Sarcopenia, predictive model, machine learning, physical characteristics, physical activity

41

## 43 I. INTRODUCTION

44 A decrease in skeletal muscle and consequent decrease in muscle strength can be seen as a 45 representative physical change in the elderly.(1) The amount of skeletal muscle decreases by about 8% 46 every 10 years (annual average of 0.8%) from the age of 40, and after the age of 70, more rapid 47 muscle loss occurs, and it appears to decrease by about 15% for 10 years.(1) As such, rapid changes in 48 body composition and body functions, such as muscle loss due to aging, can be defined as 49 sarcopenia.(1) The prevalence of sarcopenia shows a tendency to gradually increase with increasing 50 age.(2) In Korea, 13.0% of males and 21.7% of females in the middle-aged (40 to 59 years) had 51 sarcopenia, and 21.6% of males and 30.7% of females in the elderly (60 or older) had sarcopenia.(2)

52 The clinical significance of sarcopenia can be summarized as a decrease in muscle strength due to a 53 decrease in muscle mass, and an accompanying increase in physical disability and mortality. 54 Sarcopenia has been reported to be closely related to physical dysfunction, and in particular, problems 55 such as decreased gait function or increased risk of falling have been reported.(3,4) In addition, 56 sarcopenia was often accompanied by diseases such as metabolic dysfunction or geriatric chronic 57 diseases, and was also considered to be the cause of these diseases.(5,6) Therefore, sarcopenia due to 58 aging directly causes muscle strength deterioration, leading to deterioration of physical function, and 59 may cause or accompany chronic disabilities, and may increase the risk of death in the elderly.(7)

60 According to the definition of sarcopenia, measurement of absolute muscle mass is the best 61 assessment indicator for diagnosing sarcopenia.(8) Therefore, examination using imaging measures 62 such as MRI or CT to measure absolute muscle mass is currently considered the gold standard for 63 diagnosing sarcopenia.(9) These two imaging techniques show the highest accuracy and 64 reproducibility, and based on high image resolution, it is possible to selectively measure only muscles 65 excluding fat among tissues composing the body.(10) However, imaging using MRI or CT is 66 expensive and requires skilled operator experience and skill.(10) In addition, patients require high 67 compliance with MRI or CT imaging, and in the case of CT, patients may be exposed to high levels of 68 radiation.(10) For this reason, in clinical practice, sarcopenia is diagnosed using dual-energy X-ray

69 absorptiometry (DXA), which has a low radiation exposure and is relatively easy to measure (9,10)70 DXA can measure body composition by classifying it into bone mineral content, body fat mass, and 71 lean body mass, and can measure it separately for each limb and trunk.(10) The sum of the muscle 72 mass excluding the bone mineral mass from the lean body mass of both limbs is called the 73 appendicular skeletal muscle mass (ASM) and is used as a major indicator in the diagnosis of 74 sarcopenia.(11) If even these diagnostic methods are difficult to apply, sarcopenia can be diagnosed 75 with bioelectrical impedance analysis (BIA).(12) In addition to diagnosing sarcopenia using these 76 devices, attempts have been made to prevent or improve sarcopenia by more simply and easily 77 diagnosing sarcopenia through functional tests such as grip strength and walking speed.(13,14) 78 However, studies diagnosing sarcopenia using simple functional tests, such as grip strength, appear 79 differently depending on the race of subjects, or the methods of measuring functional factors used in 80 studies have not yet been standardized or consistent.(15,16)

81 Recently, to overcome the limitations of assessments for diagnosing sarcopenia, methods for 82 diagnosis through models using machine learning have been studied and presented. Machine learning 83 is a technique that can predict a specific value or classification by using a large amount of data 84 containing various variables.(17) A machine learning model that diagnoses image data through 85 clinical measurement equipment instead of an expert is being built to improve the accuracy of 86 diagnosing sarcopenia, and it is also possible to predict sarcopenia using imaging data from abdomen 87 or chest rather than limbs.(18–20) In addition, it is possible to diagnose sarcopenia or predict the risk 88 by modeling variables that can be related, such as physical characteristics, nutrition, and blood-related 89 variables, using machine learning, rather than analyzing only a few variables such as muscle strength 90 or walking speed.(21,22) These studies suggested that the prediction of sarcopenia using a machine 91 learning model would be more efficient in terms of time and cost than the traditional method of 92 diagnosing sarcopenia using imaging measurements.(19,21)

93 Although these predictors are said to be efficient in terms of time and cost, to measure these94 variables, medical devices or precise examinations by experts are required. This suggests that it may

95 be difficult to perform assessments according to the elderly living in underdeveloped facilities with 96 poor access to medical facilities or the rapidly growing elderly population.(8) Therefore, a model that 97 can be used as a screening or predict the risk of sarcopenia before using medical equipment or 98 performing a detailed examination by an expert to diagnose sarcopenia in the elderly will be needed. 99 Such a model should be composed of variables that are related to sarcopenia and are easy to measure, 100 such as physical characteristics and activities.(23,24)

101 Therefore, the purpose of this study was to create a predictive model for diagnosing sarcopenia 102 based on easily measurable variables such as physical characteristics and activity-related variables. To 103 build this predictive model, the following steps were required. First, it was to find out which variables 104 are needed to build models among variables related to physical characteristics and activities. Second, 105 it was to build several models with selected variables and improve performance through 106 hyperparameter tuning. Third, it was to compare various models and select the model with the best 107 performance among them. Fourth, it was to find out which variables had the highest importance in 108 constructing the best performing model.

### 110 II. MATERIALS AND METHODS

## 111 1. Data Source

The flow of this research process is shown in Figure 1. Data from the Korea National Health and Nutrition Examination Survey (KNHANES) conducted by the Korea Disease Control and Prevention Agency from 2008 to 2011 were used in this study. A total of 37,753 samples were investigated during that period. Among the data samples, there were a total of 5,573 data samples with dual energy x-ray data, age 60 years or older, and no missing responses to physical characteristics and activity-related variables. Data samples used in the study were provided with permission from the Korea Disease Control and Prevention Agency as data with personal identification information deleted.



119

120 Figure 1. Flowchart of the study

121

## 123 2. Data Preprocessing

124 This study used variables reflecting physical characteristics such as age, sex, height, weight and 125 BMI, and variables reflecting the degree of physical activity such as working hours or exercise 126 participation time per week. Table. 1 shows the variables and descriptions of the data samples used in 127 this study. Categorical variables were quantified through LabelEncoder, and numerical variables were 128 standardized using StandardScaler. To avoid overfitting and poor performance due to imbalanced data, 129 SMOTE was used to randomly generate data samples to match the proportions between the data. 130 LASSO algorithm was used to select features to use the multi-layer perceptron (MLP) model. And 131 RFECV (Recursive Feature Elimination with Cross-Validation) was used for feature selection of 132 XGBoost, LightGBM, and RandomForest models. Finally, before training the models, the entire 133 dataset was divided into a training dataset and a test dataset at a ratio of 8:2.

| KNHANES   | Description                         | Sarcopenia             | Normal                  |
|-----------|-------------------------------------|------------------------|-------------------------|
| age       | Age                                 | 70.76 <u>+</u> 6.17    | 68.54 <u>±</u> 5.84     |
| sex       | Gender                              | M:941, F:656           | M:1467, F:2509          |
| HE_ht     | Height                              | 159.03 <u>+</u> 8.85   | 157.03 <u>+</u> 8.89    |
| HE_wt     | Weight                              | 54.09 <u>+</u> 8.31    | 61.39 <u>+</u> 9.95     |
| HE_wc     | Waist circumference                 | 78.88 <u>+</u> 8.61    | 86.02 <u>+</u> 8.87     |
| BMI       | Body mass index                     | 21.34 <u>+</u> 2.41    | 24.81 <u>+</u> 2.94     |
| BO1_1     | Weight change                       | N:1170, I:353,<br>D:74 | N:2980, I:676,<br>D:320 |
| EQ5D      | Quality of life                     | 0.86 0.18              | 0.87 0.17               |
| EC_wht_23 | Average working hours per week      | 16.45 24.77            | 18.53 25.5              |
| BE3_11    | Vigorous physical activity per week | 1.58 1.63              | 1.76 1.78               |
| BE3_21    | Moderate physical activity per week | 2.12 2.17              | 2.56 2.45               |
| BE3_31    | Walking days per week               | 5.33 2.82              | 5.27 2.79               |
| LQ4_00    | Activity restriction                | Y:537, N:1060          | Y:1217, N:2759          |
| EC1_1     | Occupational activity               | Y:605, N:992           | Y:1704, N:2272          |

Table 1. Description of variables

134

### 136 *3. Machine learning algorithms*

137 A sarcopenia prediction model was developed using four machine learning algorithms: MLP, 138 XGBoost, LightGBM, and RandomForest. This study used a MLP model with one input layer, one 139 output layer and five hidden layers with varying numbers of nodes. The ReLU activation function was 140 used for the hidden layers, while the sigmoid activation function was used for the output layer. 141 Dropout ratio and batch normalization were applied to prevent overfitting, and the Adam optimizer 142 was used for gradient descent. The data was split with a 0.2 validation ratio, and EarlyStopping was 143 used to terminate learning when the validation loss did not improve for 20 epochs. The other three 144 models used the StratifiedKFold for cross-validation to ensure unbiased performance. This technique 145 partitions the data into K folds and ensures that each fold contains approximately equal proportions of 146 samples from each class. This allows the model to be trained and evaluated on a representative subset 147 of the data to obtain a more accurate performance estimate. In this study, a 5-fold training data set was 148 created for each model, and the model was built by repeating training and validation five times. The 149 hyperparameters of each model were then tuned using the GridSearchCV technique.

150

## 151 4. Model evaluation

The model's performance was assessed by comparing its predicted and actual classes using a confusion matrix that includes TP (true positive), FP (false positive), FN (false negative), and TN (true negative) values, based on test data. The accuracy, precision, recall, and F1-score were calculated from the matrix to evaluate the models. In addition, models were compared using the ROC AUC curve. Through the feature importance of the model with the highest accuracy, important factors in generating the model were analyzed.

### 159 III. RESULTS

- 160 1. Feature selection
- 161 All the mean test scores of the models according to variable selection using RFECV and LASSO
- 162 exceeded 0.75. The variables selected and excluded for each algorithm to build the predictive models
- 163 for sarcopenia diagnosis, and the mean test scores are shown in Table 2.

|                           | Mean test score | Selected variables   | Excluded variables             |
|---------------------------|-----------------|--|--------------------------------|
| Multi-layer<br>Perceptron | 0.773           | age, sex, LQ4_00, EQ5D,<br>EC_wht_23, BE3_11, BE3_21,<br>HE_wt, HE_wc, HE_BMI (10)                                 | EC1_1, BO1_1,<br>BE3_31, HE_ht |
| XGBoost                   | 0.835           | age, sex, EQ5D, EC_wht_23,<br>BO1_1, BE3_11, BE3_21, BE3_31,<br>HE_wt, HE_wc, HE_BMI (11)                          | LQ4_00, EC1_1,<br>HE_ht        |
| LightGBM                  | 0.834           | age, sex, EQ5D, EC_wht_23,<br>BE3_11, BE3_21, BE3_31, HE_ht,<br>HE_wt, HE_wc, HE_BMI (11)                          | LQ4_00, EC1_1,<br>BO1_1        |
| RandomForest              | 0.847           | age, sex, LQ4_00, EQ5D, EC1_1,<br>EC_wht_23, BO1_1, BE3_11,<br>BE3_21, BE3_31, HE_ht, HE_wt,<br>HE_wc, HE_BMI (14) | -                              |

Table 2. Feature selection and mean test scores through LASSO and RFECV

164

#### 165 *2. Model performance*

166 Figure 2 shows the confusion matrix created with test data to evaluate the performance of models 167 built with MLP, XGBoost, LightGBM, and RandomForest. Based on confusion matrix, the MLP 168 model showed the lowest performance with an accuracy of 0.79 when evaluated with test data. The 169 model with the highest performance was the model composed of the LightGBM algorithm. The test 170 accuracy of this model was 0.852, showing slightly higher performance than models using XGBoost 171 and RandomForest algorithms. Table 3 shows the accuracy, precision, recall, and F1-score of the 172 models using the multi-layer perceptron, XGBoost, LightGBM, and RandomForest algorithms. The 173 models compared through the ROC-AUC curve showed similar high performance except for the MLP 174 model (Figure 3).



Figure 2. Confusion matrix of sarcopenia prediction model

## 176

|                           | Training<br>Accuracy | Test<br>Accuracy | Precision | Recall | F-1 score |
|---------------------------|----------------------|------------------|-----------|--------|-----------|
| Multi-layer<br>Perceptron | 0.793                | 0.789            | 0.763     | 0.836  | 0.798     |
| XGBoost                   | 0.851                | 0.846            | 0.845     | 0.847  | 0.846     |
| LightGBM                  | 0.850                | 0.852            | 0.851     | 0.854  | 0.852     |
| RandomForest              | 0.840                | 0.850            | 0.837     | 0.869  | 0.853     |





180

## 181 *3. Feature importance*

182 The variable with the highest importance in constructing the LightGBM model in predicting 183 sarcopenia was BMI (Figure 4). In addition, variables related to individual physical characteristics 184 such as age, waist circumference, height, and weight were found to be important variables in 185 constructing a sarcopenia prediction model. EQ-5D, which represents health-related quality of life, 186 was found to be an important variable in model construction next to variables representing individual 187 physical characteristics, and variables such as the number of hours of work participation per week 188 were the next most important variables, followed by the number of days of participation in walking or 189 moderate- and high-intensity exercise. Gender appeared to be the least important variable in 190 constructing the sarcopenia prediction model.

191



Figure 4. Feature importance of LightGBM model

193

## 195 IV. DISCUSSION

The purpose of this study was to create a model to predict the diagnosis of sarcopenia in the elderly with relatively easy-to-obtain physical characteristics and activity-related variables compared to variables measured by experts with medical equipment such as DXA to measure lean muscle mass. The sarcopenia prediction model constructed in this study showed an accuracy performance of about 85% or more, except for the MLP model. In constructing a high-performance model, variables related to individual physical characteristics such as BMI and age were the most important, followed by variables related to exercise participation or occupational activity.

203 The machine learning model built with the data measured by the diagnostic imaging equipment 204 showed a relatively high level of accuracy. Burns al. (2020) built a machine learning model to predict 205 sarcopenia using abdominal cross-sectional CT taken at the lumbar level of an elderly person, and the 206 accuracy (expressed as AUC score) of this model was around 94%.(19) Ryu et al. (2022) built a 207 model to predict the prevalence of sarcopenia using chest X-ray data in a simpler way than CT scan, 208 and the accuracy (expressed as AUC score) of this model was 74-88%.(20) In addition, even when the 209 machine learning model was constructed with variables other than those measured by imaging 210 equipment, it showed high performance in diagnosing sarcopenia. Ko et al. (2021) developed a model 211 with 81-88% accuracy by analyzing patterns during timed-up-and-go (TUG) and 6-minute walk test 212 (6mWT) by machine learning using an IMU sensor.(25) Kang et al. (2019) created a model to predict 213 sarcopenia with 78-82% accuracy (expressed as AUC score) using physical variables such as BMI and 214 age, blood-related variables such as red blood cell count and white blood cell count, and dietary 215 variables such as fiber intake, fat intake, and protein intake.(21) Similarly, the AUC score of the 216 sarcopenia prediction model using body characteristics and physical activity variables in this study 217 was 87-93%. In addition, the accuracy reached about 85% for three of the four models. The 218 performance of the sarcopenia prediction model generated through this study did not show 219 outstanding performance compared to other models. Nevertheless, there are several advantages 220 compared to sarcopenia predictive models in other studies. First, the variables used to build the model

221 in this study were factors that did not require imaging techniques using expensive medical equipment 222 such as X-ray, CT, and MRI. Second, the variables in this study did not require invasive testing for 223 human specimens, such as blood-related factors or muscle biopsies. Third, the variables used in model 224 construction were not variables that required high-level skills to measure or that had to be measured 225 by experts. Considering these advantages, even if the examiners or investigators do not have high-226 level skills, they will be able to measure the data used in this model, and even the subjects will be able 227 to measure themselves using scales or questionnaires. In addition, the subject can be remotely It may 228 be possible to input data and remotely receive sarcopenia prediction results for these data.

229 Variables related to physical characteristics showed high importance in constructing the LGBM 230 model, which showed the highest accuracy among the models in this study. BMI showed the highest 231 feature importance, followed by age, waist circumference, height, and weight. In the case of 232 sarcopenia in Asian elderly with diabetes, it was found that sarcopenia decreased significantly as BMI 233 increased.(26) In addition, according to the model for predicting sarcopenia constructed with the 234 RandomForest algorithm by Kim et al (2019), BMI served as the first variable to classify the elderly 235 over 60 as sarcopenia and normal.(27) According to the deep learning model built by Somasundaram 236 et al (2022) for the diagnosis of sarcopenia in adolescents, the skeletal muscle area measured by 237 abdominal CT at the L3 level, one of the markers of sarcopenia diagnosis, showed a high correlation 238 (0.75-0.94) with weight, height, age, and BMI in adolescence. (28) Based on these studies, the model 239 constructed in our study would also have shown high importance in constructing the model for 240 physical characteristics variables. In addition, studies related to sarcopenia and quality of life have 241 been conducted. According to the review article, 4 studies measuring quality of life with the Short-242 form General Health Survey (SF-36) and 2 studies using the EQ-5D instrument, Subjects with 243 sarcopenia were found to have significantly higher rates of problems related to quality of life.(29) For 244 this reason, in the model of this study, EQ-5D would have shown the highest importance next to 245 physical characteristics. The feature importance of physical activity-related variables in the model 246 created in this study was lower than that of physical characteristics or quality of life variables. 247 Nevertheless, the importance of physical activity in preventing muscle loss has been emphasized.(30)

There is also evidence that physical activity, a modifiable lifestyle behavior, can partially reverse agerelated skeletal muscle dysfunction.(30,31) Therefore, variables related to physical activity can be used as predictive variables in constructing a sarcopenia prediction model even if the feature importance is not high.

252 There are several limitations in this study. First, the sarcopenia predictive model of this study was 253 constructed using data of the elderly over 60 years of age, so the performance of this model cannot be 254 guaranteed for predicting sarcopenia in adolescents or young people. Second, in constructing the 255 predictive model in this study, the model may have been overfitted because upsampling was used to 256 balance the number of sarcopenia and normal data. Third, according to the inherent characteristics of 257 machine learning models, it was possible to infer the correlation of variables related to sarcopenia 258 prediction by calculating the feature importance in model creation, but it was difficult to interpret it as 259 a causal relationship. Future studies will need to build a model by additionally collecting data from 260 various age groups, and additionally discover several variables that are expected to have a high impact 261 on sarcopenia and use them to construct a model. In addition, research to identify the causal 262 relationship between variables representing high feature importance and sarcopenia through a 263 machine learning model should be conducted.

# 265 V. CONCLUSION

266 In this study, a sarcopenia prediction model was created using physical characteristics and activity 267 variables of sarcopenia and normal elderly over 60 years of age. The performance of the model 268 showed an accuracy of about 85%, and physical characteristics variables such as BMI, age, height, 269 and weight were found to be important in constructing this model. In addition, through feature 270 importance, it was found that activity-related variables such as EQ-5D, occupational activity, and 271 number of days participating in physical activity were also used to construct the model. Since this 272 model was constructed using relatively easy-to-measure variables without using medical equipment, it 273 will be able to help predict and prevent sarcopenia in the elderly in areas with insufficient medical 274 resources or poor access to medical facilities.

# 276 VI. References

| 277 | 1. | Mitchell WK, Williams J, Atherton P, Larvin M, Lund J, Narici M. Sarcopenia, dynapenia,         |
|-----|----|---|
| 278 |    | and the impact of advancing age on human skeletal muscle size and strength; a quantitative      |
| 279 |    | review. Front Physiol. 2012;3:260.  |
| 280 | 2. | Oh JM, Choi J-H, Lee YJ, Lee YR, Youn NH, Song HJ. Association between sarcopenia and           |
| 281 |    | health-related quality of life in Korean adults: based on the Fifth Korean National Health and  |
| 282 |    | Nutrition Examination Survey (2010-2011). Korean J Fam Pract. 2017;7(6):870-6.                  |
| 283 | 3. | Woo N, Kim SH. Sarcopenia influences fall-related injuries in community-dwelling older          |
| 284 |    | adults. Geriatr Nurs (Minneap). 2014;35(4):279-82.  |
| 285 | 4. | Perez Sousa MA, Venegas Sanabria LC, Chavarro Carvajal DA, Cano Gutierrez CA,                   |
| 286 |    | Izquierdo M, Correa Bautista JE, et al. Gait speed as a mediator of the effect of sarcopenia on |
| 287 |    | dependency in activities of daily living. J Cachexia Sarcopenia Muscle. 2019;10(5):1009–15.     |
| 288 | 5. | Biolo G, Cederholm T, Muscaritoli M. Muscle contractile and metabolic dysfunction is a          |
| 289 |    | common feature of sarcopenia of aging and chronic diseases: from sarcopenic obesity to          |
| 290 |    | cachexia. Clin Nutr. 2014;33(5):737–48.   |
| 291 | 6. | Lim H-S, Park Y-H, Suh K, Yoo MH, Park HK, Kim HJ, et al. Association between                   |
| 292 |    | sarcopenia, sarcopenic obesity, and chronic disease in Korean elderly. J Bone Metab.            |
| 293 |    | 2018;25(3):187.   |
| 294 | 7. | Chang S, Lin P. Systematic literature review and meta analysis of the association of            |
| 295 |    | sarcopenia with mortality. Worldviews Evidence  Based Nurs. 2016;13(2):153–62.                  |
| 296 | 8. | Beaudart C, Rizzoli R, Bruyère O, Reginster J-Y, Biver E. Sarcopenia: burden and challenges     |
| 297 |    | for public health. Arch public Heal. 2014;72(1):1–8.  |
| 298 | 9. | Albano D, Messina C, Vitale J, Sconfienza LM. Imaging of sarcopenia: old evidence and new       |
| 299 |    | insights. Eur Radiol. 2020;30:2199-208.   |

| 300 | 10. | Guglielmi G, Ponti F, Agostini M, Amadori M, Battista G, Bazzocchi A. The role of DXA in |
|-----|-----|--|
| 301 |     | sarcopenia. Aging Clin Exp Res. 2016;28:1047–60.   |

- 302 11. Belarmino G, Gonzalez MC, Sala P, Torrinhas RS, Andraus W, D'Albuquerque LAC, et al.
- 303 Diagnosing sarcopenia in male patients with cirrhosis by dual  $\Box$  energy X  $\Box$  ray absorptiometry
- 304 estimates of appendicular skeletal muscle mass. J Parenter Enter Nutr. 2018;42(1):24–36.
- 305 12. Gonzalez MC, Barbosa-Silva TG, Heymsfield SB. Bioelectrical impedance analysis in the
  306 assessment of sarcopenia. Curr Opin Clin Nutr Metab Care. 2018;21(5):366–74.
- 307 13. Yuki A, Ando F, Shimokata H. Transdisciplinary approach for sarcopenia. Sarcopenia:
- definition and the criteria for Asian elderly people. Clin Calcium. 2014;24(10):1441–8.
- 309 14. Reijnierse EM, De Van Der Schueren MAE, Trappenburg MC, Doves M, Meskers CGM,
- 310 Maier AB. Lack of knowledge and availability of diagnostic equipment could hinder the
- diagnosis of sarcopenia and its management. PLoS One. 2017;12(10):e0185837.
- 312 15. Ha Y-C, Hwang S-C, Song S-Y, Lee C, Park K-S, Yoo J-I. Hand grip strength measurement in
- different epidemiologic studies using various methods for diagnosis of sarcopenia: a
  systematic review. Eur Geriatr Med. 2018;9:277–88.
- 315 16. Cawthon PM, Travison TG, Manini TM, Patel S, Pencina KM, Fielding RA, et al. Establishing
- the link between lean mass and grip strength cut points with mobility disability and other
- 317 health outcomes: proceedings of the sarcopenia definition and outcomes consortium
- 318 conference. Journals Gerontol Ser A. 2020;75(7):1317–23.
- Jordan MI, Mitchell TM. Machine learning: Trends, perspectives, and prospects. Science.
  2015;349(6245):255–60.
- 321 18. Kim YJ. Machine learning models for sarcopenia identification based on radiomic features of
  322 muscles in computed tomography. Int J Environ Res Public Health. 2021;18(16):8710.
- 323 19. Burns JE, Yao J, Chalhoub D, Chen JJ, Summers RM. A machine learning algorithm to

|  | 324 | estimate sarco | penia on | abdominal | CT. Acad | Radiol. | 2020;2 | 7(3 | ):311 | -20. |
|--|-----|----------------|----------|-----------|----------|---------|--------|-----|-------|------|
|--|-----|----------------|----------|-----------|----------|---------|--------|-----|-------|------|

- Ryu J, Eom S, Kim HC, Kim CO, Rhee Y, You SC, et al. Chest X aray based opportunistic
  screening of sarcopenia using deep learning. J Cachexia Sarcopenia Muscle. 2023;14(1):418–
- 327 28.
- 328 21. Kang Y-J, Yoo J-I, Ha Y. Sarcopenia feature selection and risk prediction using machine
  329 learning: A cross-sectional study. Medicine (Baltimore). 2019;98(43).
- 330 22. Wu L-W, OuYoung T, Chiu Y-C, Hsieh H-F, Hsiu H. Discrimination between possible
- 331 sarcopenia and metabolic syndrome using the arterial pulse spectrum and machine-learning
  332 analysis. Sci Rep. 2022;12(1):21452.
- 333 23. Beaudart C, Dawson A, Shaw SC, Harvey NC, Kanis JA, Binkley N, et al. Nutrition and
- physical activity in the prevention and treatment of sarcopenia: systematic review. Osteoporos
  Int. 2017;28:1817–33.
- 336 24. Kim KM, Jang HC, Lim S. Differences among skeletal muscle mass indices derived from
  337 height-, weight-, and body mass index-adjusted models in assessing sarcopenia. Korean J
  338 Intern Med. 2016;31(4):643.
- 339 25. Ko JB, Kim KB, Shin YS, Han H, Han SK, Jung DY, et al. Predicting Sarcopenia of Female
  340 Elderly from Physical Activity Performance Measurement Using Machine Learning Classifiers.
  341 Clin Interv Aging. 2021;1723–33.
- Fukuoka Y, Narita T, Fujita H, Morii T, Sato T, Sassa MH, et al. Importance of physical
  evaluation using skeletal muscle mass index and body fat percentage to prevent sarcopenia in
  elderly Japanese diabetes patients. J Diabetes Investig. 2019;10(2):322–30.
- 345 27. Kim S, Yi C, Lim J. Risk Factors for Sarcopenia, Sarcopenic Obesity, and Sarcopenia Without
  346 Obesity in Older Adults. Phys Ther Korea. 2021;28(3):177–85.
- 347 28. Somasundaram E, Castiglione JA, Brady SL, Trout AT. Defining normal ranges of skeletal

| 348 |     | muscle area and skeletal muscle index in children on CT using an automated deep learning     |
|-----|-----|--|
| 349 |     | pipeline: implications for sarcopenia diagnosis. Am J Roentgenol. 2022;219(2):326-36.        |
| 350 | 29. | Tsekoura M, Kastrinis A, Katsoulaki M, Billis E, Gliatis J. Sarcopenia and its impact on     |
| 351 |     | quality of life. GeNeDis 2016 Genet Neurodegener. 2017;213-8.                                |
| 352 | 30. | Pillard F, Laoudj-Chenivesse D, Carnac G, Mercier J, Rami J, Rivière D, et al. Physical      |
| 353 |     | activity and sarcopenia. Clin Geriatr Med. 2011;27(3):449-70.                                |
| 354 | 31. | Bosaeus I, Rothenberg E. Nutrition and physical activity for the prevention and treatment of |
| 355 |     | age-related sarcopenia. Proc Nutr Soc. 2016;75(2):174-80.                                    |