



E

Crosses	+/+	+/-	-/-	L	p-value	%P
<i>Pycr1</i> ^{+/-} x <i>Pycr1</i> ^{+/-}	21	50	35	12	0.092 n.s.	34
<i>Pycr1</i> ^{+/-} x <i>Pycr1</i> ^{-/-}	0	43	53	12	0.146 n.s.	54.5



Suppl Figure 1

Supplementary Figure S1: Generation of *Pycr1* knockout mice. (A) Strategy and allele description for *Pycr1* knockout mice. (B) Schematic of gDNA from WT *Pycr1*. Arrows are depicting the primers used to genotype *Pycr1*^{+/+} (619 bp, F + R1) and *Pycr1*^{-/-} (354 bp, F + R2). (C) Genotyping of *Pycr1*^{+/+}, *Pycr1*^{+/-} and *Pycr1*^{-/-} animals by PCR. (D) Quantitative real time PCR demonstrating that *Pycr1*^{-/-} dermal fibroblasts have virtually no *Pycr1* transcripts left compared to WT. Two-tailed Student's t test, ****p<0.0001 (n = 4). (E) Distribution of genotypes from heterozygous intercrosses and *Pycr1*^{+/-} x *Pycr1*^{-/-} crosses illustrate that *Pycr1*^{-/-} mice are born at expected Mendelian ratios. Two-tailed Student's t test, ns p>0.05. (F) Herovici staining shows that *Pycr1*^{-/-} have less collagen I fibers (pink), whereas collagen III fibers (blue) appear unchanged. Scale bar, 100 μm. (G) Quantification shows no significant difference in collagen I and collagen III fibers between WT and *Pycr1*^{-/-} mice. Intensity was calculated by averaging measurements from three mice/genotypes. Error bars indicate mean ± SEM. Two-tailed Student's t test, ns p>0.05, **p<0.01, ***p<0.001.