

**Supplement - Healthcare provider (HCP) gender distribution of types of HCP initially contacted by individuals with LBP**

Type of HCP		% Female	Source
Non-Rx	Doctor of Chiropractic (DC)	32%	2020 - National Board of Chiropractic Examiners
	Physical Therapist (PT)	65%	2020 - American Physical Therapy Association
	Licensed Acupuncturist (LAc)	65%	2023 - Zippia
Primary Care	Primary Care Physician (PCP)	40%	2020 - American Medical Association
	Nurse Practitioner (Nurse)	87%	2023 - Zippia
	Physician Assistant (PA)	67%	2023 - Zippia
	Doctor of Osteopathy (DO)	43%	2022 - American Osteopathic Association
Physician Specialist	Orthopedic Surgeon (OS)	7%	2020 - American Medical Association
	Physical Medicine and Rehabilitation (PMR)	36%	2020 - American Medical Association
	Pain Management (PM)	19%	2020 - American Medical Association
	Neurologist (Neuro)	31%	2020 - American Medical Association
	Rheumatologist (Rheum)	46%	2020 - American Medical Association
	Neurosurgeon (NS)	9%	2020 - American Medical Association
	MD Other (Oth)	-	
Emergency Medicine/ Urgent Care	Hospital (Hosp)	-	
	Radiologist (Rad)	26%	2020 - American Medical Association
	Emergency Medicine (EM)	28%	2020 - American Medical Association
	Urgent Care (UC)	-	