	Supplement - Table 3 - Type of healthcare service provided by patient home address 5 digit % of Episodes Including Service										5 digit zi	rip code population measures of race/ethnicity and socioeconomic status Risk Ratio and 95% Confidence Interval Comparing Female to Male Baseline						
		Comr	nercial	% of Ep Insurance (CI)	olsodes li	nciuainį	<u> </u>	e care Adv	/antage	e (MA)			nmercial Insuranc			dicare Advantage		
			COVID	-	COVID Late	COVID	Pre-C	COVID	Early C	COVID		COVID	Pre-COVID	Early COVID	Late COVID	Pre-COVID	Early COVID	Late COVID
		F	Μ	F % of P	M F	M is non-F	F Hispanic	M white (F NHW 09	M (-50%)	F and Are	M Pa Depri	vation Index (ADI	51-100) - Non-Wł	nite. Low Income			
	At least one service	3453	3208	-			14592	-			17272	9870						
	Manipulation - Chiropractic								3.9%	5.6%					,	0.68 (0.61, 0.76)		
First	Active Care Manual Therapy	18.2% 10.0%				_	9.9% 4.1%		8.0% 3.4%	7.9%	10.0% 4.0%		0.97 (0.87, 1.07)			1.03 (0.94, 1.11) 0.97 (0.85, 1.10)		
Line	Passive Therapy	10.3%					1.8%		1.3%	1.6%				0.73 (0.65, 0.82)				0.87 (0.72, 1.05)
	Manipulation - Osteopathic	0.6%	-			_	0.2%		0.2%	0.2%		0.2%		0.85 (0.39, 1.86)	1.13 (0.56, 2.28)		· · · · ·	1.07 (0.57, 2.00)
	Acupuncture	0.2%					0.1%	0.1%	0.1%	0.1%		0.1%					1.07 (0.45, 2.53)	
	Imaging - Radiography Rx NSAID	38.7% 29.2%		37.7%		_		35.0% 25.3%					0.95 (0.89, 1.00)		1.00 (0.95, 1.06)	1.04 (1.01, 1.08) 1.02 (0.98, 1.07)	1.00 (0.97, 1.04) 1.05 (1.00, 1.09)	
Second	Rx Muscle Relaxant	25.0%				_							0.84 (0.78, 0.91)		· · · · · ·		0.89 (0.84, 0.94)	
Line	Rx Oral Steroid	10.7%					7.3%		6.2%	7.9%				. , ,			,	0.80 (0.73, 0.88)
	Imaging - MRI Rx Gabapentin	8.9% 1.9%	_			_	11.9% 2.3%	14.1% 2.5%	11.6% 2.4%	13.5% 2.1%			0.84 (0.72, 0.97) 0.90 (0.64, 1.26)		0.99 (0.87, 1.13)		0.86 (0.81, 0.92) 1.14 (0.96, 1.36)	0.90 (0.85, 0.95) 0.96 (0.80, 1.15)
	Rx Other	6.4%	_			_										1.13 (1.05, 1.21)		
	Rx Opioid	11.1%		10.6%			18.6%						0.89 (0.78, 1.01)		1.02 (0.89, 1.16)		0.93 (0.88, 0.99)	
Third	Spinal Injection	4.6%				_	6.9% 7.0%		6.5%	6.7%			0.99 (0.80, 1.23)				0.98 (0.89, 1.08)	
Line	Imaging - CT Surgery	3.4% 0.4%			4.3% 4.2% 0.5% 0.5%	_			6.7% 0.3%	7.0%			0.82 (0.64, 1.04) 0.40 (0.21, 0.77)				0.95 (0.86, 1.04) 0.51 (0.35, 0.75)	0.87 (0.80, 0.94)
NHW 0-50 and ADI 0-50 - Non-White, High Income																		
	At least one service	6344				_	10708		10606		11677				0.01 (0.70, 0.07)			0.00.00.00.00.00
	Manipulation - Chiropractic Active Care	16.6% 24.9%		16.9%					3.8% 10.2%	5.8% 11.4%	4.2% 14.0%		0.81 (0.75, 0.87)	, , ,	0.81 (0.76, 0.87)	0.60 (0.53, 0.68)	0.64 (0.56, 0.74) 0.89 (0.82, 0.97)	
First	Manual Therapy	16.9%		14.0%			8.5%		5.7%	6.3%			1.05 (0.97, 1.14)	· · · · · · · · · · · · · · · · · · ·	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			0.98 (0.88, 1.08)
Line	Passive Therapy	11.7%				_	2.9%		1.9%	2.4%						0.73 (0.62, 0.85)		
	Manipulation - Osteopathic Acupuncture	0.6% 0.7%					0.3%			0.2% 0.8%	0.2%					1.13 (0.61, 2.06) 1.02 (0.78, 1.35)		
	Imaging - Radiography	37.4%		36.2%		_								, , ,	· · · · ·	1.07 (1.02, 1.12)		
	Rx NSAID	29.2%		30.3%									0.97 (0.91, 1.02)	0.99 (0.94, 1.04)	1.00 (0.95, 1.06)	0.96 (0.92, 1.01)	1.06 (1.01, 1.12)	
Second	Rx Muscle Relaxant	22.7%		22.5%									0.90 (0.85, 0.96)			0.90 (0.83, 0.97)		0.89 (0.83, 0.96)
Line	Rx Oral Steroid Imaging - MRI	8.6% 10.0%		-		6 8.8% 6 13.4%	7.1% 13.6%		5.3% 13.3%	7.1%			0.71 (0.64, 0.79) 0.86 (0.78, 0.96)	, , ,	0.78 (0.70, 0.88)	0.85 (0.76, 0.95)	0.74 (0.66, 0.83)	0.75 (0.66, 0.85) 0.89 (0.83, 0.95)
	Rx Gabapentin	1.9%				_			2.5%				0.75 (0.59, 0.96)		0.83 (0.67, 1.04)			0.94 (0.78, 1.13)
	Rx Other	6.5%	-													1.11 (1.03, 1.20)		
Third	Rx Opioid Spinal Injection	8.9% 4.8%				_			15.3% 6.2%	16.4% 6.6%			0.92 (0.83, 1.04) 0.83 (0.71, 0.97)	0.88 (0.79, 0.99)		0.93 (0.86, 0.99)		0.89 (0.83, 0.96) 0.91 (0.82, 1.01)
Line	Imaging - CT	2.3%				_			5.4%	5.8%								1.06 (0.95, 1.17)
	Surgery	0.3%	0.7%	0.3%	0.7% 0.4%	6 0.5%	0.3%		0.3%	0.7%				0.41 (0.24, 0.69)	0.86 (0.51, 1.42)	0.49 (0.32, 0.76)	0.34 (0.21, 0.54)	0.51 (0.33, 0.78)
	At least one service	8280	8202	10278	10508 9850	9564	31659				-	0 - White 26521	e, Low Income					
	Manipulation - Chiropractic					_								0.90 (0.87, 0.94)	0.88 (0.85, 0.92)	0.71 (0.68, 0.74)	0.69 (0.66, 0.72)	0.73 (0.70, 0.76)
	Active Care	16.2%	16.0%		5 15.0% 16.2%				9.0%			10.3%				, , , , , , , , , , , , , , , , , , ,		1.07 (1.03, 1.12)
First Line	Manual Therapy	9.8%				_	4.7% 1.8%		4.0% 1.3%	3.8% 1.6%					1.09 (1.00, 1.20)	, , ,		1.12 (1.04, 1.20)
Line	Passive Therapy Manipulation - Osteopathic	13.7% 0.6%				_			0.2%	0.2%			0.85 (0.79, 0.92)	0.81 (0.52, 1.27)	0.84 (0.78, 0.90)			1.04 (0.91, 1.18) 1.15 (0.83, 1.58)
	Acupuncture	0.2%					0.0%		0.1%	0.1%			1.49 (0.67, 3.31)	2.17 (0.94, 5.03)				1.60 (0.91, 2.82)
	Imaging - Radiography	34.7%		33.9%										1		1.08 (1.05, 1.10)		
	Rx NSAID Rx Muscle Relaxant	20.4% 19.7%		20.2%		6 20.3% 6 21.9%										1.10 (1.06, 1.14) 0.91 (0.87, 0.95)		0.92 (0.88, 0.95)
Second Line	Rx Oral Steroid	11.1%				6 11.1%		10.2%	7.4%	8.9%							0.84 (0.79, 0.88)	0.87 (0.82, 0.92)
Line	Imaging - MRI	8.3%														0.91 (0.87, 0.95)		
	Rx Gabapentin Rx Other	1.2% 6.5%					2.0%						0.81 (0.62, 1.06) 1.19 (1.06, 1.35)	, , ,	0.57 (0.44, 0.75)		0.98 (0.87, 1.11) 1.07 (1.02, 1.11)	0.80 (0.71, 0.91)
	Rx Opioid	10.0%											0.91 (0.83, 1.00)					0.95 (0.92, 0.99)
Third	Spinal Injection	4.3%				_			7.9%	8.1%				0.95 (0.83, 1.08)	0.81 (0.71, 0.92)			0.99 (0.95, 1.04)
Line	Imaging - CT	2.7%		-		_			6.7%	6.3%						0.98 (0.92, 1.05) 0.50 (0.40, 0.62)		1.01 (0.95, 1.06)
	Surgery	0.770	0.3%	0.4/0	, 1.070 0.27	0.7/0	0.576						High Income	0.00)	0.52 (0.20, 0.52)	0.00 (0.40, 0.02)	0.50, 0.50)	5.55 (0.41, 0.01)
	At least one service				17405 18252									1				
	Manipulation - Chiropractic Active Care				26.9% 23.7%											0.75 (0.72, 0.79) 1.01 (0.97, 1.05)		
First	Manual Therapy				5 24.0% 24.37 5 14.8% 15.4%				7.1%							1.04 (0.98, 1.10)		
Line	Passive Therapy	13.4%	14.7%	11.5%	5 12.7% 10.6%	6 12.1%	2.6%	2.6%	2.0%	2.1%	2.0%	2.1%	0.91 (0.86, 0.96)	0.91 (0.86, 0.96)	0.88 (0.83, 0.93)	1.00 (0.90, 1.12)	0.97 (0.86, 1.10)	0.94 (0.84, 1.05)
	Manipulation - Osteopathic	0.8%							0.4%							1.39 (1.04, 1.85)		
	Acupuncture Imaging - Radiography	0.9% 38.4%								0.2%						1.39 (0.95, 2.03)1.07 (1.04, 1.10)		
	Rx NSAID															1.07 (1.04, 1.10)		
Second	Rx Muscle Relaxant	16.5%	20.2%	16.8%	5 20.2% 15.7%	6 18.0%	10.9%	11.8%	11.0%	12.2%	10.9%	12.6%	0.81 (0.78, 0.85)	0.83 (0.80, 0.87)	0.87 (0.83, 0.91)	0.92 (0.88, 0.97)	0.90 (0.86, 0.95)	0.87 (0.83, 0.91)
Line	Rx Oral Steroid	9.0% 10.9%				6 10.0%		10.1%			5.9%					0.80 (0.76, 0.85) 0.84 (0.80, 0.88)		
	Imaging - MRI Rx Gabapentin	10.9%					2.4%									0.94 (0.80, 0.88)		
	Rx Other	5.9%	5.9%	6.9%	6.4% 5.9%	6.1%	11.5%	10.9%	14.4%	13.4%	11.4%	11.2%	1.01 (0.93, 1.10)	1.07 (0.99, 1.16)	0.97 (0.89, 1.05)	1.06 (1.00, 1.11)	1.07 (1.03, 1.12)	1.02 (0.98, 1.07)
Thind	Rx Opioid	7.7%														1.01 (0.96, 1.05)		
Third Line	Spinal Injection Imaging - CT	5.4% 2.0%			+ +	_	8.7% 5.5%			9.9% 5.6%						0.89 (0.84, 0.95) 1.09 (1.01, 1.17)		
	Surgery	0.4%			1 1	-	0.5%									0.46 (0.37, 0.57)		

Pre=Pre-COVID period, Early=first 12 months post-COVID, Late=13-24 months post-COVID

For risk ratio, cells in red indicate the confidence interval includes 1 and likelihood of HCP being initially contacted is not different than the male reference for the same time period