

The Impact of Post-Acute Care on Depression and Anxiety in Stroke Patients: A Prospective Study to Explore the Mediating Effect of Cognitive Function

Shi-Jer Lou, PhD^{1,2*}, Hsiu-Fen Lin, MD^{3,4*}, Yu-Tsz Shiu, MS², Hon-Yi Shi, DrPH, MS^{1,2,5,6,7}

¹Graduate Institute of Technological and Vocational Education, National Pingtung University of Science and Technology, Pingtung 91201, Taiwan.

²Department of Healthcare Administration and Medical Informatics, Kaohsiung Medical University, Kaohsiung 80708, Taiwan.

³Department of Neurology, Kaohsiung Medical University Hospital, Kaohsiung 80756, Taiwan.

⁴Department of Neurology, Kaohsiung Medical University, Kaohsiung 80708, Taiwan.

⁵Department of Business Management, National Sun Yat-sen University, Kaohsiung 80424, Taiwan.

⁶Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung 80756, Taiwan.

⁷Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan.

Running title: Impact of PAC on depression and anxiety

*Shi-Jer Lou and Hsiu-Fen Lin contributed equally to this work and share first authorship.

Correspondence to: Hon-Yi Shi, DrPH, Department of Healthcare Administration and Medical Informatics, Kaohsiung Medical University, No. 100, Shih-Chuan 1st Road, 80708 Kaohsiung, Taiwan. E-mail: hshi@kmu.edu.tw.

TEL: +886-7-3211101#2648.

FAX: +886-7-3137487.

WHAT IS KNOWN

- Research suggests that cognitive function, including depression and anxiety, significantly improved for patients using post-acute care (PAC).
- Cognitive function was significantly associated with post-stroke depression and anxiety in patients with stroke.

WHAT THE STUDY ADDS

- PAC had a mediating effect on cognition and depression at week 12 on cognition and anxiety at week 12, and it also had a direct effect on depression and anxiety in the first year.
- Acute lengths of stay had a significant moderation effect in the model, but the model lost its mediating effect when the moderation variable was added.
- Patients with stroke should receive post-acute care as soon as possible to improve their cognitive function after rehabilitation, to maximize the effectiveness of treatment for mental disorders, and to reduce the burden of the disease.

ABSTRACT

BACKGROUND: Cognitive function was significantly associated with post-stroke depression and anxiety in stroke patients. However, no studies have examined whether there is an interaction. This study purposed to investigate whether cognitive function mediates the effect of enrollment in post-acute care (PAC) programs on depression or anxiety in stroke patients and whether the indicators are moderated in the pathway.

METHODS: This is a prospective observational cohort study. A group of patients who had received PAC for stroke at one of two medical centers (PAC group, $n = 2,087$) was compared with a group who had received standard care for stroke at one of four hospitals (three regional hospitals and one district hospital; non-PAC group, $n = 1,591$) in Taiwan from March, 2015, to March, 2022. The effects of PAC on cognitive function and depression and anxiety at baseline, 12th week, and 1st year after rehabilitation were investigated using structural equation modeling (SEM). The effect of each variable on the moderation of different pathways in the model was analyzed using AMOS 23.0, and The SPSS PROCESS macro also was used to perform mediation analysis.

RESULTS: The PAC program had a mediating effect on cognition and depression at week 12 ($a*b = -0.098$, $P < 0.05$) on cognition and anxiety at week 12 ($a*b = -0.107$, $P < 0.05$), and the PAC program had a direct effect on depression and anxiety in the first year. It was found that acute lengths of stay had a significant moderation effect in the model ($X*W \rightarrow M = 0.204$, $P = 0.002$), but the model lost its mediating effect when the moderation variable was added.

CONCLUSIONS: Patients with stroke should receive post-acute care as soon as possible to improve their cognitive function after rehabilitation, to maximize the effectiveness of treatment for mental disorders, and to reduce the burden of the disease.

Key Words: post-acute care, stroke, structural equation model, depression, anxiety, cognitive function

INTRODUCTION

The post-acute care (PAC) program can significantly reduce disability and death in patients with stroke.^{1,2} The National Health Insurance Agency announced the PAC program in 2014, designed to replace long-term hospitalization in medical centers.³ It allows patients to continue to receive appropriate care after treating acute illness. Initially, the PAC program was established for acute stroke patients who are medically stable with mild to moderate to severe functional impairment and who have the potential for active rehabilitation. After PAC was implemented, it was found that the patients receiving post-acute care showed improvement in all indicators, including activities of daily living, mobility, swallowing, and cognitive function.⁴⁻⁶

Research suggests that cognitive function, including depression and anxiety, significantly improved for patients using PAC.^{7,8} Moreover, cognitive function was significantly associated with post-stroke depression and anxiety in stroke patients.^{9,10} However, no studies have examined whether there is an interaction. Since depression and anxiety in patients with stroke vary over time, research must examine the mediating and moderating effects at different time points to gain a more detailed understanding of the onset of depression and anxiety symptoms in stroke patients after PAC rehabilitation. Therefore, the present study investigated whether cognitive functioning affected depression and anxiety in stroke patients in post-acute care and whether demographic,

clinical, and quality-of-care characteristics had a moderating effect on the model.

METHODS

Study Design and Sample

This study prospectively recruited 3,678 stroke patients who received rehabilitation program at two medical centers, three regional hospitals, and one district hospital in southern Taiwan from March 1, 2014 to March 1, 2022. If patients were eligible for the PAC program and agreed to be referred to a district or regional hospital for PAC, they were selected for the PAC group. If the patients continued to receive acute care or were discharged from the medical center, they were selected for the non-PAC group. The patients' physical functional status at the time of admission, at week 12, and at year one was tracked using questionnaires administered by trained research assistants.

These inclusion criteria were as follows: (1) Patients with acute cerebrovascular disease who were medically stable (controllable vital characteristics within 72 hours); (2) Patients with functional status assessed as mild to moderate or severe functional impairment (Modified Rankin Scale, mRS 2-4);¹¹ (3) Patients with active participation in rehabilitation and the potential for positive rehabilitation. The exclusion criteria included patients who were under 18 with cognitive

impairment (Mini-Mental State Examination, MMSE=0)¹² and the inability to communicate. The study protocol was approved by the Institutional Review Board of the Kaohsiung Medical University Hospital (KMUH-IRB-20140308).

Research Instruments

The medical record review included three main attributes: demographic attributes, clinical attributes, and healthcare quality attributes. The mRS assesses the degree of dependence on daily living activities in stroke patients and is divided into seven levels, with 0 representing no symptoms, 5 representing severe disability, and 6 representing death. The MMSE is an 11-question scale measuring time and place orientation, attention and computation, immediate and short-term memory, language, and visuospatial abilities. The maximum score is 30, with higher scores indicating better ability and scores below 25 indicating severe (≤ 9), moderate (10-20), or mild (21-24) impairment, respectively. The Beck Depression Inventory (BDI) comprises 21 questions, each with four options, ranging from 0 (not at all) to 3 (severe).¹³ The total score is 63, with higher scores indicating more severe depression. The Beck Anxiety Inventory (BAI) comprises 21 questions, each with four options, ranging from 0 (not at all) to 3 (severe).¹⁴ Its total score is 63, with higher scores indicating more severe anxiety.

Research Variables

The independent variable was participating in PAC program or not. The dependent variables were the BDI and BAI scores. The mediating variable was the MMSE score. The moderating variables were age, gender, education, body mass index (BMI), smoking, alcohol drinking, acute lengths of stay, readmission in 30 days, stroke recurrence, use of a urinary catheter or nasogastric tube, hypertension, diabetes mellitus, hyperlipidemia, atrial fibrillation, and previous stroke.

Statistical Analysis

The unit of analysis in this study was the individual patient with stroke. This study weighted the independent and moderating variables with the inverse probability of treatment weighting (IPTW) to balance the differences in the relevant variables between the two groups.

We used the structural equation model (SEM) to determine the mediating effects, explain the path coefficients, and investigate the mediating effects at different time points. We also used the cross-delay model with PAC as the independent variable, cognitive function at different time points as the mediating variable, and depression symptom and anxiety symptom at different time points as the dependent variables.

The mediating and moderating effect analysis focused on the critical moderators, and the model with the most mediating effect in SEM was used to explore the moderating variables.

Demographic, clinical, and quality of care characteristics were used to find the most moderating variables, which moderated the path between the independent and mediating variables and between the mediating and dependent variables. Therefore, the mediating and moderating effect model was built accordingly.

We performed statistical analysis using AMOS 23.0 and SPSS 21.0 software. The PROCESS module provided by Hayes (2013) for SPSS syntax was installed in SPSS. The models (Model 7) (Model 14) and (Model 58) were also used to elaborate the notion of mediation and interference and to analyze the mixed models.

RESULTS

Before and After IPTW

2,087 patients were in the PAC group with a mean age of 68.41 years (standard deviation, SD 13.02 years), and the majority were male patients (1,345, 64.4%); 1,591 patients were in the non-PAC group with a mean age of 70.12 years (SD 13.89 years), and the majority were male patients (1,011, 63.5%). They were weighted by IPTW and the distribution of the variables in the

two groups was not significantly different (Table 1).

Structural Equation Modeling of Overall Depression or Anxiety

The participation of PAC positively predicted cognitive function at week 12 ($b=1.42$, $P<0.001$)

(Figure 1). This subsequently influenced patients' depression scores at week 12 and at year 1.

Particularly, pre-rehabilitation (baseline) cognitive function positively predicted depression scores

at week 12 ($b=0.05$, $P<0.001$). However, whereas the path of PAC on cognitive function at year 1

was not significant but had a direct effect on depression at year 1 ($b=0.88$, $P<0.001$). Cognitive

function at week 12 fully mediated between PAC participation and depression at week 12

($b=-0.07$, $P<0.001$). At year 1, the participation of PAC had a direct effect on depression.

Furthermore, at week 12, cognitive function fully mediated between PAC participation and

anxiety at week 12 ($b= -0.07$, $P<0.001$) (Figure 2). However, whereas at year 1, PAC participation

had a direct impact on anxiety ($b=1.01$, $P<0.001$). Overall, besides the mediating effect of the

depression-anxiety model, there were positive prediction effects among cognitive function,

depression, and anxiety at different assessment time points, with all paths being significant

($P<0.05$). In Figure 1, the pre-rehabilitation cognitive function affected cognitive function at week

12 ($b=0.76$, $P<0.001$), and the cognitive function at week 12 affected the cognitive function at

year 1 ($b=0.85$, $P<0.001$). Regarding depression, pre-rehabilitation depression affected depression

at week 12 ($b=0.18$, $P<0.001$), and depression at week 12 also affected depression at year 1 ($b=0.19$, $P<0.001$). In Figure 2, regarding anxiety, pre-rehabilitation anxiety affected anxiety at week 12 ($b=0.15$, $P<0.001$). Anxiety at week 12 also affected anxiety at year 1 ($b=0.20$, $P<0.001$). These findings suggest that the patient's cognitive function, depression, and anxiety were affected over time.

Moderating Effect

First, in the path from the independent variables to the mediating variables of the 12th week depression model (education ($b=0.582$) and acute lengths of stay ($b=0.203$)) and anxiety model (education ($b=0.517$) and acute lengths of stay ($b=0.131$)), had a moderating effect ($P<0.05$) (Table 2). Furthermore, in the path from the mediating variables to the dependent variables of the 12th week depression model (age ($b=-0.002$), alcohol drinking ($b=-0.145$), acute lengths of stay ($b=-0.003$), foley catheter ($b=-0.112$), nasogastric tube ($b=-0.629$), and diabetes mellitus ($b=-0.647$)) and anxiety model (body mass index ($b=0.006$), acute lengths of stay ($b=-0.004$), and previous stroke ($b=-0.092$)) had a significant moderating effect ($P<0.05$) (Table 3). Additionally, the results in Tables 2 and 3 indicate that acute lengths of stay as a moderating variable was significant in both paths of the depression and anxiety models. Therefore, acute lengths of stay was added as a moderating variable in the mediating and moderating effects analysis.

Analysis of Mediating and Moderating Effects

Regarding the depression and anxiety mediating and moderating models, PAC significantly predicted cognitive function at week 12 ($a_1=-2.617$, $P=0.043$) (Table 4). Acute lengths of stay significantly predicted cognitive function at week 12 ($a_2=-0.214$, $P<0.001$). The interaction between PAC and acute lengths of stay significantly predicted cognitive function at week 12 ($a_3=0.204$, $P=0.002$). In the mediating and moderating models, cognitive function at week 12 did not significantly predict depression and anxiety at week 12. In contrast, acute lengths of stay significantly predicted depression and anxiety at week 12 ($b_2=0.074$, $P=0.015$ and $b_2=0.094$, $P=0.001$, respectively). The interaction between cognitive function at week 12 and acute lengths of stay also significantly predicted depression and anxiety at week 12 ($b_3=-0.003$, $P=0.021$ and $b_3=-0.004$, $P=0.002$, respectively). Additionally, PAC positively influenced the direct effect of depression and anxiety at week 12 but not at a significant level. This indicates that although acute lengths of stay had a significant moderating effect, the original mediating effect of the depression and anxiety models was lost when the moderating effect was added.

DISCUSSION

Although PAC has been promoted for the past decades, evidence of its mediating effect of

cognitive function has been limited.^{15,16} To address this issue, we used IPTW matching in this study. It has been demonstrated that there was a significant improvement in cognitive function at week 12 in the PAC group compared to the non-PAC group ($P < 0.05$), but there was no significant difference at year 1.¹⁷ It also has noted significant improvements in cognitive function scores after a mean of 43.57 days of PAC participation,¹⁸ which is consistent with the path significance in the current study. There was no correlation between PAC participation and cognitive function in the long term, but there was a more significant correlation in the short term. One study was designed to examine how long that cognitive improvement lasts and to compare depressed patients' cognitive status with that of non-depressed patients with comparable lesions.¹⁹ They concluded that stroke patients who receive treatment consistently in the early stages of depression can sustain cognitive improvement for more than two years. Therefore, providing intensive rehabilitation therapy early in the progression course can help improve patients' depression symptoms.

The present study is consistent with previous literatures in that anxiety symptoms after stroke were associated with cognitive function, physical function, and quality of life.²⁰⁻²² Moreover, anxiety symptoms after stroke were significantly associated with lower cognitive function before rehabilitation. However, no significant correlation was observed between the vertical association of anxiety symptoms after stroke with changes in cognitive function over one year.^{23,24}

Education had a moderating effect on the PAC's prediction of cognitive function at week 12. Age and alcohol drinking had a significant moderating effect on cognitive function at week 12 for predicting depression at week 12. However, it had no moderating effect on cognitive function at week 12 for predicting anxiety at week 12. It has suggested that education is associated with cognitive function and that those with higher education had significantly better cognitive performance than those with lower education.^{25,26} In the present study, at three months after stroke, 37% of the patients were at risk for depressive symptoms. Further, depression scores at three months after stroke were higher in patients aged 25-54 years than in those aged 75 years and older, as well as in those aged 55-64.²⁷ Past studies indicated that alcohol drinking was significantly associated with the severity of depressive symptoms and that depressive symptoms were more severe in women than men.^{28,29} PAC was not a significant predictor of cognitive function at week 12. In contrast, a nasogastric tube, a foley catheter, and diabetes mellitus were significant moderators in the path of cognitive function at week 12, predicting depression at week 12. Previous stroke was a significant moderator in the path of cognitive function at week 12, predicting anxiety at week 12. Therefore, depressive symptoms will increase the risk of developing diabetes mellitus by 1.62 to 2.52 times, suggesting that the product (interaction) of diabetes mellitus and cognitive function also predicts depression.^{30,31} Another study found a

significant association between severe anxiety symptoms and an increased risk of stroke.³²

However, age, education, alcohol drinking, a nasogastric tube, a foley catheter, diabetes mellitus, and previous stroke had only moderating effects in a single path, not in both paths. Thus, demographic attributes and clinical attributes were not included in the final mediating and moderating effects analysis of the present study.

Acute lengths of stay was significant in each model path for depression and anxiety symptoms but was not significant for stroke recurrence and readmission in 30 days. First, the average lengths of stay for the patients with stroke in Taiwan was longer than it in foreign countries.^{3,33} It also indicated that acute lengths of stay was negatively associated with depression and anxiety symptoms after one month's hospital treatment.³⁴ In contrast, readmission in 30 days was not associated with depression and anxiety symptoms. The mediating model lost its mediating effect when acute lengths of stay was added as a moderating variable; the week 12 cognitive function predicting week 12 depression and the path of depression was not significant. This could be explained as the mediating effect of acute lengths of stay in the PAC prediction of the week 12 cognitive function path. Acute lengths of stay was significantly correlated to post-rehabilitation cognitive function and was an independent predictor, and could have a more significant effect as a mediating variable in the model than as a moderating variable.

There are three limitations to the current study. First, we only included stroke patients from six hospitals in Taiwan, which is not representative of patients with stroke across Taiwan or other regions and is regionally limited. Furthermore, the follow-up period was one year, which lacks long-term follow-up of patients and cannot investigate the long-term effects of PAC or changes in cognitive function and psychological status. Finally, the patients dictated the questionnaire, and interviews were conducted in multiple stages, which may be influenced by patients' subjective judgment, recall bias, or social expectations.

PAC improved the anxiety and depression symptoms of patients with stroke after 12 weeks via the effects of cognitive function and directly affected anxiety and depression symptoms after one year of rehabilitation. These findings suggest that participation in PAC program can improve anxiety and depression symptoms in stroke patients. The cognitive function, anxiety, and depression symptoms of patients with stroke who participated in PAC program changed over time after rehabilitation, i.e., the earlier the patients received rehabilitation, the more their anxiety and depression symptoms improved. Thus, cognitive function mediated the relationship between PAC and depression and anxiety symptoms. Other moderating variables influenced it, while the mediating role may be lost with the addition of moderation. Therefore, we recommend that

healthcare providers should give multiple cognitive rehabilitation or psychotherapy to the

long-term inpatients participating in PAC.

Sources of Funding:

This study was supported by funding from the Kaohsiung Medical University

(KMU-DK(A)112009) and the Ministry of Science and Technology (MOST

108-2410-H-037-006-SS3 & 111-2410-H-037-002-MY3) in Tai-wan.

Disclosures:

The authors declare no conflict of interest.

REFERENCES

1. Ackerly DC, Grabowski DC. Post-acute care reform--beyond the ACA. *N Engl J Med*. 2014;370:689-691.
2. Bushnell CD, Duncan PW, Lycan SL, Condon CN, Pastva AM, Lutz BJ, Halladay JR, Cummings DM, Arnan MK, Jones SB, et al. A Person-Centered Approach to Poststroke Care: The COMprehensive Post-Acute Stroke Services Model. *J Am Geriatr Soc*. 2018;66:1025-1030.
3. Wang CY, Chen YR, Hong JP, Chan CC, Chang LC, Shi HY. Rehabilitative post-acute care for stroke patients delivered by per-diem payment system in different hospitalization paths: A Taiwan pilot study. *Int J Qual Health Care*. 2017;29:779-784.
4. Skopec L, Huckfeldt PJ, Wissoker D, Aarons J, Dey J, Oliveira I, Zuckerman S. Home Health And Postacute Care Use In Medicare Advantage And Traditional Medicare. *Health Aff (Millwood)*. 2020;39:837-842.
5. Werner RM, Konetzka RT. Trends in Post-Acute Care Use Among Medicare Beneficiaries: 2000 to 2015. *JAMA*. 2018;319:1616-1617.
6. McArthur KS, Quinn TJ, Higgins P, Langhorne P. Post-acute care and secondary prevention after ischaemic stroke. *BMJ*. 2011;342:d2083.
7. Shier V, Edelen MO, McMullen TL, Ahluwalia SC, Chen EK, Dalton SE, Paddock S,

- Rodriguez A, Sherbourne CD, Mandl S, et al. Standardized assessment of cognitive function of post-acute care patients. *J Am Geriatr Soc.* 2022;70:1001-1011.
8. Middleton A, Graham JE, Lin YL, Goodwin JS, Bettger JP, Deutsch A, Ottenbacher KJ. Motor and Cognitive Functional Status Are Associated with 30-day Unplanned Rehospitalization Following Post-Acute Care in Medicare Fee-for-Service Beneficiaries. *J Gen Intern Med.* 2016;31:1427-1434.
9. Zhang SH, Wang YL, Zhang CX, Zhang CP, Xiao P, Li QF, Liang WR, Pan XH, Zhou MC. Effect of Interactive Dynamic Scalp Acupuncture on Post-Stroke Cognitive Function, Depression, and Anxiety: A Multicenter, Randomized, Controlled Trial. *Chin J Integr Med.* 2022;28:106-115.
10. Lu W, Pai M, Scholes S, Xue B. Do depressive symptoms link chronic diseases to cognition among older adults? Evidence from the Health and Retirement Study in the United States. *J Affect Disord.* 2021;294:357-365.
11. Banks JL, Marotta CA. Outcomes validity and reliability of the modified Rankin scale: Implications for stroke clinical trials: A literature review and synthesis. *Stroke.* 2007;38:1091-1096.
12. Katzman R, Zhang MY, Qu OY, Wang ZY, Liu WT, Yu E, Wong SC, Salmon DP, Grant I. A Chinese version of the Mini-Mental State Examination; impact of illiteracy in a Shanghai

- dementia survey. *J Clin Epidemiol.* 1988;41:971-978.
13. Beck AT, Steer RA. Manual for the Revised Beck Depression Inventory. San Antonio, TX: Psychological Corporation; 1987.
 14. Beck AT, Steer RA. Manual for the Beck Anxiety Inventory. San Antonio, TX: Psychological Corporation; 1990.
 15. Rao BR, Jung EH, Dickert NW. Getting Cost Discussions Right: Nudging Patients to Avoid Cognitive Pitfalls. *Circ Cardiovasc Qual Outcomes.* 2023;16:e009447.
 16. Shier V, Edelen MO, McMullen TL, Ahluwalia SC, Chen EK, Dalton SE, Paddock S, Rodriguez A, Sherbourne CD, Mandl S, et al. Standardized assessment of cognitive function of post-acute care patients. *J Am Geriatr Soc.* 2022;70:1001-1011.
 17. Chen YC, Yeh YJ, Wang CY, Lin HF, Lin CH, Hsien HH, Hung KW, Wang JD, Shi HY. Cost Utility Analysis of Multidisciplinary Postacute Care for Stroke: A Prospective Six-Hospital Cohort Study. *Front Cardiovasc Med.* 2022;9:826898.
 18. Lai CL, Tsai MM, Luo JY, Liao WC, Hsu PS, Chen HY. Post-acute care for stroke - a retrospective cohort study in Taiwan. *Patient Prefer Adherence.* 2017;11:1309-1315.
 19. Narushima K, Chan KL, Kosier JT, Robinson RG. Does cognitive recovery after treatment of poststroke depression last? A 2-year follow-up of cognitive function associated with poststroke depression. *Am J Psychiatry.* 2003;160:1157-1162.

20. Tang WK, Lau CG, Mok V, Ungvari GS, Wong KS. Impact of anxiety on health-related quality of life after stroke: a cross-sectional study. *Arch Phys Med Rehabil.* 2013;94:2535-2541.
21. Pinter D, Fandler-Höfler S, Fruhwirth V, Berger L, Bachmaier G, Horner S, Eppinger S, Kneihsl M, Enzinger C, Gattringer T. Relevance of Cognition and Emotion for Patient-Reported Quality of Life After Stroke in Working Age: An Observational Cohort Study. *Front Neurol.* 2022;13:869550.
22. Kovacs AH, Brouillette J, Ibeziako P, Jackson JL, Kasparian NA, Kim YY, Livecchi T, Sillman C, Kochilas LK; American Heart Association Council on Lifelong Congenital Heart Disease and Heart Health in the Young; and Stroke Council. Psychological Outcomes and Interventions for Individuals With Congenital Heart Disease: A Scientific Statement From the American Heart Association. *Circ Cardiovasc Qual Outcomes.* 2022;15:e000110.
23. Lee EH, Kim JW, Kang HJ, Kim SW, Kim JT, Park MS, Cho KH, Kim JM. Association between Anxiety and Functional Outcomes in Patients with Stroke: A 1-Year Longitudinal Study. *Psychiatry Investig.* 2019;16:919-925.
24. Kleineidam L, Stark M, Riedel-Heller SG, Pabst A, Schmiedek F, Streit F, Rietschel M, Klinger-König J, Grabe HJ, Erhardt A, et al. The assessment of cognitive function in the German National Cohort (NAKO) - Associations of demographics and psychiatric symptoms

- with cognitive test performance. *World J Biol Psychiatry*. 2022:1-15.
25. Cheng HM, Chuang SY, Ko YT, Liao CF, Pan WH, Liu WL, Hung CY, Chen CH. Education level may modify the association between cardiac index and cognitive function among elders with normal ejection function. *Front Cardiovasc Med*. 2022;9:844396.
26. Chen Y, Lv C, Li X, Zhang J, Chen K, Liu Z, Li H, Fan J, Qin T, Luo L, et al. The positive impacts of early-life education on cognition, leisure activity, and brain structure in healthy aging. *Aging (Albany NY)*. 2019;11:4923-4942.
27. McCarthy MJ, Sucharew HJ, Alwell K, Moomaw CJ, Woo D, Flaherty ML, Khatri P, Ferioli S, Adeoye O, Kleindorfer DO, et al. Age, subjective stress, and depression after ischemic stroke. *J Behav Med*. 2016;39:55-64.
28. Bellos S, Skapinakis P, Rai D, Zitko P, Araya R, Lewis G, Lionis C, Mavreas V. Longitudinal association between different levels of alcohol consumption and a new onset of depression and generalized anxiety disorder: results from an international study in primary care. *Psychiatry Res* 2016;243:30-34.
29. Franco-Urbano MS, Rodríguez-Martínez MDC, García-Pérez P. The Impact of Depression on the Functional Outcome of the Elderly Stroke Victim from a Gender Perspective: A Systematic Review. *Healthcare (Basel)*. 2022;10:2110.
30. Mezuk B, Eaton WW, Albrecht S, Golden SH. Depression and type 2 diabetes over the


lifespan: a meta-analysis. *Diabetes Care*. 2008;31:2383-2390.

31. Demakakos P, Pierce MB, Hardy R. Depressive symptoms and risk of type 2 diabetes in a national sample of middle-aged and older adults: the English longitudinal study of aging. *Diabetes Care*. 2010;33:792-797.
32. Lambiase MJ, Kubzansky LD, Thurston RC. Prospective study of anxiety and incident stroke. *Stroke*. 2014;45:438-443.
33. Tung YC, Chang GM, Cheng SH. Long-term effect of fee-for-service-based reimbursement cuts on processes and outcomes of care for stroke: interrupted time-series study from Taiwan. *Circ Cardiovasc Qual Outcomes*. 2015;8:30-37.
34. Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, Melloni EMT, Furlan R, Ciceri F, Rovere-Querini P, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behav Immun*. 2020;89:594-600.

FIGURE LEGENDS

FIGURE 1. The whole model using structural equation model. The solid line represents a significant relationship between the two variables. Numbers are path coefficients. PAC, post-acute care; MMSE, mini-mental state examination; BDI, Beck Depression Inventory; BAI, Beck Anxiety Inventory. *P<0.05, **P<0.01, *P<0.001. To investigate whether cognitive function mediates the effect of enrollment in PAC programs on depression in stroke patients and whether the indicators are moderated in the pathway;**

FIGURE 2. The whole model using structural equation model. The solid line represents a significant relationship between the two variables. Numbers are path coefficients. PAC, post-acute care; MMSE, mini-mental state examination; BDI, Beck Depression Inventory; BAI, Beck Anxiety Inventory. *P<0.05, **P<0.01, *P<0.001. To investigate whether cognitive function mediates the effect of enrollment in PAC programs on anxiety in stroke patients and whether the indicators are moderated in the pathway.**






Table 1. Baseline Characteristics of the Study Population Before and After Inverse Probability of Treatment Weighting (IPTW) (N=3,678)

Characteristics	Before IPTW			After IPTW		
	Non-PAC(n=1,591)	PAC (n=2,087)	P value	Non-PAC	PAC	P value
Age, years	70.12±13.89	68.41±13.02	0.168	69.93±14.28	69.36±12.59	0.741
Gender	Male	1,011(63.5%)	1,345(64.4%)	291	300	0.693
	Female	580(36.5%)	742(35.6%)	178	174	
Education, years	8.54±5	8.02±1.77	0.114	8.11±5.05	7.98±1.79	0.593
Body mass index, kg/m ²	24.29±3.42	24.32±4.01	0.918	24.48±3.61	24.37±3.95	0.673
Acute lengths of stay, days	15.56±9.88	18.71±9.55	<0.001	17.07±11.01	17.22±9.24	0.815
Readmission in 30 days	Yes	176(11.1%)	107(5.1%)	38	34	0.591
	No	1,415(88.9%)	1,980(94.9%)	430	439	
Stroke recurrence	Yes	217(13.6%)	55(2.6%)	35	31	0.573
	No	1,374(86.4%)	2,032(97.4%)	433	443	
Nasogastric tube	Yes	449(28.2%)	456(21.8%)	125	119	0.587
	No	1,142(71.8%)	1,631(78.2%)	354	343	
Foley catheter	Yes	279(17.5%)	154(7.4%)	57	53	0.650
	No	1,312(82.5%)	1,933(92.6%)	412	420	
Hypertension	Yes	1,129(71.0%)	1,559(74.7%)	347	352	0.924
	No	462(29.0%)	528(25.3%)	121	121	
Hyperlipidemia	Yes	456(28.7%)	736(35.3%)	150	152	0.996
	No	1,135(71.3%)	1,351(64.7%)	318	322	
Diabetes mellitus	Yes	585(36.8%)	890(42.6%)	191	203	0.513
	No	1,006(63.2%)	1,197(57.4%)	278	271	
Atrial fibrillation	Yes	140(8.8%)	176(8.4%)	45	42	0.697
	No	1,451(91.2%)	1,911(91.6%)	423	431	
Previous stroke	Yes	379(23.8%)	302(14.5%)	89	88	0.859
	No	1,212(76.2%)	1,785(85.5%)	379	386	
MMSE score	2	84(5.3%)	77(3.7%)	21	21	0.989
	3	254(16.0%)	302(14.5%)	67	66	
	4	1,253(78.7%)	1,708(81.8%)	381	386	

Continuous variables are presented as mean ± standard deviation, and categorized variables are presented as n (%). PAC, post-acute care; MMSE, Mini-Mental State Examination.

Table 2. The Moderation Effect in the Path from the Independent Variables to the Mediating Variables of the 12th Week Depression or Anxiety Model After Post-Acute Care (PAC) Program

Moderator (W)	Depression Model		Anxiety Model	
	b	P value	b	P value
Age, years	-0.092	0.051	-0.081	0.059
Gender, female vs. male	0.752	0.579	0.827	0.599
Body mass index, kg/m ²	0.123	0.488	0.130	0.540
Education, years	0.582	0.024	0.517	0.020
Smoking, yes vs. no	0.628	0.686	0.670	0.699
Drinking wine, yes vs. no	-0.394	0.841	-0.434	0.887
Acute lengths of stay, days	0.203	0.002	0.131	0.001
Readmission in 30 days, yes vs. no	0.904	0.714	0.947	0.841
Stroke recurrence, yes vs. no	-2.269	0.456	-2.137	0.407
Foley catheter, yes vs. no	1.818	0.380	1.751	0.307
Nasogastric tube, yes vs. no	-0.489	0.740	-0.402	0.674
Hypertension, yes vs. no	-0.414	0.775	-0.497	0.798
Diabetes mellitus, yes vs. no	1.206	0.366	1.362	0.398
Hyperlipidemia, yes vs. no	1.442	0.305	1.167	0.275
Atrial fibrillation, yes vs. no	1.501	0.517	1.576	0.501
Previous stroke, yes vs. no	1.419	0.394	1.294	0.300
MMSE scores	0.400	0.739	0.412	0.727

MMSE, Mini-Mental State Examination.

Table 3. The Moderation Effect in the Path from the Mediating Variables to the Dependent Variables of the 12th Week Depression or Anxiety Model After Post-Acute Care (PAC) Program

Moderator (W)	Depression Model		Anxiety Model	
	b	P value	b	P value
Age, years	-0.002	0.028	-0.001	0.366
Gender, female vs. male	0.036	0.224	-0.037	0.162
Body mass index, kg/m ²	-0.001	0.693	0.006	0.050
Education, years	-0.001	0.953	-0.006	0.069
Smoking, yes vs. no	-0.016	0.626	0.018	0.553
Drinking wine, yes vs. no	-0.145	<0.001	-0.042	0.270
Acute lengths of stay, days	-0.003	0.021	-0.004	0.002
Readmission in 30 days, yes vs. no	0.054	0.296	0.042	0.375
Stroke recurrence, yes vs. no	0.046	0.354	-0.006	0.897
Foley catheter, yes vs. no	-0.112	0.009	-0.009	0.812
Nasogastric tube, yes vs. no	-0.629	0.039	-0.018	0.511
Hypertension, yes vs. no	-0.430	0.219	0.054	0.087
Diabetes mellitus, yes vs. no	-0.647	0.019	-0.009	0.714
Hyperlipidemia, yes vs. no	-0.001	0.981	0.005	0.850
Atrial fibrillation, yes vs. no	0.049	0.381	-0.065	0.198
Previous stroke, yes vs. no	-0.011	0.758	-0.092	0.005
MMSE scores	0.005	0.900	-0.012	0.733

MMSE, Mini-Mental State Examination.

Table 4. The Mediation Effect of Depression and Anxiety and the Moderation Effect on All Paths

Estimated variables		12 th week cognitive function (M)				12 th week depression (Y)			12 th week anxiety (Y)		
		b	t value	P value		b	t value	P value	b	t value	P value
Post-acute care program (X)	a1	-2.617	-2.032	0.043	é	0.173	0.878	0.381	0.231	1.309	0.191
12 th week cognitive function (M)					b1	0.021	0.737	0.461	0.049	1.921	0.055
Acute lengths of stay (W)	a2	-0.214	0.049	<0.001	b2	0.074	2.451	0.015	0.094	0.027	0.001
X*W	a3	0.204	3.102	0.002							
M*W					b3	-0.003	-2.310	0.021	-0.004	-3.199	0.002
		R ² =0.041				R ² =0.029			R ² =0.036		
		F(3, 472)=6.647 < 0.05				F(4, 471)=3.534 < 0.05			F(4, 471)=4.414 < 0.05		