

Supplementary Table 1: Diagnostic codes and subcodes for patients included in the study between January 2020 and December 2021.

	Diagnostic code in ANZICS-APD	Diagnostic subcode for patients without COVID-19	Diagnostic subcode for patients with COVID-19*
Viral pneumonia	213	213.01	213.02
ARDS	204	204.01	204.22
ANZICS – Australia New Zealand Intensive Care Society; APD – adult patient database; ARDS – acute respiratory distress syndrome * The patients were considered highly likely to be positive for COVID-19 (“Suspected or confirmed pandemic infection”), as ANZICS does not have this information but infers it.			

Supplementary Table 2: Missing data comparison for patients with and without available CFS scores.

Variable	Frailty data present (n=3064)	Missing frailty data (n=645)	p-value
PerCI, n (%)	847 (27.6%)	134 (20.9%)	<0.001
Indigenous, n (%)	78 (2.5%)	10 (1.6%)	0.32
Male sex, n (%)	1879 (56.2%)	382 (59.6%)	0.41
Age (years) (median [IQR])	56.9 [44.7, 68.2]	59.0 [48.3, 69.9]	0.007
Hospital admission source, n (%)			
- Home	2473 (80.7%)	539 (84.1%)	0.012
- Other acute hospital (not ICU/ED)	129 (4.2%)	16 (2.5%)	
- Nursing home or chronic care	15 (0.5%)	8 (1.2%)	
- Other hospital ICU	257 (8.4%)	38 (5.9%)	
- Other hospital Emergency department	151 (4.9%)	28 (4.4%)	
- Rehabilitation	3 (0.1%)	2 (0.3%)	
ICU admission source, n (%)			
- Emergency department	1191 (38.9%)	280 (43.7%)	<0.001
- Ward	1483 (48.4%)	278 (43.4%)	
- Other hospital	103 (3.4%)	33 (5.1%)	
- Other hospital ICU	272 (8.9%)	36 (5.6%)	
- Operating theatre / Recovery	1 (0.0%)	0 (0)	
- Direct admit	14 (0.5%)	14 (2.2%)	
Documented co-morbidities, n (%)			
- Chronic respiratory condition	200 (6.5%)	15 (2.3%)	<0.001
- Chronic cardiovascular condition	179 (5.8%)	19 (3.0%)	0.003
- Chronic renal failure	74 (2.4%)	6 (0.9%)	0.019
- Chronic liver disease	21 (0.7%)	6 (0.9%)	0.50
- Diabetes mellitus	911 (29.7%)	89 (13.9%)	<0.001
- Immune suppressive therapy	142 (4.6%)	27 (4.2%)	0.64
- Lymphoma	13 (0.4%)	1 (0.2%)	0.31
- Leukaemia	26 (0.8%)	6 (0.9%)	0.83
- Metastatic cancer	24 (0.8%)	4 (0.6%)	0.67
- Delirium, n (%)	259 (8.5%)	8 (1.2%)	<0.001
Organ failure score			
- APACHE III score (mean [SD])	50.0 [20.0]	50.4 [21.7]	0.47
- ANZROD (%) (mean [SD])	9.5 [12.2]	10.1 [12.1]	0.61
ICU admission post MET call, n (%)	1105 (36.3%)	182 (28.4%)	<0.001
Cardiac arrest, n (%)	6 (0.2%)	0 (0)	0.004
Treatment limitation, n (%)	255 (8.4%)	55 (8.6%)	0.40
Pre-ICU (days) (median [IQR])	12.4 [4.0, 56.8]	9.5 [3.7, 63.6]	0.31
ICU supports			
Mechanical ventilation (MV), n (%)	1306 (43.1%)	195 (32.0%)	<0.001
Non-invasive ventilation (NIV), n (%)	1260 (41.8%)	203 (33.4%)	<0.001
Vasopressor and inotropes, n (%)	1188 (39.2%)	200 (32.8%)	0.003
Renal replacement therapy, n (%)	180 (6.0%)	33 (5.5%)	0.63
ECMO, n (%)	104 (3.5%)	9 (1.5%)	0.011
Tracheostomy, n (%)	186 (6.2%)	13 (2.2%)	<0.001
CFS – clinical frailty scale, SD – standard deviation, IQR – interquartile range, MET – medical emergency team, APACHE - Acute Physiology and Chronic Health Evaluation, ED – emergency department, ICU – intensive care unit, ROD – risk of death, ANZROD – Australia New Zealand risk of death, PerCI - Persistent critical illness			

Supplementary Table 3a: Baseline characteristics of patients with and without Persistent critical illness (PerCI), for clinical frailty scale (CFS) categories CFS-1-2 and CFS-3-4.

Variable	CFS-1-2 (n=1123)			CFS-3-4 (n=1575)		
	Patients without PerCI (n=835)	Patients with PerCI (n=288)	p-value	Patients without PerCI (n=1109)	Patients with PerCI (n=466)	p-value
Male sex, n (%)	523 (62.6%)	203 (70.5%)	0.012	656 (59.2%)	295 (63.3%)	0.12
Indigenous status, n (%)	24 (2.9%)	8 (2.8%)	0.92	32 (2.9%)	9 (1.9%)	0.46
Age, (years), median (IQR)	46.5 (34.1, 56.6)	54.0 (43.5, 63.9)	<0.001	59.1 (47.5, 69.3)	61.7 (51.7, 69.8)	0.010
Hospital admission source, n (%)						
- Home	715 (85.6%)	191 (66.3%)	<0.001	950 (85.7%)	337 (72.3%)	<0.001
- Other acute hospital	103 (12.4%)	90 (31.3%)		148 (13.3%)	126 (26.9%)	
- Nursing home or chronic care	0 (0)	1 (0.3%)		1 (0.1%)	0 (0)	
- Rehabilitation	0 (0)	0 (0)		2 (0.2%)	0 (0)	
- Missing	17 (2.0%)	6 (2.1%)		8 (0.7%)	4 (0.9%)	
ICU admission source, n (%)						
- Emergency department (ED)	355 (42.5%)	93 (32.3%)	<0.001	459 (41.4%)	138 (29.6%)	<0.001
- Ward	411 (49.2%)	116 (40.3%)		340 (45.1%)	40 (43.0%)	
- Other hospital (ED and ICU)	62 (7.4%)	77 (26.7%)		179 (23.7%)	30 (32.3%)	
- Operating theatre / Recovery	1 (0.1%)	0 (0)		0 (0)	0 (0)	
- Direct admit	6 (0.7%)	2 (0.7%)		2 (0.2%)	2 (0.4%)	
Documented co-morbidities, n (%)						
- Chronic respiratory condition	25 (3.0%)	7 (2.4%)	0.62	72 (6.5%)	17 (3.6%)	<0.001
- Chronic CVS condition	18 (2.2%)	6 (2.1%)	0.94	71 (6.4%)	16 (3.4%)	0.019
- Chronic renal failure	3 (0.4%)	1 (0.3%)	0.98	30 (2.7%)	10 (2.1%)	0.52
- Chronic liver disease	0 (0)	1 (0.3%)	0.09	7 (0.6%)	4 (0.9%)	0.62
- Diabetes mellitus	138 (16.9%)	63 (22.1%)	0.05	375 (35.6%)	181 (41.5%)	0.031
- Immune suppressive therapy	12 (1.4%)	30 (11.0%)	0.45	58 (5.2%)	26 (5.6%)	0.78
- Lymphoma	0 (0)	0 (0)	-	5 (0.5%)	3 (0.6%)	0.62
- Leukaemia	1 (0.1%)	1 (0.3%)	0.43	8 (0.7%)	5 (1.1%)	0.48
- Metastatic cancer	2 (0.2%)	0 (0)	0.41	9 (0.8%)	2 (0.4%)	0.41
- Obese (BMI \geq 30 kg.m ⁻²)	410 (49.1%)	100 (34.7%)	<0.001	440 (39.7%)	161 (34.5%)	0.11
- Delirium	16 (1.9%)	44 (15.3%)	<0.001	279 (25.2%)	124 (26.6%)	<0.001
- Pregnancy status	31 (3.7%)	6 (2.1%)	<0.001	31 (2.8%)	4 (0.9%)	0.004
Pre-ICU (days) (median [IQR])	13.6 (4.6, 54.4)	6.7 (0.9, 36.8)	<0.001	13.5 (4.6, 59.3)	9.9 (1.6, 57.9)	0.004
ICU discharge delay, (median [IQR])	4.0 (2.2, 7.2)	5.0 (2.9, 11.4)	<0.001	4.0 (1.9, 7.1)	4.5 (1.5, 8.2)	0.48
Organ failure scores						
- APACHE III (mean [SD])	39.6 (16.6)	53.2 (18.4)	<0.001	49.9 (11.9)	57.3 (18.1)	<0.001
- ANZROD (%) (mean [SD])	4.6 (6.0)	9.1 (10.6)	<0.001	9.3 (11.9)	11.1 (10.7)	<0.001
ICU admission post MET call, n (%)	282 (34.0%)	88 (30.9%)	0.34	432 (39.3%)	164 (35.5%)	0.16
Treatment limitations	7 (0.8%)	6 (2.1%)	0.13	86 (7.8%)	19 (4.1%)	0.026
Cardiac arrest, n (%)	0 (0)	0 (0)	-	4 (0.4%)	0 (0)	0.39
ICU Supports, n (%)						
Mechanical ventilation	168/823 (20.4%)	247/286 (86.4%)	<0.001	365/1097 (33.3%)	402/465 (86.5%)	<0.001
Non-invasive ventilation	270/823 (32.8%)	111/282 (39.4%)	0.046	476/1093 (43.5%)	234/461 (50.8%)	0.009
Vasopressor and inotropes	130/824 (15.8%)	216/285 (75.8%)	<0.001	339/1097 (30.9%)	369/464 (79.5%)	<0.001
Renal replacement therapy	6/822 (0.7%)	39/282 (13.8%)	<0.001	29/1091 (2.7%)	85/461 (18.4%)	<0.001
ECMO	6/822 (0.7%)	45/282 (16.0%)	<0.001	6/1092 (0.5%)	45/461 (9.8%)	<0.001
Tracheostomy	3/822 (0.4%)	55/281 (19.6%)	<0.001	3/1090 (0.3%)	111/458 (24.2%)	<0.001
CFS – clinical frailty scale, SD – standard deviation, IQR – interquartile range, BMI – body mass index, MET – medical emergency team, APACHE - Acute Physiology and Chronic Health Evaluation, ED – emergency department, ICU – intensive care unit, ANZROD – Australia New Zealand risk of death. ECMO – extracorporeal membrane oxygenation						


Supplementary Table 3b: Baseline characteristics of patients with and without Persistent critical illness (PerCI), for clinical frailty scale (CFS) categories CFS-5-6 and CFS-7-8.

Variable	CFS-5-6 (n=299)			CFS-7-8 (n=67)		
	Patients without PerCI (n=219)	Patients with PerCI (n=80)	p-value	Patients without PerCI (n=54)	Patients with PerCI (n=13)	p-value
Male sex, n (%)	105 (47.9%)	50 (62.5%)	0.026	36 (66.7%)	11 (84.6%)	0.20
Indigenous status, n (%)	1 (0.5%)	0 (0)	0.75	4 (7.4%)	0 (0)	0.31
Age, (years), median (IQR)	72.8 (64.5, 81.6)	68.3 (58.0, 75.4)	<0.001	68.7 (55.8, 78.5)	62.1 (52.1, 70.3)	0.20
Hospital admission source, n (%)						
- Home	185 (84.5%)	48 (60.0%)	<0.001	41 (75.9%)	6 (46.2%)	0.003
- Other acute hospital	27 (12.3%)	29 (36.4%)		8 (14.9%)	7 (53.9%)	
- Nursing home or chronic care	7 (3.2%)	2 (2.5%)		4 (7.4%)	0 (0)	
- Rehabilitation	0 (0)	0 (0)		1 (1.9%)	0 (0)	
- Missing	0 (0)	1 (1.3%)		0 (0)	0 (0)	
ICU admission source, n (%)						
- Emergency department (ED)	99 (45.2%)	21 (26.3%)	<0.001	24 (44.4%)	2 (15.4%)	0.018
- Ward	111 (50.7%)	35 (43.8%)		23 (42.6%)	5 (38.5%)	
- Other hospital (ED and ICU)	8 (3.6%)	24 (30.1%)		6 (11.1%)	6 (46.2%)	
- Operating theatre / Recovery	0 (0)	0 (0)		0 (0)	0 (0)	
- Direct admit	1 (0.5%)	0 (0)		1 (1.9%)	0 (0)	
Documented co-morbidities, n (%)						
- Chronic respiratory condition	48 (21.9%)	9 (11.3%)	0.038	18 (33.3%)	4 (30.8%)	0.86
- Chronic CVS condition	45 (20.5%)	7 (8.8%)	0.017	15 (27.8%)	1 (7.7%)	0.13
- Chronic renal failure	24 (11.0%)	2 (2.5%)	0.022	4 (7.4%)	0 (0)	0.31
- Chronic liver disease	5 (2.3%)	2 (2.5%)	0.91	2 (3.7%)	0 (0)	0.48
- Diabetes mellitus	97 (45.8%)	34 (43.6%)	0.74	22 (41.5%)	1 (7.7%)	0.022
- Immune suppressive therapy	25 (11.4%)	9 (11.3%)	0.97	5 (9.3%)	1 (7.7%)	0.040
- Lymphoma	3 (1.4%)	1 (1.3%)	0.94	0 (0)	1 (7.7%)	0.62
- Leukaemia	6 (2.7%)	2 (2.5%)	0.91	3 (5.6%)	0 (0)	0.39
- Metastatic cancer	7 (3.2%)	1 (1.3%)	0.36	2 (3.7%)	1 (7.7%)	0.53
- Obese (BMI \geq 30 kg.m ²)	55 (25.1%)	32 (40.0%)	0.028	19 (35.2%)	7 (53.8%)	0.41
- Delirium	23 (10.5%)	18 (22.5%)	0.006	7 (13.0%)	3 (23.1%)	0.50
- Pregnancy status	0 (0)	0 (0)	-	0 (0)	0 (0)	-
Pre-ICU (days) (median [IQR])	17.9 (6.4, 75.9)	5.5 (0.1, 69.4)	<0.001	15.6 (5.5, 88.9)	13.6 (0.1, 61.6)	0.20
ICU Discharge delay	4.2 (1.3, 8.8)	4.9 (1.9, 9.9)	0.57	4.8 (2.0, 11.5)	0 (0, 2.9)	0.006
Organ failure scores						
- APACHE III (mean [SD])	62.1 (20.6)	62.8 (20.3)	0.61	62.5 (24.0)	66.5 (26.8)	0.80
- ANZROD (%) (mean [SD])	20.1 (18.4)	16.4 (14.6)	0.10	23.8 (24.0)	19.6 (17.3)	0.86
ICU admission post MET call, n (%)	91 (41.6%)	25 (32.1%)	0.14	18 (33.3%)	5 (38.5%)	0.73
Treatment limitations	94 (43.0%)	11 (13.8%)	<0.001	30 (55.6%)	2 (14.4%)	0.016
Cardiac arrest, n (%)	0 (0)	0 (0)	-	1 (1.9%)	1 (7.7%)	0.54
ICU Supports, n (%)						
Mechanical ventilation	37/213 (17.4%)	61/80 (76.3%)	<0.001	14/53 (26.4%)	12/13 (92.3%)	<0.001
Non-invasive ventilation	102/213 (47.9%)	38/77 (49.4%)	0.83	27/54 (50.0%)	2/13 (15.4%)	0.024
Vasopressor and inotropes	48/214 (22.4%)	58/80 (72.5%)	<0.001	17/53 (32.1%)	11/13 (84.6%)	<0.001
Renal replacement therapy	9/213 (4.2%)	8/77 (10.4%)	0.048	1/53 (1.9%)	3/413 (23.1%)	0.004
ECMO	0/213 (0)	2/77 (2.6%)	0.018	0 (0)	0 (0)	-
Tracheostomy	2/213 (0.9%)	11/77 (14.3%)	<0.001	0 (0)	1/13 (7.7%)	0.042
CFS – clinical frailty scale, SD – standard deviation, IQR. – interquartile range, BMI – body mass index, MET – medical emergency team, APACHE - Acute Physiology and Chronic Health Evaluation, ED – emergency department, ICU – intensive care unit, ANZROD – Australia New Zealand risk of death. ECMO – extracorporeal membrane oxygenation						

Supplementary Table 4: Adjusted odds of hospital mortality in patients with and without persistent critical illness. The graphical version is presented in Figure 1.


Predictor	Patients without Persistent Critical illness		Patients with Persistent Critical illness	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Logistic Regression Model 1 – Frailty assessed using CFS categories				
CFS 1-2	Reference		Reference	
CFS 3-4	3.25 (1.86-5.66)	<0.001	2.09 (1.44-3.04)	<0.001
CFS 5-6	5.54 (2.88-10.66)	<0.001	1.87 (1.03-3.40)	0.041
CFS 7-8	11.02 (4.52-26.85)	<0.001	7.01 (1.94-25.26)	0.003
Male sex	1.77 (1.20-2.59)	<0.001	1.31 (0.93-1.85)	0.12
ANZROD	1.08 (1.07-1.10)	<0.001	1.05 (1.04-1.07)	<0.001
Logistic Regression Model 2 – Frailty assessed using CFS as a continuous variable				
CFS	1.46 (1.29-1.64)	<0.001	1.30 (1.14-1.49)	<0.001
ANZROD	1.08 (1.07-1.10)	<0.001	1.05 (1.04-1.07)	<0.001
Male sex	1.82 (1.24-2.67)	0.002	1.32 (0.94-1.86)	0.11
CFS = clinical frailty scale, ANZROD = Australia and New Zealand risk of death				

Supplementary Figure 1: Flow diagram demonstrating patient inclusion.



CFS – clinical frailty scale, PerCI - Persistent critical illness
Non-frail (CFS 1-4), Frail (CFS 5-8)

Supplementary Figure 2: The relationship between frailty and mortality in those with and without PerCI, before and after adjusting for ANZROD and male sex, using AUROC (95%CI). The Clinical Frailty Scale (CFS) was treated as a continuous variable.



Comparison between unadjusted and adjusted PerCI vs. no PerCI are both <0.001
* Adjusted CFS for ANZROD and male sex

Supplementary Figure 3: Proportion of patients with ICU length of stays and those who died in hospital, based on CFS categories. *Error bars are standard errors of the mean.

