

The experience of family carers for people with moderate to advanced dementia within a domestic home setting: a systematically constructed narrative synthesis

Charles James PhD, Catherine Walshe PhD, Caroline Swarbrick PhD

Abstract

Background

Caring for someone with moderate to advanced dementia within a domestic home setting can be burdensome and time-consuming. To ensure the effectiveness of care planning and delivery, especially towards the end of life, understanding the nature and impact of such caregiving on the family carer is important. Synthesising existing research will allow greater insight into this experience.

Review question

'How do family carers describe their experience of providing home-based care for people with moderate to advanced dementia?'

Design

A narrative synthesis of qualitative research exploring the experiences of people with moderate to advanced dementia and their family carers was conducted. Databases (MEDLINE, CINAHL, EMBASE, PsychINFO, Web of Science and Academic Search Complete) were searched from 1984 to 2020. Similarities and differences between papers were grouped using textual narrative synthesis.

Findings

This paper reports findings from a PhD thesis (James, 2021). Included papers (n=17) incorporated those focused on caring for people with advanced dementia (n=8), and those with moderate dementia (n=9). Family carers reported an experience of loss, burden, and

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

grief. Experiences of internal conflict also led to psychological distress. These experiences had a negative impact on the general health of the family carer. Improvement in the family carer's inter-relationships and the feeling of being useful or having a sense of meaning were reported as positive aspects of caregiving.

Conclusion

A complete description of family carers' experiences may be inadequate in conveying an acknowledgement of dementia as an illness within the domestic home. Their justifications and determinants for balancing family carers' challenges and distress to morals also remain unclear. Further research is required to ascertain how family carers may proportionally balance their personified loss with their personified value earlier in the disease trajectory.

Introduction

Accounts of caring for someone living with dementia are mostly rooted within negative experiences, such as the associated burden (Lindeza et al., 2020; Bieber et al., 2019; Farina et al., 2017). Burden, in this respect, may be described as the impact of the difficulties associated with the dementia diagnosis, which increases the level of uncertainty about the future for both the person with the disease and their family (Galvin & Sadowski, 2012; Phillips, 2011). However, not all family carers, who are mostly unpaid, perceive their experiences as burdensome.

Being diagnosed with dementia is often regarded as a burden which increases the level of uncertainty about the future for both the person with the disease and their family (Galvin & Sadowski, 2012; Phillips, 2011). As a factor which also increases the social implications for the people diagnosed and their families, given their difficulty in preparedness, receiving a diagnosis of dementia is often difficult and sometimes described in terms of a feeling of shame and loss of oneself (Xanthopoulou & McCabe, 2019; van Gennip et al., 2016; Lee & Weston, 2011; Aminzadeh et al., 2007).

Life expectancy in dementia varies between individuals. Although a duration of less than 10 years after diagnosis is regarded as a common assumption (Joling et al., 2020), it is unpredictable in comparison with other terminal diseases, such as cancer (Vestergaard et al., 2020; Hall & Sikes, 2018; Arcand, 2015; Harris, 2007). The duration of caring for someone with dementia is therefore likely to last longer than in most other terminal illnesses. The level of decline in dementia is often described in stages, although the duration of each stage is usually unknown. While a cure or effective prevention for dementia is currently unavailable, its stages represent the degree of its severity and are usually identified between the early to the most severe level of decline (Alzheimer's Society, 2015; Schmidt, 2014; Auer & Reisberg, 1997). A progressive decline in self-care activities of daily living, such as dressing, bathing, and toileting by the person diagnosed is often useful for its stage categorisation (Kumar et al., 2020; Mlinac & Feng, 2016; Marshall et al., 2012).

The responsibilities of family carers usually increase in line with its severity and vary according to the needs of the person cared for (Mesterton et al., 2010). Fulfilling a home home-based care wish in dementia often relies on the willingness of a family carer to take on the role. In some cases, such care provision may be viewed as the family carer's choice. However, the assumption of the role is mostly associated with a lack of choice (Pertl et al., 2019; Reinhard et al., 2012; Schulz et al., 2012; Al-janabi et al., 2018). A view of family caregiving as performing a moral obligation is therefore plausible.

Given evidence suggests that around 36% of family carers spend more than 100 hours per week, while about 40% provide round-the-clock care (Dementia Carers Count, 2019). In some cases, instances of further unpredictable difficulties, such as behavioural changes are likely over the disease trajectory (van Wijngaarden et al., 2018; Schulz & Eden, 2016). The likelihood of this long caregiving combined with a possible lack of choice leading to family carers' increased burden and depression is high (Greenwood & Smith, 2019; Karg et al., 2018; Schulz et al., 2012).

Experiences of decline in dementia are unique within its different stages. Within the moderate stage, for example, variations of decline in individual capabilities are possible (Clemmensen et al., 2016; Alzheimer's Society, 2015). This may be experienced earlier or later by some people, or not at all by others. It is also possible for stages to overlap (Alzheimer's Society, 2015), which suggests the likelihood of a variation of family carers' experiences while caregiving within dementia stages. The focus of this review is to explore and synthesise the experiences of family carers while caregiving for people living with dementia within a home setting.

Aim & objectives

To understand the experiences of family carers who provide care for people with moderate to advanced dementia within a domestic home setting.

Methods

Design

A textual narrative synthesis approach was adopted due to its appropriateness in ensuring that similarities and differences between multiple studies were systematically explored (Popay et al., 2006). The stages suggested by Popay et al. (2006), were followed, including identifying the review focus, specifying the review question, identifying studies to include data extraction and quality appraisal, synthesising of the findings, and reporting of findings. The study protocol is published (James et al., 2020).

Specifying the review question and inclusion criteria

Family carers are people who may not reside with the care recipient, but provide unpaid and significant care (Lindeza et al., 2020; Evans et al., 2019; Woodman et al., 2016). Whilst they are often crucial to the quality of life of the care recipient, they are also regarded as the invisible second patient due to their own care needs (Karg et al, 2018; Sanders, 2016; Brodaty & Donkin, 2009). A purposive search of the key terms, dementia and family carers, and their variations was conducted for the comprehensiveness of the inclusion criteria (Table 1).

Table 1: Review inclusion and exclusion criteria

| Inclusion Criteria | Exclusion Criteria |
|--|---|
| Type of papers: peer-reviewed empirical papers, qualitative papers including case studies and mixed-methods (both telephone and postal surveys are acceptable only if open-ended questions are asked), qualitative evidence collected from interviews, observations | Type of papers: Papers that are not primary research e.g. systematic reviews, meta-analysis |
| Language of papers: English | Language of papers: Papers in other languages |
| Date: Papers published from 1984 onwards | Date: Papers were written before 1984 |
| Population of focus: unpaid family members or family carer | Population of focus: Population of focus relates to other groups of carers e.g. paid carers, professionals |
| Age group: 18 and above | Age group: Below 18 |
| Primary illness of interest: moderate to advanced dementia as defined by the authors of identified studies in the abstract or full paper. | Primary illness of interest: Other illnesses |

| | |
|--|--|
| Setting: domestic home setting, unpaid home-based care | Setting: Other formal care establishments where care provision is paid for |
| Focus of papers: current experiences or views or needs of family carer | Focus of papers: Focus on the views of others, or where death has already occurred. |
| Experience: Subjective experiences, such as family carers' views, opinion, attitudes, and the reasons for this, with respect to home-based caregiving for people with moderate to advanced dementia. | Experience: other non-family carers' views |
| Research Type: Research adopting either qualitative or mixed methods (described in terms of exposure to specific elements of caregiving responsibilities leading to a noticeable change in behaviour or wellbeing of the sample, accounts of the duration of time spent caregiving and family carers' feelings or emotions during care provision were also of interest. | Research Type: other types of research |

Search strategy and process

Searches were conducted in MEDLINE Complete, CINAHL, EMBASE, PsycINFO, Web of Science, and Academic Search Complete. Websites and grey literature were also searched. For all searches, the terms and strategy used a broad range of terms and relevant keywords related to dementia, caregivers, and qualitative studies. These were checked by CW and CS, who were familiar with the domain of dementia and care provision. The search strategy was also tailored for use with each database, using Boolean operators, truncations, and Medical Subject Heading (MESH) as appropriate for each database (Table 2).

Table 2: Search terms and strategy

| Search terms and strategy |
|--|
| MH "Dementia+" OR TX dement* OR TX alzheimer * OR TX "lew* bod*" OR TX FTLD OR FTD OR frontotemporal |
| AND TX carer* OR TX caregiv* OR TX care-giver OR TX spouse-caregiver* |
| AND MH "Qualitative Research+" OR MH "qualitative studies" OR MH "Focus Groups" OR MH "exploratory+" OR MH "qualitative interview+" OR MH "ethnography+" OR MH "ethnographic research" OR MH "content analysis" OR MH "Grounded theory" |

The decision to include all papers published from 1984 to 2020 was based on the reported increase in diagnosis and people caring informally (Lewis et al., 2014). Supplementary searches of the reference lists of identified studies were also conducted. The list of studies returned from the searches was imported into Endnote, an online reference management software (Lorenzetti & Ghali, 2013), to remove duplicates and ensure the validity and reliability of the review process' (Kwon et al., 2015) (See Table 2 above).

Data extraction

Key data such as author's name, publication origin and year, setting, population and sample size, aims and objectives, data collection method, and main findings, which show the family carers' actions were extracted from the included papers.

Quality and bias assessment

A non-discriminatory tool developed by Hawker et al. (2002) and adapted by Lorenc et al. (2014) was used in assessing the quality of the included papers. Based on the adaptability of its structure to a variety of methodologically distinctive designs, the purpose of this tool was to assess the understanding of quality within the synthesis, rather than to exclude papers of lower quality. The studies included in this review were assessed by two study supervisors, CW and CS. The purpose was to check the results of the appraisal process and ensure the appropriateness of the selected tool as well as the overall process of quality assessment (See Supplementary Table 1).

Approach to synthesis

A textual narrative synthesis approach was adopted in accordance with guidance from Popay et al. (2006), ensuring the exploration of synthesised words or texts from systematically selected studies using their similarities and differences. A total of 2248 full-text articles were reviewed, with seventeen papers included in the narrative synthesis (Fig 1). Eight studies presented data from those only caring for people with advanced dementia, nine included data from those caring for people with both moderate dementia and advanced dementia. While the review was not focused on establishing a comparison of experiences between genders, it is noteworthy that two studies were dedicated solely to accounts from female family carers (De Silva & Curzio, 2009; de La Cuesta-Benjumea, 2011).

Review findings




Fig 1: Sources of studies included (PRISMA diagram adapted from Page et al., 2020).

The review findings are presented in three stages. Systematically extracted data are first tabulated to display the similarities and differences between the findings of each selected paper (Popay et al., 2006) (Table 3).

Table 3: Characteristics of included studies

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|-----------|--|---|---|---|---|---------------|---------------|
| Adams and Sanders (2004) [USA] | Homebased | Sample: Informal caregivers (n=74) Sampling method: Convenience sampling Stage: Moderate to advanced | To explore self-reported losses, grief reactions and depressive symptoms among a sample of caregivers of persons with Alzheimer's disease or other progressive dementia | Type: Mixed method Design: Cross-sectional | Data collection: Open-ended postal survey Analysis: Cross-sectional analysis | a) Caregivers for individuals within the late stage of dementia reported significantly more symptoms of grief and depression than those in the early or middle stages; b) The late-stage experience mostly resembled bereavement. | 29 | B |
| Brunton et al. (2008) [New Zealand] | Homebased | Sample: Family caregivers 1 st stage: (n=5) 2 nd stage: (n=64) Sampling method: | To explore the lived experience and perceptions of a sample of caregivers who are providing informal care 24hr per day, seven days per week for those with Alzheimer's | Type: Mixed method Design: Exploratory | Data collection: Semi-structured interview and questionnaires (using open-ended questions) | a) High cost of caring is reported as a form of negative impact on caregiver's physical and psychological health; b) Difficulty in asking for help from family; c) Caregivers were concerned about finances and fears for the future; d) Caregivers reported also reported the | 34 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|---------------------|---|---|--|---|--|---------------|---------------|
| | | Purposive, non-probability Stage: Stage 2 to Stage 5 | disease in New Zealand. | | Analysis: Thematic analysis | need to be free, need for a time out, need for a role change and a need to be someone else. | | |
| Chan et al. (2010) [Hong Kong] | Community homebased | Sample: Informal caregivers (n=27) Sampling method: Purposive Stage: Moderate to advanced | To report the lived experience of caregivers of persons with dementia in Hong Kong, and to explore their service needs. | Type: Qualitative Design: Focus group | Data collection: Focus group interviews Analysis: Colaizzi's phenomenological analysis | a) Caregivers were confused about the nature of their care recipients' disease; b) Negative report of caregiving in terms of grief, sadness, guilt, anger and fear; c) The needs of caregivers changed in line with the care recipient's disease progression; d) Caregiving responsibilities increased as the illness progressed; (e) Difficulty in coping with increasing care recipient's behavioural changes; (f) Increased caregiving responsibilities conflicted with other social roles, such as employment and other social activities; g) Burden reduces in the advanced | 36 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--------------------------------|-----------|--|--|---|---|---|---------------|---------------|
| | | | | | | stage of dementia as role acceptance increases. | | |
| Colling (2004) [USA] | Homebased | <p>Sample: Family caregivers</p> <p>Moderate (n=16), Advanced (n=19)</p> <p>Sampling method: Purposive</p> <p>Stage: Moderate to advanced</p> | To describe how the experience of passivity was for the caregiver and the person with Alzheimer's disease. | <p>Type: Qualitative</p> <p>Design: Phenomenology</p> | <p>Data collection: Semi-structured interviews</p> <p>Analysis: content analysis (using Colaizzi's thematic extraction)</p> | a) Passivity increased over time as the disease worsened;b) Experience of emotions such as fatigue, sadness, tears, frustration and a sense of helplessness; c) Caregivers of moderately impaired persons found it stressful to cope with lifestyle changes associated with caregiving; d) Caregivers of severely impaired persons expressed the most sadness associated with experiencing the process of cognition deterioration in their loved ones; e) at the moderate stage found it stressful to cope with lifestyle changes associated with caregiving, a sense of helplessness and frustration at watching behavioural changes in the care recipient | 35 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|---|-----------|---|--|--|---|---|---------------|---------------|
| de La Cuesta-Benjumea (2011) [Spain] | Homebased | Sample: Female family caregivers (n=23) Sampling method: Purposeful and theoretical Stage: Advanced | To explore strategies women caregivers of relatives with advanced dementia use to rest from caregiving | Type: Qualitative Design: Grounded theory | Data collection: Semi-structured interviews Analysis: Grounded theory (constant comparison analysis) | a) Taking leave from the life of caregiving is used by the caregivers as a strategy to rest from their caregiver selves; (b) Caregivers did not perceive respite to mean resting or being idle. For them, respite is considered as doing something different, and by thinking and relating to others in different ways than those linked to caregiving. | 36 | A |
| De Silva and Curzio (2009) [UK] | Homebased | Sample: Female family caregivers (n=10) Sampling method: Purposive Stage: Advanced | To explore the life experiences of family carers in the transition from home into long-term care | Type: Qualitative Design: Phenomenology | Data collection: One-to-one interviews Analysis: Thematic analysis | a) Deterioration of behaviour is a major reason for care recipient transitioning into long-term care; (b) Although alternatives to long-term care were offered before placement, several did not receive much support and felt isolated; c) Spousal caregivers reported great sadness in placing their loved one; d) Difficulty in finding the right place; e) Severity of dementia and the | 29 | B |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|---|-----------|--|---|--|--|---|---------------|---------------|
| | | | | | | complex health needs of a loved one left caregivers no other choice but long-term care; f) Experience of guilt or emotional distress during the transition. | | |
| Hirschman et al. (2006) [USA] | Homebased | Sample: Family caregivers (n=30) Sampling method: Purposive Stage: Advanced | To identify what standard of decision making a family member uses when making medical decisions for their relative with advanced dementia | Type: Qualitative Design: Qualitative | Data collection: Semi-structured interviews Analysis: Thematic analysis | a) Making medical decisions is difficult for caregivers, and there is no guidance as to how this is done; b) Either substituted judgement (what the care recipient would have wanted) (n=2) or best interest (opinion of the caregiver for deciding what is best for the care recipient) (n=17). In some cases, decisions are made by combining both as care recipients (n=11) often fail to have discussions about end of life with their family; c) Barriers highlighted include: avoiding the topic, waiting too long and denial of dementia by the care | 35 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|-----------|---|---|--|---|--|---------------|---------------|
| | | | | | | recipient and sometimes the caregivers themselves. | | |
| Idura et al. (2018) [Malaysia] | Homebased | Sample: Family caregivers (n=12) Sampling method: Purposive Stage: Moderate to advanced | To describe the experience of Malaysian family caregivers' experiences in providing care to their family members with moderate to advanced dementia | Type: Qualitative Design: Qualitative | Data collection: In-depth semi-structured interviews Analysis: Thematic analysis | a) Increased severity did not increase caregivers' burden due to acceptance of the caregiving role; b) Caregivers' grief increases with severity; c) Burden increases at the earlier stages; d) Caregivers felt it was their responsibility to provide care; e) Family support as a possible reason for a reduced burden; f) Family support is a positive experience in helping to improve relationships among family members; g) Family support was not always available. | 35 | A |
| Karger (2018) [Germany] | Homebased | Sample: Relatives (n=20) Sampling method: Purposive | To gain insight into families and professionals' understanding of the emotional experience in patients with | Type: Qualitative Design: Focus group | Data collection: Group interviews Analysis: | a) Relatives perceived emotionality as one of the essential characteristics retained despite the illness; b) loss of emotional experience increased with loss of cognitive capacity | 32 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|------------------------------------|-----------|--|---|--|--|---|---------------|---------------|
| | | Stage: Advanced | advanced Alzheimer's disease | | Qualitative content analysis and framework analysis | and verbal expression; c) shared memories faded with illness progression; d) memories forgotten no longer held any emotional significance for the care recipient; e) Perceived discontinuity of patient's emotional responses contrary to relatives' expectations creates ambiguity; f) Patients' behaviour which is different to expectations is a distressing experience for the relatives. | | |
| Lamahewa et al. (2018) [UK] | Homebased | Sample: Family caregivers Current (n=6) Sampling method: Purposive | To explore the difficulties in decision making for family carers and practitioners at the end of life | Type: Qualitative Design: Focus group | Data collection: Semi-structured interviews Analysis: Thematic analysis | Making decisions is particularly difficult at the end of life and associated with caregivers' unpreparedness | 32 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|---|---|---|---|--|---|---------------|---------------|
| | | Stage: Advanced | | | | | | |
| Lethin et al. (2016) [Sweden] | Homebased (also <i>nursing home</i>) | Sample: Family caregivers (n=23) Sampling method: Strategic Stage: <i>Advanced (includes accounts of early stages)</i> | To investigate family caregivers' experiences of formal care when caring for a person with dementia through the stages of the disease | Type: Qualitative Design: Focus group | Data collection: Focus group interviews Analysis: Content analysis | a) Caregivers' feeling of inadequacy and striving to remain in control, while avoiding being a burden and struggling to avoid dependence on formal care; b) Worry about the future and the capacity to continue caregiving; c) Feelings of being alone and isolated because formal care was not as supportive as expected; d) Increasingly isolated when friends and family no longer came to visit | 36 | A |
| Lindauer et al. (2016) [USA] | Homebased | Sample: Family caregivers (n=11) Sampling method: | To explore the meaning AA family caregivers ascribed to dementia-related changes in care recipients | Type: Qualitative Design: Interpretative Phenomenolo | Data collection: In-depth interviews Analysis: Interpretative Phenomenolo | a) Dementia-related changes meant that caregivers had to hang on to the care recipient for as long as possible; b) Caregiver's recognition that the valued care recipients were changed but still here | 36 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|------------------------------------|---|--|---|--|---|--|---------------|---------------|
| | | Purposive Stage: Moderate to <i>advanced</i> | | gical Approach | gical Approach (IPA) | and worthy of respect and compassion; c) Valued care recipients were changed, but still here, was regarded as a paradox; d) Family values, shaped by historical oppression, influenced caregiving meanings; e) Caregivers did not consider they were grieving, as their care recipient was <i>still here</i> . | | |
| Moore et al. (2017) [UK] | Homebased (Part of a larger study including care-home as the setting) | Sample: Family carers (n=6) Sampling method: Stratified Stage: Advanced | To understand the experiences of carers during advanced dementia and to explore the links between mental health and experiences of end of life care. | Type: Mixed methods Design: Longitudinal cohort study | Data collection: In-depth interviews Analysis: Thematic analysis | a) Difficulty in understanding dementia progression and being involved during advanced stages; b) Placing care recipient in a care home is perceived as a loss of control, and a need for heightened vigilance; c) The unpredictability of dementia led to challenges in preparing for the end of life; d) Grief is a staged-process and high levels of guilt and grief into bereavement is linked to the inability to effect change in care provision | 36 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|---|-----------|---|--|---|--|--|---------------|---------------|
| Robinson et al. (2010) [Canada] | Homebased | Sample: Family caregivers (n=29) Sampling method: Purposive Stage: Advanced | To understand how family caregivers of institutionalised relatives with dementia perceived and experienced relocation of their relative to a more home-like setting | Type: Mixed methods Design: Focus group | Data collection: Focus group and individual interviews Analysis: Thematic analysis | a) Caregivers often exhaust their physical, cognitive, emotional resources before agreeing to have their family member placed in care; b) The family caregiver's relational connection to the member with dementia remained strong and generated suffering when placement occurred that persisted over time and, for some, increased. This suffering took the form of grieving the losses arising from placement, guilt for being unable to prevent placement, second-guessing themselves with regards to the necessity of placement, and concern over the circumstances of the care. | 35 | A |
| Smith et al. (2001) [USA] | Homebased | Sample: Primary caregivers (family) (n=45) | To explore caregivers' narratives of their experience and needs, and | Type: Qualitative Design: Ethnography | Data collection: Semi- structured | a) Family caregivers had financial worries, with those identified as spouses expressing higher levels of anxiety; b) Some caregivers' | 35 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|-----------|--|---|---|--|--|---------------|---------------|
| | | <p>Sampling method: Purposive</p> <p>Stage: Mild to advanced</p> | how the disease and their interaction with multiple practitioners have impacted their lives | | <p>intensive interviews</p> <p>Analysis: Thematic analysis</p> | needs seemed more dependent upon unfulfilled expectations from family, support systems, or practitioners; c) Disappointment at the level of care, emotional support, or love expressed by family members | | |
| Tretteteig et al. (2017) [Norway] | Homebased | <p>Sample: Family caregivers (n=5)</p> <p>Sampling method: Purposive</p> <p>Stage: Moderate to advanced</p> | To gain knowledge about family caregivers' experiences of meaning in their caring role | <p>Type: Qualitative</p> <p>Design: Narrative (using Riessman's method)</p> | <p>Data collection: In-depth semi-structured interviews</p> <p>Analysis: Narrative analysis method using a case study approach</p> | a) Caregivers' roles and coping strategies were related to their relational ties; b) Difficulty in accepting their care recipient had changed; c) Caregivers had to make decisions whether to enhance, maintain or let go of emotional ties to find a good balance between meeting own needs and care recipient's needs; d) Letting go is almost impossible due to having a shared history with care recipient | 35 | A |

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|----------------------------------|-----------|---|---|---|---|--|---------------|---------------|
| Wladkowski (2016) [UK] | Homebased | <p>Sample: Family caregivers (n=24)</p> <p>Sampling method: <i>Purposive</i></p> <p>Stage: <i>Advanced</i></p> | To explore the grief experience in end-of-life care by caregivers | <p>Type: Qualitative</p> <p>Design: Qualitative</p> | <p>Data collection: Semi-structured interviews</p> <p>Analysis: Thematic analysis</p> | a) Unpreparedness for a change in terminal prognosis and uncertainty about being able to resume caregiving for their loved one following a live discharge; b) Emotional conflict in understanding disease progression following a live discharge, which also caused a struggle in understanding care recipient's terminal prognosis as being temporary; c) A level of ambiguity in the caregivers' grief process as caregivers become unsure of how or when to grieve. | 35 | A |

Synthesising the findings

Papers were grouped in the second stage according to the disease stage, country, authors, and population, and the common focus identified in the literature. A textual narrative synthesis of the common focus between the included studies was presented, which shows the interrelationship between the different studies in terms of their commonality on how caregiving experiences differ at both the moderate and advanced stages of dementia (See Supplementary Table 2). In the third stage, a preliminary synthesis is also presented, as a test for the robustness of the synthesised results and as an initial stage to result presentation. Popay et al. (2006) consider this a crucial interrogation of the initial synthesis required to understand the reasons for the findings presented by each study included (See Supplementary Table 3).

Textual narrative synthesis and presentation of findings

In congruence with Popay et al. (2006), the navigation of the textual narrative synthesis is sequential, showing the family carers' differentiated accounts of reported burden, loss, grief, how they coped with challenges and the motivation for carrying on with care provision within the explored stages of dementia. These findings are not presented in any hierarchy of importance. However, given its commonality of experience in dementia caregiving, the account of burden is presented first.

Burden: experience at moderate and advanced stages

Family carers in the moderate stage experienced increased caregiving responsibility due to increased care recipient's needs (Idura et al., 2018; Chan et al., 2010; Adams & Sanders, 2004). Extended time was spent caregiving as a result, which led to a clash with other social roles and a difficulty in meeting the family carers' own needs (Idura et al., 2018; Chan et al., 2010). In the case of spousal caregivers, extended time spent on caregiving was also highlighted (Idura et al., 2018; Chan et al., 2010; Adams & Sanders, 2004). Whilst the extended time caregiving was considered a necessity in some cases as highlighted by

Adams and Sanders (2004), the reason for this was not given. Swall et al. (2019) and Hellström et al. (2005) suggested, however, that this may be attributable to their couplehood.

Increased caregiving responsibility caused family carers' increased burden at the moderate stage of dementia (Adams & Sanders, 2004). Their difficulty in meeting own needs also intensified over time (Chan et al., 2010). This finding represented a high cost of caring with possible risk to family carers' mental and physical health (Idura et al., 2018; Brunton et al., 2008; Adams & Sanders, 2004). Furthermore, contrary to the report of an increase in caregiving being the only association to caregivers' increased burden, evidence also showed that family carers' increased burden was linked to progressive decline and changes in the care recipient's behaviour (Idura et al., 2018; Karger, 2018; Lindauer et al., 2016; Chan et al., 2010).

The collective account at the advanced stage further associated family carers' burden with care recipient's behavioural changes, which existed as part of progressive decline in dementia (Colling, 2004). Passivity, a distinctive change in care recipient's behaviour, which also affects functionality in thinking, movement, and interactions with others, was cited as an example (Colling, 2004). However, the severity of this change did not increase the family carers' burden (Idura et al., 2018; Chan et al., 2010). Passivity also changed the care recipient's emotional responses and caused the family carers' difficulty in accepting care recipients' changes (Karger, 2018). Overall, as the care recipient's passivity increased, the family carers' helplessness, sadness, and frustration at the inability to help also increased (Colling, 2004).

Passivity was, however, not only distinctive at the advanced stage. Common symptoms of increased agitation, aggression, and withdrawal were also reported at the moderate stage (Lindauer et al., 2016; Chan et al., 2010; Colling, 2004). As care recipients became more passive and required help with functional activities such as personal care, the demand for

their carer's attention also increased (Colling, 2004). This also caused the family carers' burden to increase in line with care recipients' increasing passivity at this stage of the disease.

Some family carers found it difficult to accept the process of care recipients' decline and expressed the most sadness at the advanced stage (Colling, 2004). Despite this, they accepted their role better, as they were adjusted to their new lifestyle (Chan et al., 2010; Colling, 2004). In comparison, less burden was reported by family carers at the moderate stage (Colling, 2004), which was attributed to their reliance on external resources such as extended family support (Idura et al., 2018; Chan et al., 2010; Brunton et al., 2008). While the availability of external support was commonly perceived as being able to reduce burden by family carers for recipients at both stages of dementia (Idura et al., 2018; Chan et al., 2010), evidence of consistency in burden reduction at either stage was not discussed in the reviewed literature.

Some family carers were assisted by their extended family (Brunton et al., 2008; Smith et al., 2001). Others experienced difficulty in asking for support (Chan et al., 2010; Brunton et al., 2008; Smith et al., 2001). Some family carers had to "*apply pressure*" to receive any assistance at all (Brunton et al., 2008), and disappointment was sometimes experienced at the level of assistance offered (Smith et al., 2001). Although some positive experiences were reported, such as the report of an improvement in interrelationships (Idura et al., 2018; Chan et al., 2010), tension within the family was a possibility (Chan et al., 2010). This also supports the finding that the reduction of family carer's burden through extended family support at both stages of dementia was not true in all cases.

Poor understanding of family carers' needs (Idura et al., 2018; Chan et al., 2010), was commonly evidenced as one of the factors for inadequate familial assistance at both stages (Chan et al., 2010; Brunton et al., 2008). Other reasons include living away from the care recipient (Chan et al., 2010; Brunton et al., 2008), and familial indifference (Chan et al.,

2010; Brunton et al., 2008). Receiving assistance from the family was therefore not always available. Moreover, it was possible for family expectations to attach a sense of duty to caregiving (Idura et al., 2018; Tretteteig et al., 2017; Chan et al., 2010). As such, caregiving therefore became an obligation at both stages (Idura et al., 2018; Chan et al., 2010).

Loss: experience at moderate and advanced stages

Family carers' loss was explored from two perspectives. The first, mostly descriptive of the narrative at the moderate stage, was related to their personal changes. This was described in relation to own time, freedom, independence, and being able to socialise (Idura et al., 2018; Chan et al., 2010; Brunton et al., 2008; Adams & Sanders, 2004). These lifestyle changes were stressful (Colling et al., 2004), and attributable to their increased caregiving responsibilities (Chan et al., 2010; Adams & Sanders, 2004; Colling, 2004). It was further reported by those who identified themselves as spouses, that this type of loss was the highest, especially in cases where caregiving ceases to be provided at home at the advanced stage of the disease (Adams & Sanders, 2004). Their loss was also accompanied by guilt (Adams & Sanders, 2004).

The second perspective was associated with the care recipient's changes, mostly at the advanced stage. This was described as the care recipient's altered behaviour (Colling, 2004), and the difference over time (Adams & Sanders, 2004; Colling, 2004). For this type of loss, care recipients were described in relation to who they once were (Lindauer et al., 2016; Adams & Sanders, 2004), in terms of their personality (Tretteteig et al., 2017; Lindauer et al., 2016; Adams & Sanders, 2004), and relationship once jointly shared with the family carer (Tretteteig et al., 2017; Lindauer et al., 2016; Brunton et al., 2008; Adams & Sanders, 2004).

Some family carers did not emphasise their care recipient's personality changes (Lindauer et al., 2016). For them, the care recipient's significance or value was instead considered of more relevance (Lindauer et al., 2016), than the changes attributed to what has been lost

to dementia (Lindauer et al., 2016). These family carers therefore held on to “*what remained*” of the care recipient for as long as possible (Lindauer et al., 2016, p.738). However, some family carers placed more emphasis on their care recipient’s lost personality (Adams & Sanders, 2004), instead of what remained (Lindauer et al., 2016). In comparison to those who considered the value of their care recipients to be of more significance, these carers regarded their care recipients as already gone (Adams & Sanders, 2004).

A commonality in the description of loss was therefore established as the recognition of the absence of the care recipient’s valued personality (Lindauer et al., 2016; Adams & Sanders, 2004). Both of these studies also gave evidence of a common experience of a paradoxical feeling, described as holding on to ‘what remains of the care recipient’ (Lindauer et al., 2016), or holding on to ‘someone who was already gone’ (Lindauer et al., 2016; Adams & Sanders, 2004). These two expressions appear synonymous. However, while evidence of historical hardship was cited between carers and care recipients in Lindauer et al. (2016), the existence of such a relationship was not described by Adams and Sanders (2004).

Grief: experience at moderate and advanced stages

Across the trajectory of dementia, variance in grief level existed (Lindauer et al., 2016; Adams & Sanders, 2004). As an emotional response to losing a loved one (Moore et al., 2017; Adams & Sanders, 2004; Colling, 2004), its severity was described as being at its highest at the advanced stage (Idura et al., 2018; Wladkowski, 2016; Adams & Sanders, 2004). In some cases, while family carers’ perception of grief was regarded as part of the pre-death experience (Moore et al., 2017; Lindauer et al., 2016; Adams & Sanders, 2004), it was also associated with the observation of the care recipient’s decline (Colling, 2004). Irrespective of the relationship between the family carer and their care recipient, the

association between higher grief level and higher loss reported at the advanced stage was unclear.

Some family carers reported a no-grief experience at both stages of dementia. At the moderate stage, for instance, this was considered a response to the delay in the care recipient's decline (Adams & Sanders, 2004). As a result, carers at this stage concentrated on their care recipient's lost personhood as an outlet for their perceived loss (Adams & Sanders, 2004). Although this claim was also supported by Lindauer et al. (2016), both studies differed in some of their findings. While Lindauer et al. (2016) asserted that carers who reported a no-grief experience focused on their care recipient's personality which remained, Adams and Sanders (2004) reported that family carers' emphasis was on their care recipient's already lost personality. The carers' no-grief experience was therefore related to anger, self-pity, or shame. (Adams & Sanders, 2004). On the other hand, the no-grief experience reported, especially by those with a shared history of hardship (Lindauer et al., 2016), establishes a defining characteristic of a shared identity.

Holding on to what remains caused some family carers' unpreparedness for the care recipient's death (Moore et al., 2017; Lindauer et al., 2016), as caregiver-attributed value was preserved longer. Indeed, for some, grief was irrelevant, and the reality of actual death was unacceptable (Lindauer et al., 2016). In this situation, a prolonged after-death grieving was therefore possible.

Dealing with challenges and continuing with caregiving

Caregiving was perceived as a natural obligation by some family carers (Lindauer et al., 2016), and they used religion to cope with the caregiving challenges encountered (Idura et al., 2018; Lindauer et al., 2016; Colling, 2004; Smith et al., 2001). Through religion, caregiving was perceived by some as rewarding or as the 'inheritance of good morals' (Idura et al., 2018), while others considered it as fulfilling an important role (Lindauer et al., 2016),

or doing the right thing (Tretteteig et al., 2017). These perceptions were common across both stages of the disease.

Caregiving expectations carried a sense of obligation, attributed by societal and cultural values (Idura et al., 2018; Chan et al., 2010), or the carer's self-induced expectations to provide care (Idura et al., 2018; Tretteteig et al., 2017). This, however, varied with relationship between the carer and the care recipient (Idura et al., 2018; Tretteteig et al., 2017; Chan et al., 2010). In some societies, for instance, the obligation to provide care to older people was placed on their family (Lindauer et al., 2016; Chan et al., 2010) by sources such as the society (Tretteteig et al., 2017), religion (Idura et al., 2018), culture (Chan et al., 2010), a sense of history, and identity shared with the care recipient (Lindauer et al., 2016). In these societies, the perceived sense of duty, therefore, influenced caregiving (Tretteteig et al., 2017).

The fear of inadequacy in knowledge and skills existed in some family carers' preparation for their increased caregiving responsibilities (Idura et al., 2018). At the moderate stage, this was managed by finding a balance between own expectations and the reality of the care recipient's behavioural changes (Idura et al., 2018). Although some carers used their prior knowledge of the care recipient's background for managing challenges faced in caregiving (Colling, 2004), others considered care recipient's placement in a care home (Lethin et al., 2016; Adams & Sanders, 2004; Smith et al., 2001), or receiving support at home (Lethin et al., 2016). For most carers, placing the care recipients away from home was however inconceivable, especially at the advanced stage.

A preference for continuing with home-based caregiving was highlighted, especially in situations where the care recipient's decline necessitates consideration for care provision outside of the home setting (Lindauer et al., 2016). Some family carers found this decision difficult, and the wish to maintain control on caregiving persisted (De Silva & Curzio, 2009). This wish to maintain control increased their ability to make meaning of the caregiving role

(Tretteteig et al., 2017). Although this was considered a characteristic of carer's grief (Adams & Sanders, 2004), maintaining control was also crucial for balancing the carer-to-recipient relationship.

Loss of control was accompanied by a feeling of guilt and sadness, especially where the care recipient's placement away from home was necessary (Moore et al., 2017; Tretteteig et al., 2017; De Silva & Curzio, 2009; Adams & Sanders, 2004; Chan et al., 2010). The fear of poor caregiving by others outside of the home setting was cited as a reason (Moore et al., 2017). Situations occurred whereby some of the family carers' guilt and sadness persisted beyond bereavement, due to self-blame (Moore et al., 2017). Although evidence was given which showed that this further increased the reluctance to seek outside care away from home, the influence of family carers' perceived obligation to continue caregiving, and their need to maintain control in such cases was unclear.

Moving the person cared for away from home was difficult (Lindauer et al., 2016). Some family carers used respite as their preferable means for managing their caregiving-related challenges (Idura et al., 2016; Lethin et al., 2016). Generally, respite is used in briefly taking away caregiving responsibilities (de La Cuesta-Benjumea, 2011). Variation in its description, however, exists within the reviewed literature. According to Brunton et al. (2008), respite is described as a physical location, where care recipients reside in order for their carers to have a break, or a situation where caregiving responsibility is shifted to another person, such as a family member (Idura et al., 2018; Brunton et al., 2008). However, de La Cuesta-Benjumea (2011) described respite as a situation whereby family carers may separate themselves from their caregiving identity and briefly assume a different identity (de La Cuesta-Benjumea, 2011). This therefore suggests that achieving rest is possible without the need for family carers' physical separation from their care recipients.

Alternative identities were achievable through mental isolation, whereby caregiving responsibilities were mentally paused by the family carer to give them an opportunity to

explore other identities (de La Cuesta-Benjumea, 2011). As such, managing caregiving challenges was possible, regardless of the carer's physical location (de La Cuesta-Benjumea, 2011). This may be useful in situations where a respite facility is unavailable or where there is a reluctance to place the care recipient away from home.

Discussion

The central focus of experiences was mainly negative, and broadly on family carers' accounts of loss, burden, and grief. These experiences are reported in the wider literature as attributes commonly associated with general experiences of dementia caregiving. Some positive aspects of caregiving were also reported, including improvement in family interrelationships, and the family carers' feeling of being useful and having a sense of meaning. An association between loss, burden, and grief of the caregiver exists, according to Meuser and Marwit (2001). The experience of these was not descriptive of all family carers' narratives in this review. Furthermore, these are not the only dementia-caregiving related experiences, as a variety of other experiences, such as stress, anxiety, and depression are also reported in the wider literature (Blandin & Pepin, 2017; Chan et al., 2010; Adams & Sanders, 2004). The order of these experiences by the family carers, however, remains unclear.

Narratives of loss were explored from two different perspectives in this review. Firstly, it was described in relation to family carers' unmet needs, as a loss of social roles and difficulty in fulfilling own needs. This was termed '*personified loss*', as descriptions represented the family carers' perception of own loss. The description of loss in terms of family carers' own needs has been given in other studies. Brodaty and Donkin (2009) describe this in terms of social isolation which may result in a loss of meaning in life. Similar findings have also been reported in this current review, and it is suggested that this experience may worsen over time due to the possibility of emotional burden exposure from increased caregiving (Greenwood et al., 2019; Brodaty & Hadzi-Pavlovic, 1990).

Changes to the way of life were described by the family carers' increased burden through caregiving responsibilities. Inability to socialise at this stage of dementia was viewed as a conflict between increased psychosocial distress and associated changes to caregiving focus. In relevance to this experience, the description of family carers' *personified loss* may therefore be a precursor to a *personified cost* in terms of opportunities forfeited for accepting caregiving responsibilities. Hence, the likelihood of family carers' *personified loss* being interpreted as a forfeited way of life rather than expressed as part of the normal progression of dementia caregiving is elevated.

Secondly, loss was also described as a caregiver-attributed value in relation to the care recipient's changes and termed '*personified value*' in this review. Dupuis (2002) identifies this as a detrimental attribute due to the possible closeness between the carer and the person cared for. This type of loss was further described in terms of ambiguity given that it acknowledges the absence of a valued personality (Dupuis, 2002; Boss, 1999; Jones & Martinson, 1992). In similarity to the findings by Dupuis (2002), evidence by Boss (1999), and Jones and Martinson (1992) suggest that family carers experience a state of loss where the person cared for was considered gone while still physically present. Similarity of findings in this review shows that although loss suggests a *personified value*, the view of the person cared for as 'already gone' while still alive was also shared.

The possibility of increasing family carers' emotional hold to the personalities that remain is considerable, given the fear of losing *personified value*. Subsequently, holding on persists longer as a paradox where balancing "*what remains*" of the care recipient with "*what is already lost*" becomes difficult. As in other terminal conditions, evidence suggests that this experience, also referred to as anticipatory grief, is a pre-death response to the process of progressive decline in dementia (Pérez-González et al., 2021; Cheung et al., 2018; Blandin & Pepin, 2017; Meuser & Marwit, 2001). Thus, in cases where the realisation of dementia contributes to the early attribution of *personified value*, family carers' fear of losing *what remains* may cause a higher level of grief in both stages of dementia.

The resolution of grief corresponds with family carers' acceptance of own loss (Prigerson & Maciejewski, 2008). As an “*emotional inability to accept the loss of something cherished*”, non-acceptance results in grief and attributes such as guilt, frustration, and helplessness (Prigerson & Maciejewski, 2008, p.435). While grief may be described as a natural response to family carers' loss, those who report higher levels of pre-death grief have a higher risk of health complications post-death (Shuter et al., 2014; Chan et al., 2010; Givens et al., 2011; Rando, 2000). The risk of health complications from earlier stages of the disease and possibly post-death may therefore increase. An early resolution of caregiver-attributed *personified value* is therefore necessary.

Grief was, however, not an experience reported by family carers who shared a historical hardship with the care recipient in this review. In commonality with other caregivers, holding on to *personified value* for longer was highlighted. While it is appropriately assumed that “*something cherished*” (Prigerson & Maciejewski, 2008, p.435) may refer to factors such as family carers' own time, freedom, or care recipient's value, as well as the identity shared as a result, the loss described may also be personified. Further research is, however, required for differentiating between a caregiver-attributed value of what remains and what is already lost to dementia, and its influence on the possible duration of prolonged after-death grieving.

It is further appropriate to also consider that, for most family carers, a grief-experience is reported through both stages of dementia irrespective of their relationship with the care recipient (Meuser & Marwit, 2001). As the care recipient declines, *personified loss* increases in line with the increased demand for the carers' attention. Consequently, loss of *personified value* and family carers' grief may also increase at both stages. The justification and determinants for balancing *personified value* against family carers' *personified loss* are however, currently unclear. Given these findings, the issue of balancing family carers' reported experiences against their reasons for continuing caregiving at both stages of the disease therefore arises. It is assumed that other factors may also be present in the family carers' experiences, whereby powerlessness is felt in how future priorities are perceived and

how the balance between *personified value* and *personified loss* may influence continued caregiving. This also calls for an understanding of a possible distress at choosing the right course of action to take in such situations (Oh & Gastmans, 2015; Silén, 2011; Jameton, 1993; 1984).

Some gaps in current knowledge were also identified during this review. It was claimed that variation in family carers' grief-experience across dementia stages increases with severity (Idura et al., 2018; Lindauer et al., 2016; Wladkowski, 2016; Adams & Sanders, 2004). Moreover, a higher level of grief-experience is associated with the advanced stages (Adams & Sanders, 2004). Clarity is however required to determine whether a higher grief level at the advanced stage is resultant from a higher loss experienced from earlier stages.

Placement of caregiving expectations from different sources was suggested as a reason for an induced sense of obligation on family carers (Idura et al., 2018; Tretteteig et al., 2017; Chan et al., 2010). This varied according to the relationship between the family carers and their care recipients (Idura et al., 2018; Tretteteig et al., 2017; Chan et al., 2010), and serves as a factor which influenced behaviour (Tretteteig et al., 2017). While the possibility of negative outcomes was reported, the relationship between the family carers' obligation and grief level remained unexplored. Although similarity of experience of holding on to care recipients' personality was evidenced in the literature, an ambiguity remained between clear definition of what remains of the care recipient (Lindauer et al., 2016) and what was already lost to dementia (Adams & Sanders, 2004), as well as its influence on determining the duration of prolonged after-death grieving.

Implications

The impact of higher levels of pre-death grief in caregiving for someone with dementia is widely recognised in association with health complications post-death (Shuter et al., 2014; Chan et al., 2010; Givens et al., 2011). Although higher grief level is commonly attributed to the advanced stages, the possibility of attributing a *personified value* earlier in caregiving is

evident from this review. For the family carer, the fear of losing care recipients' *personified value* carries a higher level of grief in both stages, thereby increasing the risk of health complications earlier and possibly post-death. This may influence how home-based caregiving is viewed by family carers and their willingness to accept or continue caregiving responsibilities at home. Given a projected increase to over two million dementia family carers by 2051 (Knapp et al., 2014), the implication is therefore severe for sustaining home-based caregiving for people with moderate to advanced dementia in the UK.

Strengths and limitations of the review

The adoption of a narrative synthesis approach allowed for heterogeneity of evidence in this review. This approach may have contributed to the robustness of synthesised evidence. A difficulty in managing the scope of possible evidence was however high, as a large number of individual papers were checked for the inclusion criteria to be met. Many papers were found where the population of interest was either missing or incorrectly reported. This difficulty was further exacerbated by the inaccurate classification of some UK papers by their authors, thereby making both the screening stage and inclusion problematic from reading their abstracts. Variation of accounts contributes to how family carers' experience of loss may be explored in future research. Given that some accounts used were extrapolated from a wider context of discussion within the reviewed literature, full justification of accounts may therefore be difficult (Polit & Beck, 2010). Generalisation of the findings at either of the moderate and advanced stages of dementia may also be difficult. However, the main purpose of any qualitative endeavour, such as this review, is to provide a rich understanding of people's accounts. Polit and Beck (2010) highlight that the transferability of these accounts across clearly defined settings and contexts is therefore appropriate.

Conclusions and recommendations

Family carers' descriptions of experiences at the moderate to advanced stages of dementia vary. Overall, accounts suggest that spending time with the person cared for enhances the acknowledgement of the illness experience by the family carer. Using a narrative synthesis approach, the findings of this review suggest that the level of understanding of the illness experience differs. A complete description of all experiences may therefore be inadequate in conveying an acknowledgement of the illness within home-based caregiving. Given accounts of burden, loss, grief, and dealing with challenges in making meaning show that increasing caregiving responsibilities increased the family carer's difficulties, especially at the moderate stage. Understanding the experience of loss through to the advanced stages is also crucial. It is suggested that the caregiver-attributed *personified value* resulted in holding on to what remains of the care recipient for longer. Research is however required to understand whether this may lead to a higher grief experience from the moderate stage or earlier. Further research is also required to explore how family carers may proportionally balance their *personified loss* with their *personified value* earlier in the disease trajectory, given that the justification and determinants for balancing these, and whether distress to morals is felt, remains unclear.

Acknowledgement

We would like to express our gratitude to Professor Katherine Froggatt for her advice.

Funding information

This review received no grant from any funding body.

Conflicts of interest

The authors declare that there are no conflicts of interests.

Ethics statement

Ethical approval was not required for this review.

Data availability

Data supporting this review are included within the paper.

References

- Adams, K. B., & Sanders, S. (2004). Alzheimer's caregiver differences in experience of loss, grief reactions and depressive symptoms across stage of disease: a mixed-method analysis. *Dementia*, 3(2), 195-210. Retrieved 2nd November 2020, from: <https://doi.org/10.1177/1471301204042337>
- Al-Janabi, H., Carmichael, F., & Oyebode, J. (2018). Informal care: choice or constraint? *Scandinavian Journal of Caring Sciences*, 32(1), 157-167. Retrieved 27th October 2020, from: <https://doi.org/10.1111/scs.12441>
- Alzheimer's Society. (2015). The progression of Alzheimer's disease and other dementias. Retrieved 21st January 2021, from: https://www.alzheimers.org.uk/sites/default/files/pdf/factsheet_the_progression_of_alzheimers_disease_and_other_dementias.pdf
- Aminzadeh, F., Byszewski, A., Molnar, F. J., & Eisner, M. (2007). Emotional impact of dementia diagnosis: Exploring persons with dementia and caregivers' perspectives. *Aging & Mental Health*, 11(3), 281-290. Retrieved 15th March 2020, from: <https://doi.org/10.1080/13607860600963695>
- Arcand, M. (2015). End-of-life issues in advanced dementia: Part 1: goals of care, decision-making process, and family education. *Canadian Family Physician Medecin De Famille Canadien*, 61(4), 330-334. Retrieved 10th January 2021, from: <https://pubmed.ncbi.nlm.nih.gov/25873700>
- Auer, S., & Reisberg, B. (1997). The GDS/FAST staging system. *International Psychogeriatrics*, 9(S1), 167-171. Retrieved 28th February 2021, from: <https://doi.org/10.1017/S1041610297004869>
- Bieber, A., Nguyen, N., Meyer, G., & Stephan, A. (2019). Influences on the access to and use of formal community care by people with dementia and their informal caregivers: a scoping review. *BMC Health Services Research*, 19(1), 88. Retrieved 27th January 2021, from: <https://doi.org/10.1186/s12913-018-3825-z>
- Blandin, K., & Pepin, R. (2017). Dementia grief: A theoretical model of a unique grief experience. *Dementia (London)*, 16(1), 67-78. Retrieved 27th January 2021, from: <https://doi.org/10.1177/1471301215581081>
- Boss, P. (1999). *Ambiguous loss: Learning to live with unresolved grief*. Harvard University Press.
- Brody, H., & Donkin, M. (2009). Family caregivers of people with dementia. *Dialogues in Clinical Neuroscience*, 11(2), 217-228. Retrieved 20th March 2020, from: <https://pubmed.ncbi.nlm.nih.gov/19585957/>
- Brody, H., & Hadzi-Pavlovic, D. (1990). Psychosocial effects on carers of living with persons with dementia. *Australian & New Zealand Journal of Psychiatry*, 24(3), 351-361. Retrieved 20th March 2020, from: <https://doi.org/10.3109/00048679009077702>
- Brunton, M., Jordan, C., & Fouche, C. (2008). Managing public health care policy: who's being forgotten? *Health Policy*, 88(2-3), 348-358. Retrieved 27th January 2021, from: <https://doi.org/10.1016/j.healthpol.2008.04.007>

- Chan, W. C., Ng, C., Mok, C.C. M., Wong, F. L. F., Pang, S. L., & Chiu, H. F. K. (2010). Lived experience of caregivers of persons with dementia in Hong Kong: a qualitative study. *East Asian Archives of Psychiatry*, 20(4), 163-168. Retrieved 12th March 2020, from: <https://search.informit.org/doi/10.3316/INFORMIT.934619447079403>
- Cheung, D.S.K., Ho, K.H.M., Cheung, T.F., Lam, S.C., Tse, M.M.Y. (2018). Anticipatory grief of spousal and adult children caregivers of people with dementia. *BMC Palliat Care* 17, 124. Retrieved 8th January 2022, from: <https://doi.org/10.1186/s12904-018-0376-3>
- Clemmensen, T. H., Busted, L. M., Søbørg, J., & Bruun, P. (2016). The family's experience and perception of phases and roles in the progression of dementia: An explorative, interview-based study. *Dementia*, 18(2), 490-513. Retrieved 20th March 2020, from: <https://doi.org/10.1177/1471301216682602>
- Colling, K. B. (2004). Special section - behavioral symptoms of dementia: their measurement and intervention. Caregiver interventions for passive behaviors in dementia: links to the NDB model. *Aging & Mental Health*, 8(2), 117-125. Retrieved 20th March 2020, from: <https://doi.org/10.1080/13607860410001649626>
- de la Cuesta-Benjumea, C. (2011). Strategies for the relief of burden in advanced dementia care-giving. *Journal of Advanced Nursing*, 67(8), 1790-1799. Retrieved 20th March 2020, from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2648.2010.05607.x>
- Dementia Carers Count. (2019). Dementia care statistics: Key facts & figures. Retrieved 22nd February 2021, from: <https://dementiacarers.org.uk/for-dementia-professionals/key-facts-figures/>
- De Silva, N., & Curzio, J. (2009). Transition into long-term care: the lived experience of carers of individuals with dementia. *Mental Health Nursing*, 29(5), 10-14. Retrieved 25th March 2020, from: <https://www.proquest.com/openview/63367e292b9e8c5f279df23c1dca883f/1?pq-origsite=gscholar&cbl=135348>
- Dupuis, S. L. (2002). Understanding ambiguous loss in the context of dementia care. *Journal of Gerontological Social Work*, 37(2), 93-115. Retrieved 18th March 2020, from: https://doi.org/10.1300/J083v37n02_08
- Evans, S., Waller, S., Bray, J., & Atkinson, T. (2019). Making homes more dementia-friendly through the use of aids and adaptations. *Healthcare (Basel, Switzerland)*, 7(1), 43. Retrieved 17th March 2020, from: <https://doi.org/10.3390/healthcare7010043>
- Farina, N., Page, T. E., Daley, S., Brown, A., Bowling, A., Basset, T., Livingston, G., Knapp, M., Murray, J., & Banerjee, S. (2017). Factors associated with the quality of life of family carers of people with dementia: A systematic review. *Alzheimer's & Dementia*, 13(5), 572-581. Retrieved 17th March 2020, from: <https://doi.org/https://doi.org/10.1016/j.jalz.2016.12.010>

- Galvin, J. E., & Sadowsky, C. H. (2012). Practical guidelines for the recognition and diagnosis of dementia. *The Journal of the American Board of Family Medicine*, 25(3), 367. Retrieved 10th January 2021, from: <https://doi.org/10.3122/jabfm.2012.03.100181>
- Givens, J. L., Prigerson, H. G., Kiely, D. K., Shaffer, M. L., & Mitchell, S. L. (2011). Grief among family members of nursing home residents with advanced dementia. *American Journal of Geriatric Psychiatry*, 19(6), 543-550. Retrieved 10th January 2020, from: <https://doi.org/10.1097/JGP.0b013e31820dcbe0>
- Greenwood, N., Pound, C., Smith, R., & Brearley, S. (2019). Experiences and support needs of older carers: A focusgroup study of perceptions from the voluntary and statutory sectors. *Maturitas*, 123, 40-44. Retrieved 10th January 2021, from: <https://doi.org/10.1016/j.maturitas.2019.02.003>
- Greenwood, N., & Smith, R. (2019). Motivations for being informal carers of people living with dementia: a systematic review of qualitative literature. *BMC Geriatrics*, 19(1), 169. Retrieved 10th January 2021, from: <https://doi.org/10.1186/s12877-019-1185-0>
- Hall, M., & Sikes, P. (2018). From "What the Hell Is Going on?" to the "Mushy Middle Ground" to "Getting Used to a New Normal": Young people's biographical narratives around navigating parental dementia. *Illness, Crises & Loss*, 26(2), 124-144. Retrieved 10th October 2020, from: <https://doi.org/10.1177/1054137316651384>
- Harris, D. (2007). Forget me not: palliative care for people with dementia. *Postgraduate Medical Journal*, 83(980), 362-366. Retrieved 10th January 2021, from: <https://doi.org/10.1136/pgmj.2006.052936>
- Hawker, S., Payne, S., Kerr, C., Hardey, M., & Powell, J. (2002). Appraising the evidence: Reviewing disparate data systematically. *Qualitative Health Research*, 12(9), 1284-1299. Retrieved 10th October 2020, from: <https://doi.org/10.1177/1049732302238251#>
- Hellström, I., Nolan, M., & Lundh, U. (2005). 'We do things together': A case study of 'couplehood' in dementia. *Dementia*, 4(1), 7-22. Retrieved 10th January 2020, from: <https://doi.org/10.1177/1471301205049188>
- Idura, A. M., Noorlaili, M. T., Rosdinom, R., Azlin, B., & Iryani, M. D. (2018). Caring for moderate to severe dementia patients - Malaysian family caregivers experience. *International Medical Journal Malaysia*, 17(1), 101-110.
- James, C.O. (2021). *Moral Distress in the Care of People Living with Moderate to Advanced Dementia: A Narrative Exploration of Family Carers' Experience of Home-Based Care Provision towards the End of Life*. PhD Thesis, Lancaster University.
- James, C., Walshe, C., & Froggatt, K. (2020). Protocol for a systematic review on the experience of informal caregivers for people with a moderate to advanced dementia within a domestic home setting. *Systematic Reviews*, 9(1), 270. Retrieved 28th February 2021, from: <https://doi.org/10.1186/s13643-020-01525-0>
- Jameton, A. (1984). *Nursing practice: The ethical issues*. Prentice-Hall.

- Jameton, A. (1993). Dilemmas of moral distress: moral responsibility and nursing practice. *AWHONNS Clinical Issues in Perinatal and Women's Health Nursing*, 4(4), 542-551. Retrieved 28th February 2019, from: <https://pubmed.ncbi.nlm.nih.gov/8220368/>
- Joling, K. J., Janssen, O., Francke, A. L., Verheij, R. A., Lissenberg-Witte, B. I., Visser, P. J., & van Hout, H. P. (2020). Time from diagnosis to institutionalization and death in people with dementia. *Alzheimer's & Dementia*, 16(4), 662-671. Retrieved 28th January 2021, from: <https://doi.org/10.1002/alz.12063>
- Jones, P. S., & Martinson, I. M. (1992). The experience of bereavement in caregivers of family members with Alzheimer's disease. *Image Journal of Nursing Scholarship*, 24(3), 172-176. Retrieved 28th January 2021, from: <https://doi.org/10.1111/j.1547-5069.1992.tb00714.x>
- Karg, N., Graessel, E., Randzio, O., & Pendergrass, A. (2018). Dementia as a predictor of care-related quality of life in informal caregivers: a cross-sectional study to investigate differences in health-related outcomes between dementia and non-dementia caregivers. *BMC Geriatrics*, 18(1), 189. Retrieved 28th January 2021, from: <https://doi.org/10.1186/s12877-018-0885-1>
- Karger, C. R. (2018). Emotional experience in patients with advanced Alzheimer's disease from the perspective of families, professional caregivers, physicians, and scientists. *Aging & Mental Health*, 22(3), 316-322.
- Knapp, M., Prince, M., Guerchet, M., McCrone, P., Prina, M., Comas-Herrera, A., Wittenberg, R., Adelaja, B., Hu, B., King, D., Rehill, A., & Salimkumar, D. (2014). *Dementia UK: Update (second edition)*.
- Kumar, A. S., J.; Goyal, A.; Tsao, J.W. (2020). *Alzheimer disease*. Stat Pearls Publishing; 2020. Retrieved 8th January 2021, from: <https://www.ncbi.nlm.nih.gov/books/NBK499922/>
- Kwon, Y., Lemieux, M., McTavish, J., & Wathen, N. (2015). Identifying and removing duplicate records from systematic review searches. *Journal of the Medical Library Association: JMLA*, 103(4), 184-188. Retrieved 15th January 2021, from: <https://doi.org/10.3163/1536-5050.103.4.004>
- Lee, L., & Weston, W. W. (2011). Disclosing a diagnosis of dementia: helping learners to break bad news. *Canadian Family Physician Medecin De Famille Canadien*, 57(7), 851-e272. Retrieved 25th January 2020, from: <https://pubmed.ncbi.nlm.nih.gov/21753111>
- Lethin, C., Hallberg, I. R., Karlsson, S., & Janlöv, A. (2016). Family caregivers experiences of formal care when caring for persons with dementia through the process of the disease. *Scandinavian Journal of Caring Sciences*, 30(3), 526-534. Retrieved 27th March 2021, from: <https://doi.org/10.1111/scs.12275>
- Lewis, F., Sussex, J., Schaffer, S. K., O'Neill, P., & Cockcroft, L. (2014). The trajectory of dementia in the UK - Making a difference. IDEAS Working Paper Series from RePEc. Retrieved 15th February 2020, from: https://www.ohe.org/system/files/private/publications/401%20-%20Trajectory_dementia_UK_2014.pdf

- Lewis, L. F. (2014). Caregivers' experiences seeking hospice care for loved ones with dementia. *Qualitative Health Research*, 24(9), 1221-1231. Retrieved 15th February 2020, from: <https://doi.org/10.1177/1049732314545888>
- Lindauer, A., Harvath, T. A., Berry, P. H., & Wros, P. (2016). The meanings African American caregivers ascribe to dementia-related changes: The paradox of hanging on to loss. *Gerontologist*, 56(4), 733-742. Retrieved 15th February 2020, from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6282689/pdf/gnv023.pdf>
- Lindeza, P., Rodrigues, M., Costa, J., Guerreiro, M., & Rosa, M. M. (2020). Impact of dementia on informal care: a systematic review of family caregivers' perceptions. *BMJ Supportive & Palliative Care*, bmjpspcare-2020-002242. Retrieved 15th February 2021, from: <https://doi.org/10.1136/bmjpspcare-2020-002242>
- Lorenc, T., Petticrew, M., Whitehead, M., Neary, D., Clayton, S., Wright, K., Thomson, H., Cummins, S., Sowden, A., & Renton, A. (2014). Appendix 5: Quality assessment for the systematic review of qualitative evidence. In *Crime, fear of crime and mental health: synthesis of theory and systematic reviews of interventions and qualitative evidence*. NIHR Journals Library. Retrieved 25th January 2020, from: <https://doi.org/10.3310/phr02020>
- Lorenzetti, D. L., & Ghali, W. A. (2013). Reference management software for systematic reviews and meta-analyses: an exploration of usage and usability. *BMC Med Res Methodol*, 13, 141. Retrieved 25th January 2020, from: <https://doi.org/10.1186/1471-2288-13-141>
- Marshall, G. A., Amariglio, R. E., Sperling, R. A., & Rentz, D. M. (2012). Activities of daily living: where do they fit in the diagnosis of Alzheimer's disease? *Neurodegenerative Disease Management*, 2(5), 483-491. Retrieved 25th August 2020, from: <https://doi.org/10.2217/nmt.12.55>
- Mesterton, J., Wimo, A., By, A., Langworth, S., Winblad, B., & Jönsson, L. (2010). Cross sectional observational study on the societal costs of Alzheimer's disease. *Current Alzheimer Research*, 7(4), 358-367. Retrieved 23rd January 2021, from: <https://doi.org/10.2174/156720510791162430>
- Meuser, T. M., & Marwit, S. J. (2001). A comprehensive, stage-sensitive model of grief in dementia caregiving. *The Gerontologist*, 41(5), 658-670. Retrieved 13th June 2020, from: <https://doi.org/10.1093/geront/41.5.658>
- Mlinac, M. E., & Feng, M. C. (2016). Assessment of activities of daily living, self-care, and independence. *Archives of Clinical Neuropsychology*, 31(6), 506-516. Retrieved 17th January 2021, from: <https://doi.org/10.1093/arclin/acw049>
- Moore, K. J., Davis, S., Gola, A., Harrington, J., Kupeli, N., Vickerstaff, V., King, M., Leavey, G., Nazareth, I., Jones, L., & Sampson, E. L. (2017). Experiences of end of life amongst family carers of people with advanced dementia: longitudinal cohort study with mixed methods. *BMC Geriatrics*, 17(135).
- Oh, Y., & Gastmans, C. (2015). Moral distress experienced by nurses: A quantitative literature review. 22(1), 15-31. Retrieved 18th October 2020, from: <https://doi.org/10.1177/0969733013502803>
- Page, M. J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., et al. (2020). The PRISMA 2020 statement: an updated guideline for reporting systematic

- reviews. *BMJ* 2021; 372. Retrieved 15th July 2021, from: <https://doi.org/10.1136/bmj.n160>
- Pérez-González, A., Vilajoana-Celaya, J, Guàrdia-Olmos, J. (2021). Alzheimer's Disease Caregiver Characteristics and Their Relationship with Anticipatory Grief. *International Journal of Environmental Research and Public Health*. 18(16):8838. Retrieved 8th January 2022, from: <https://doi.org/10.3390/ijerph18168838>
- Pertl, M. M., Sooknarine-Rajpatty, A., Brennan, S., Robertson, I. H., & Lawlor, B. A. (2019). Caregiver choice and caregiver outcomes: A longitudinal study of Irish spousal dementia caregivers. *Frontiers in Psychology*, 10(1801). Retrieved 18th October 2020, from: <https://doi.org/10.3389/fpsyg.2019.01801>
- Phillips, J., Pond, D., & Goode, S. M. (2011). Timely diagnosis of dementia: Can we do better? A report for Alzheimer's Australia. *Alzheimer's Australia*. Paper 24. Retrieved 18th October 2020, from: https://www.dementia.org.au/sites/default/files/Timely_Diagnosis_Can_we_do_better.pdf
- Polit, D. F., & Beck, C. T. (2010). Generalization in quantitative and qualitative research: Myths and strategies. *International Journal of Nursing Studies*, 47(11), 1451-1458. Retrieved 18th October 2020, from: <https://doi.org/10.1016/j.ijnurstu.2010.06.004>
- Popay, J., Roberts, H., Sowden, A., Petticrew, M., Arai, L., Rodgers, M., Britten, N., Roen, K., & Duffy, S. (2006). Guidance on the conduct of narrative synthesis in systematic Reviews. Retrieved 18th March 2020, from: <https://www.lancaster.ac.uk/media/lancaster-university/content-assets/documents/fhm/dhr/chir/NSsynthesisguidanceVersion1-April2006.pdf>
- Prigerson, H. G., & Maciejewski, P. K. (2008). Grief and acceptance as opposite sides of the same coin: setting a research agenda to study peaceful acceptance of loss. *British Journal of Psychiatry*, 193(6), 435-437. Retrieved 18th March 2020, from: <https://doi.org/10.1192/bjp.bp.108.053157>
- Rando, T. A. (2000). *Clinical dimensions of anticipatory mourning: Theory and practice in working with the dying, their loved ones, and their caregivers*. Research Press.
- Reinhard, S., Levine, C., & Samis, S. (2012). Home alone: Family caregivers providing complex chronic care. Retrieved 18th January 2021, from: https://www.aarp.org/content/dam/aarp/research/public_policy_institute/health/home-alone-family-caregivers-providing-complex-chronic-care-rev-AARP-ppi-health.pdf
- Sanders, A. E. (2016). Caregiver stress and the patient with dementia. *Continuum: Lifelong Learning in Neurology*, 22(2), 619-625. Retrieved 27th March 2020, from: <https://doi.org/10.1212/con.0000000000000301>
- Schmidt, K. (2014). Clinical dementia rating scale. In A. C. Michalos (Ed.), *Encyclopedia of Quality of Life and Well-Being Research* (pp. 957-960). Springer Netherlands. Retrieved 20th January 2021, from: https://doi.org/10.1007978-94-007-0753-5_690
- Schulz, R., Beach, S. R., Cook, T. B., Martire, L. M., Tomlinson, J. M., & Monin, J. K. (2012). Predictors and consequences of perceived lack of choice in becoming an informal caregiver. *Aging & Mental Health*, 16(6), 712-721. Retrieved 27th January 2021, from: <https://doi.org/10.1080/13607863.2011.651439>

- Schulz, R., & Eden, J. (2016). Family caregiving roles and impacts. In R. Schulz & J. Eden (Eds.), *Families Caring for an Aging America*. National Academies Press (US). Retrieved 27th January 2021, from: <https://doi.org/10.17226/23606>
- Shuter, P., Beattie, E., & Edwards, H. (2014). An exploratory study of grief and health-related quality of life for caregivers of people with dementia. 29(4), 379-385. Retrieved 21st July 2020, from: <https://doi.org/10.1177/1533317513517034>
- Silén, M. (2011). Encountering ethical problems and moral distress as a nurse (Publication Number 20), Jönköping University.
- Smith, A. L., Lauret, R., Peery, A., & Mueller, T. (2001). Caregiver needs: a qualitative exploration. *Clinical Gerontologist*, 24(1), 3-26. Retrieved 20th March 2020, from: https://doi.org/10.1300/J018v24n01_02
- Swall, A, Williams, C, Marmstål Hammar, L. (2020). The value of “us”—Expressions of togetherness in couples where one spouse has dementia. *International Journal of Older People Nursing*. 15:e12299. Retrieved 4th April 2021, from: <https://doi.org/10.1111/opn.12299>
- Tretteteig, S., Vatne, S., & Rokstad, A. M. (2017). The influence of day care centres designed for people with dementia on family caregivers - a qualitative study. *BMC Geriatrics*, 17(1), 5. Retrieved 20th March 2020, from: <http://dx.doi.org/10.1186/s12877-016-0403-2>
- Tretteteig, S., Vatne, S., & Rokstad, A. M. (2017). Meaning in family caregiving for people with dementia: a narrative study about relationships, values, and motivation, and how day care influences these factors. *Journal of Multidisciplinary Healthcare*, 10, 445-455. Retrieved 20th March 2020, from: <https://doi.org/10.2147/JMDH.S151507>
- van Gennip, I.E., Roeline, H., Pasman, W., Oosterveld-Vlug, M.G., Willems, D.L., Onwuteaka-Philipsen, B.D. (2016). How Dementia Affects Personal Dignity: A Qualitative Study on the Perspective of Individuals with Mild to Moderate Dementia. *The Journals of Gerontology: Series B*, 71(3), 491–501, Retrieved 4th January 2022, from: <https://doi.org/10.1093/geronb/gbu137>
- van Wijngaarden, E., van der Wedden, H., Henning, Z., Komen, R., & The, A. (2018). Entangled in uncertainty: The experience of living with dementia from the perspective of family caregivers. *PLOS ONE*, 13(6), e0198034-e0198034. Retrieved 12th February 2021, from: <https://doi.org/10.1371/journal.pone.0198034>
- Vestergaard, A.H.S., Sampson, E.L., Johnsen, S.P., Petersen, I. (2020). Social Inequalities in Life Expectancy and Mortality in People with Dementia in the United Kingdom. *Alzheimer Disease & Associated Disorders*. 34(3), 254-261. Retrieved 4th January 2022, from: <https://doi.org/10.1097/WAD.0000000000000378>
- Wladkowski, S. P. (2016). Live discharge from hospice and the grief experience of dementia caregivers. *Journal of Social Work*. In: *End-of-Life & Palliative Care*, 12(1), 47-62. Retrieved 5th March 2020, from: <https://doi.org/10.1080/15524256.2016.1156600>
- Woodman, C., Baillie, J., & Sivell, S. (2016). The preferences and perspectives of family caregivers towards place of care for their relatives at the end-of-life. A systematic review and thematic synthesis of the qualitative evidence. *BMJ*

Supportive & Palliative Care, 6(4), 418. Retrieved 27th May 2020, from:
<https://doi.org/10.1136/bmjspcare-2014-000794>

Xanthopoulou, P., McCabe, R. (2019). Subjective experiences of cognitive decline and receiving a diagnosis of dementia: qualitative interviews with people recently diagnosed in memory clinics in the UK. *BMJ Open* 2019;**9**: e026071.
Retrieved 4th January 2022, from: <https://doi.org/10.1136/bmjopen-2018-026071>




Fig 1: Sources of studies included (PRISMA diagram adapted from Page et al., 2020).

Table 1: Review inclusion and exclusion criteria

| Inclusion Criteria | Exclusion Criteria |
|--|---|
| Type of papers: peer-reviewed empirical papers, qualitative papers including case studies and mixed-methods (both telephone and postal surveys are acceptable only if open-ended questions are asked), qualitative evidence collected from interviews, observations | Type of papers: Papers that are not primary research e.g. systematic reviews, meta-analysis |
| Language of papers: English | Language of papers: Papers in other languages |
| Date: Papers published from 1984 onwards | Date: Papers were written before 1984 |
| Population of focus: unpaid family members or family carer | Population of focus: Population of focus relates to other groups of carers e.g. paid carers, professionals |
| Age group: 18 and above | Age group: Below 18 |
| Primary illness of interest: moderate to advanced dementia as defined by the authors of identified studies in the abstract or full paper | Primary illness of interest: Other illnesses |
| Setting: domestic home setting, unpaid home-based care | Setting: Other formal care establishments where care provision is paid for |
| Focus of papers: current experiences or views or needs of family carer | Focus of papers: Focus on the views of others, or where death has already occurred |
| Experience: Subjective experiences, such as family carers' views, opinion, attitudes, and the reasons for this, with respect to home-based caregiving for people with moderate to advanced dementia. | Experience: other non-family carers' views |
| Research Type: Research adopting either qualitative or mixed methods (described in terms of exposure to specific elements of caregiving responsibilities leading to a noticeable change in behaviour or wellbeing of the sample, accounts of the duration of time spent caregiving and family | Research Type: other types of research |

| | |
|---|--|
| carers' feelings or emotions during care provision were also of interest. | |
|---|--|

Table 2: Search terms and strategy

| Search terms and strategy |
|--|
| MH "Dementia+" OR TX dement* OR TX alzheimer * OR TX "lew* bod*" OR TX FTL D OR FTD OR frontotemporal |
| AND TX carer* OR TX caregiv* OR TX care-giver OR TX spouse-caregiver* |
| AND MH "Qualitative Research+" OR MH "qualitative studies" OR MH "Focus Groups" OR MH "exploratory+" OR MH "qualitative interview+" OR MH "ethnography+" OR MH "ethnographic research" OR MH "content analysis" OR MH "Grounded theory" |

Table 3: Characteristics of included studies

| Author/date/ Country | Setting | Population/ sample size/ sampling method/ illness stage | Aim/objectives | Research type/ research design | Data collection & analysis | Findings | Quality score | Quality Grade |
|--|-----------|---|--|---|--|---|---------------|---------------|
| Adams and Sanders (2004) [USA] | Homebased | Sample: Informal caregivers (n=74) Sampling method: Convenience sampling Stage: Moderate to advanced | To explore self-reported losses, grief reactions and depressive symptoms among a sample of caregivers of persons with Alzheimer's disease or other progressive dementia | Type: Mixed method Design: Cross-sectional | Data collection: Open-ended postal survey Analysis: Cross-sectional analysis | a) Caregivers for individuals within the late stage of dementia reported significantly more symptoms of grief and depression than those in the early or middle stages; b) The late-stage experience mostly resembled bereavement. | 29 | B |
| Brunton et al. (2008) [New Zealand] | Homebased | Sample: Family caregivers 1 st stage: (n=5) 2 nd stage: (n=64) Sampling method: Purposive, non-probability | To explore the lived experience and perceptions of a sample of caregivers who are providing informal care 24hr per day, seven days per week for those with Alzheimer's disease in New Zealand. | Type: Mixed method Design: Exploratory | Data collection: Semi-structured interview and questionnaires (using open-ended questions) Analysis: | a) High cost of caring is reported as a form of negative impact on caregiver's physical and psychological health; b) Difficulty in asking for help from family; c) Caregivers were concerned about finances and fears for the future; d) Caregivers reported also reported the need to be free, need for a time out, need for a role | 34 | A |

| | | | | | | | | |
|--|---------------------|---|---|--|--|--|----|---|
| | | Stage: Stage 2 to Stage 5 | | | Thematic analysis | change and a need to be someone else. | | |
| Chan et al. (2010) [Hong Kong] | Community homebased | Sample: Informal caregivers (n=27) Sampling method: Purposive Stage: Moderate to advanced | To report the lived experience of caregivers of persons with dementia in Hong Kong, and to explore their service needs. | Type: Qualitative Design: Focus group | Data collection: Focus group interviews Analysis: Colaizzi's phenomenological analysis | a) Caregivers were confused about the nature of their care recipients' disease; b) Negative report of caregiving in terms of grief, sadness, guilt, anger and fear; c) The needs of caregivers changed in line with the care recipient's disease progression; d) Caregiving responsibilities increased as the illness progressed; (e) Difficulty in coping with increasing care recipient's behavioural changes; (f) Increased caregiving responsibilities conflicted with other social roles, such as employment and other social activities; g) Burden reduces in the advanced stage of dementia as role acceptance increases. | 36 | A |
| Colling (2004) [USA] | Homebased | Sample: Family caregivers Moderate (n=16), Advanced (n=19) | To describe how the experience of passivity was for the caregiver and the person with Alzheimer's disease. | Type: Qualitative Design: Phenomenology | Data collection: Semi-structured interviews Analysis: content analysis (using Colaizzi's thematic extraction) | a) Passivity increased over time as the disease worsened;b) Experience of emotions such as fatigue, sadness, tears, frustration and a sense of helplessness; c) Caregivers of moderately impaired persons found it stressful to cope with lifestyle changes associated with caregiving; d) Caregivers of severely impaired persons expressed the most sadness | 35 | A |

| | | | | | | | | |
|---|-----------|---|--|---|--|---|----|---|
| | | <p>Sampling method: Purposive</p> <p>Stage: Moderate to advanced</p> | | | | associated with experiencing the process of cognition deterioration in their loved ones; e) at the moderate stage found it stressful to cope with lifestyle changes associated with caregiving, a sense of helplessness and frustration at watching behavioural changes in the care recipient | | |
| de La Cuesta-Benjumea (2011) [Spain] | Homebased | <p>Sample: Female family caregivers (n=23)</p> <p>Sampling method: Purposeful and theoretical</p> <p>Stage: Advanced</p> | To explore strategies women caregivers of relatives with advanced dementia use to rest from caregiving | <p>Type: Qualitative</p> <p>Design: Grounded theory</p> | <p>Data collection: Semi-structured interviews</p> <p>Analysis: Grounded theory (constant comparison analysis)</p> | a) Taking leave from the life of caregiving is used by the caregivers as a strategy to rest from their caregiver selves; (b) Caregivers did not perceive respite to mean resting or being idle. For them, respite is considered as doing something different, and by thinking and relating to others in different ways than those linked to caregiving. | 36 | A |
| De Silva and Curzio (2009) [UK] | Homebased | <p>Sample: Female family caregivers (n=10)</p> <p>Sampling method: Purposive</p> <p>Stage: Advanced</p> | To explore the life experiences of family carers in the transition from home into long-term care | <p>Type: Qualitative</p> <p>Design: Phenomenology</p> | <p>Data collection: One-to-one interviews</p> <p>Analysis: Thematic analysis</p> | a) Deterioration of behaviour is a major reason for care recipient transitioning into long-term care; (b) Although alternatives to long-term care were offered before placement, several did not receive much support and felt isolated; c) Spousal caregivers reported great sadness in placing their loved one; d) Difficulty in finding the right place; e) Severity of dementia and the complex health needs of a | 29 | B |

| | | | | | | | | |
|--|-----------|---|---|--|---|--|----|---|
| | | | | | | loved one left caregivers no other choice but long-term care; f) Experience of guilt or emotional distress during the transition. | | |
| Hirschman et al. (2006) [USA] | Homebased | Sample: Family caregivers (n=30) Sampling method: Purposive Stage: Advanced | To identify what standard of decision making a family member uses when making medical decisions for their relative with advanced dementia | Type: Qualitative Design: Qualitative | Data collection: Semi-structured interviews Analysis: Thematic analysis | a) Making medical decisions is difficult for caregivers, and there is no guidance as to how this is done; b) Either substituted judgement (what the care recipient would have wanted) (n=2) or best interest (opinion of the caregiver for deciding what is best for the care recipient) (n=17). In some cases, decisions are made by combining both as care recipients (n=11) often fail to have discussions about end of life with their family; c) Barriers highlighted include: avoiding the topic, waiting too long and denial of dementia by the care recipient and sometimes the caregivers themselves. | 35 | A |
| Idura et al. (2018) [Malaysia] | Homebased | Sample: Family caregivers (n=12) Sampling method: Purposive Stage: Moderate to advanced | To describe the experience of Malaysian family caregivers' experiences in providing care to their family members with moderate to advanced dementia | Type: Qualitative Design: Qualitative | Data collection: In-depth semi-structured interviews Analysis: Thematic analysis | a) Increased severity did not increase caregivers' burden due to acceptance of the caregiving role; b) Caregivers' grief increases with severity; c) Burden increases at the earlier stages; d) Caregivers felt it was their responsibility to provide care; e) Family support as a possible reason for a reduced burden; f) | 35 | A |

| | | | | | | | | |
|--------------------------------|-----------|---|--|--|--|---|----|---|
| | | | | | | Family support is a positive experience in helping to improve relationships among family members; g) Family support was not always available. | | |
| Karger (2018) [Germany] | Homebased | Sample: Relatives (n=20) Sampling method: Purposive Stage: Advanced | To gain insight into families and professionals' understanding of the emotional experience in patients with advanced Alzheimer's disease | Type: Qualitative Design: Focus group | Data collection: Group interviews Analysis: Qualitative content analysis and framework analysis | a) Relatives perceived emotionality as one of the essential characteristics retained despite the illness; b) loss of emotional experience increased with loss of cognitive capacity and verbal expression; c) shared memories faded with illness progression; d) memories forgotten no longer held any emotional significance for the care recipient; e) Perceived discontinuity of patient's emotional responses contrary to relatives' expectations creates ambiguity; f) Patients' behaviour which is different to expectations is a distressing experience for the relatives. | 32 | A |
| Lamahewa et al. (2018) [UK] | Homebased | Sample: Family caregivers Current (n=6) Sampling method: Purposive | To explore the difficulties in decision making for family carers and practitioners at the end of life | Type: Qualitative Design: Focus group | Data collection: Semi-structured interviews Analysis: Thematic analysis | Making decisions is particularly difficult at the end of life and associated with caregivers' unpreparedness | 32 | A |

| | | | | | | | | |
|---|----------------------------------|---|---|---|--|---|----|---|
| | | Stage: Advanced | | | | | | |
| Lethin et al. (2016) [Sweden] | Homebased (also nursing home) | Sample: Family caregivers (n=23) Sampling method: Strategic Stage: Advanced (includes accounts of early stages) | To investigate family caregivers' experiences of formal care when caring for a person with dementia through the stages of the disease | Type: Qualitative Design: Focus group | Data collection: Focus group interviews Analysis: Content analysis | a) Caregivers' feeling of inadequacy and striving to remain in control, while avoiding being a burden and struggling to avoid dependence on formal care; b) Worry about the future and the capacity to continue caregiving; c) Feelings of being alone and isolated because formal care was not as supportive as expected; d) Increasingly isolated when friends and family no longer came to visit | 36 | A |
| Lindauer et al. (2016) [USA] | Homebased | Sample: Family caregivers (n=11) Sampling method: Purposive Stage: Moderate to advanced | To explore the meaning AA family caregivers ascribed to dementia-related changes in care recipients | Type: Qualitative Design: Interpretative Phenomenological Approach | Data collection: In-depth interviews Analysis: Interpretative Phenomenological Approach (IPA) | a) Dementia-related changes meant that caregivers had to hang on to the care recipient for as long as possible; b) Caregiver's recognition that the valued care recipients were changed but still here and worthy of respect and compassion; c) Valued care recipients were changed, but still here, was regarded as a paradox; d) Family values, shaped by historical oppression, influenced caregiving meanings; e) Caregivers did not consider they were grieving, as their care recipient was <i>still here</i> . | 36 | A |

| | | | | | | | | |
|--|---|---|--|---|--|--|-----------|----------|
| <p>Moore et al. (2017)</p> <p>[UK]</p> | <p>Homebased</p> <p>(Part of a larger study including care-home as the setting)</p> | <p>Sample: Family carers (n=6)</p> <p>Sampling method: Stratified</p> <p>Stage: Advanced</p> | <p>To understand the experiences of carers during advanced dementia and to explore the links between mental health and experiences of end of life care.</p> | <p>Type: Mixed methods</p> <p>Design: Longitudinal cohort study</p> | <p>Data collection: In-depth interviews</p> <p>Analysis: Thematic analysis</p> | <p>a) Difficulty in understanding dementia progression and being involved during advanced stages; b) Placing care recipient in a care home is perceived as a loss of control, and a need for heightened vigilance; c) The unpredictability of dementia led to challenges in preparing for the end of life; d) Grief is a staged-process and high levels of guilt and grief into bereavement is linked to the inability to effect change in care provision</p> | <p>36</p> | <p>A</p> |
| <p>Robinson et al. (2010)</p> <p>[Canada]</p> | <p>Homebased</p> | <p>Sample: Family caregivers (n=29)</p> <p>Sampling method: Purposive</p> <p>Stage: Advanced</p> | <p>To understand how family caregivers of institutionalised relatives with dementia perceived and experienced relocation of their relative to a more home-like setting</p> | <p>Type: Mixed methods</p> <p>Design: Focus group</p> | <p>Data collection: Focus group and individual interviews</p> <p>Analysis: Thematic analysis</p> | <p>a) Caregivers often exhaust their physical, cognitive, emotional resources before agreeing to have their family member placed in care; b) The family caregiver's relational connection to the member with dementia remained strong and generated suffering when placement occurred that persisted over time and, for some, increased. This suffering took the form of grieving the losses arising from placement, guilt for being unable to prevent placement, second-guessing themselves with regards to the necessity of placement, and concern over the circumstances of the care.</p> | <p>35</p> | <p>A</p> |

| | | | | | | | | |
|---|-----------|--|--|--|---|--|----|---|
| Smith et al. (2001) [USA] | Homebased | Sample: Primary caregivers (family) (n=45) Sampling method: Purposive Stage: Mild to advanced | To explore caregivers' narratives of their experience and needs, and how the disease and their interaction with multiple practitioners have impacted their lives | Type: Qualitative Design: Ethnography | Data collection: Semi-structured intensive interviews Analysis: Thematic analysis | a) Family caregivers had financial worries, with those identified as spouses expressing higher levels of anxiety; b) Some caregivers' needs seemed more dependent upon unfulfilled expectations from family, support systems, or practitioners; c) Disappointment at the level of care, emotional support, or love expressed by family members | 35 | A |
| Tretteteig et al. (2017) [Norway] | Homebased | Sample: Family caregivers (n=5) Sampling method: Purposive Stage: Moderate to advanced | To gain knowledge about family caregivers' experiences of meaning in their caring role | Type: Qualitative Design: Narrative (using Riessman's method) | Data collection: In-depth semi-structured interviews Analysis: Narrative analysis method using a case study approach | a) Caregivers' roles and coping strategies were related to their relational ties; b) Difficulty in accepting their care recipient had changed; c) Caregivers had to make decisions whether to enhance, maintain or let go of emotional ties to find a good balance between meeting own needs and care recipient's needs; d) Letting go is almost impossible due to having a shared history with care recipient | 35 | A |
| Wladkowski (2016) [UK] | Homebased | Sample: Family caregivers (n=24) Sampling method: Purposive | To explore the grief experience in end-of-life care by caregivers | Type: Qualitative Design: Qualitative | Data collection: Semi-structured interviews Analysis: Thematic analysis | a) Unpreparedness for a change in terminal prognosis and uncertainty about being able to resume caregiving for their loved one following a live discharge; b) Emotional conflict in understanding disease progression following a live discharge, which also caused a struggle | 35 | A |

| | | | | | | | | |
|--|--|----------------------------------|--|--|--|---|--|--|
| | | Stage: <i>Advanced</i> | | | | in understanding care recipient's terminal prognosis as being temporary; c) A level of ambiguity in the caregivers' grief process as caregivers become unsure of how or when to grieve. | | |
|--|--|----------------------------------|--|--|--|---|--|--|