

## Supplemental Materials

**Supplemental Table E1.** Databases used for cohort construction and variable definitions

**Supplemental Table E2.** Variables used propensity score matching and hard matching

**Supplemental Table E3:** Interaction by sex (propensity score-matched cohort)

**Supplemental Figure E1.** Cohort construction

**Supplemental Table E4.** Cohort characteristics before propensity score matching

**Supplemental Figure E2:** Difference in annual median health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex

**Supplemental Figure E3:** Difference in annual 95th percentile health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex

**Supplemental Figure E4:** Sensitivity analyses censored at entrance to long-term care

**Supplemental Figure E5:** Differences in 6-month health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex, at the following distributions: (A) mean, (B) median, (C) 95<sup>th</sup> percentile, (D) 99<sup>th</sup> percentiles

**Supplemental Table E5:** Risk differences between test-positive and test-negative people for being a high-cost user within 6 months and 1 year, overall and by sex, defined as (A) >95th percentile and (B) any post-acute health care cost (>0)

**Supplemental Table E1.** Databases used for cohort construction and variable definitions. Additional details are available at <https://datadictionary.ices.on.ca/Applications/DataDictionary/Default.aspx>

<b>Database</b>	<b>Cost Components</b>	<b>Costing Methods (standardized to 2020CAD)</b>
Ontario Health Insurance Plan (OHIP)	Outpatient physician visits Laboratory services Non-physician services	-Fee-for-service + 'shadow billing' -Technical + professional fees
National Ambulatory Care Reporting System (NACRS)	Emergency department visits Dialysis visits Oncology clinic visits	Resource weight intensity x cost/weighted case
Discharge Abstract Database (DAD)	Inpatient hospitalizations	Resource weight intensity x cost/weighted case; as appropriate, prorated to exclude days prior to start of follow up
Client Agency Program Enrolment	Capitation costs	Base rate x age-sex multiplier + applicable premiums
Ontario Drug Benefit (ODB)	Medication use	Total payments
National Rehabilitation Reporting System	Rehabilitation admissions	Rehabilitation patient group x rehabilitation cost-weight
Continuing Care Reporting System	Complex continuing care admissions Long-term care	-Case mix index x cost/weighted day -Length of stay x cost/day
Home Care Database	Home care services	Standard cost per service
Mental Health Reporting System	Mental health admissions	(Length of stay x case mix index) x cost /weighted day
Same Day Surgery	Same day surgery	Resource weight intensity x cost/weighted case
<b>Other Databases</b>		
Registered Persons Database (RPDB)		Congestive Heart Failure (CHF)
Canadian Institute for Health Information (CIHI)		Ontario Asthma dataset (ASTHMA)
Yearly Health Services Contact		Chronic Obstructive Pulmonary Disease (COPD)
Ontario Marginalization Index (ONMARG)		Ontario Cancer Registry
Ontario COVID-19 Vaccine Data		Ontario Dementia Database (DEMENTIA)
COVID19 Integrated Testing Data		ICES Physician Database
Ontario Laboratories Information System		Corporate Provider Database
Ontario Hypertension Dataset (HYPER)		Ontario Diabetes Dataset (DIABETES)
<b>Descriptions of specialized health care services</b>		
Homecare	Providing of services for people of all ages who require care in their home, school, or in the community, and seniors with complex medical conditions who otherwise would be able to live in their home. Includes health professional home visits (e.g., nursing care, physio/physical therapy, occupational therapy, speech-language therapy, social work, nutrition support, home healthcare supplies, personal care support for activities of daily living, homemaking, and family-managed home care/self-directed care, and end-of-life care at home.	
Complex continuing care	Complex Continuing Care (CCC) provides continuing, medically complex and specialized services to people of all ages, sometimes over extended periods of time. CCC is offered in healthcare centres for people who have long-term illnesses or disabilities typically requiring skilled, technology-based care not available at home or in long-term care facilities. Patients must be medically stable and are often eligible if they have an uncuffed tracheostomy, peripherally inserted central catheter, feeding tube, or nephrostomy tube.	
Rehabilitation	Hospital beds for medically complex patients provide	

	<p>rehabilitation to gradually improve strength and movement. Care includes access to doctors or nurse practitioners, nursing, and physio (physical) and occupational therapy. Typical short-term stays are up to three months. Long-term patients are understood to be permanent residents unless conditions change.</p>
<p>Long-term care</p>	<p>Similar to nursing home care for medically complex patients in the United States. Long-term care homes provide homecare and services for people whose needs cannot be met in the community. Monthly out-of-pocket patient costs are determined by the Ministry and as of July 1, 2023 range from \$1,986.92-\$2,838.49 per month.</p>

**Supplemental Table E2.** Variables used propensity score matching and hard matching. Hard matched variables indicated by\*: sex, outbreak period, and 1-year baseline total healthcare costs. Please see Supplemental Table E1 for datasets.

Variable	Definition and/or source
Age	Years, restricted cubic spline
Sex*	Male, female (yes/no pregnant)
Outbreak period*	Two week blocks, beginning January 28, 2020
Public health unit	34 units
Neighbourhood income	Quintiles
Residential instability <sup>1</sup>	Quintiles, Ontario Marginalization Index; e.g., percentage living alone, dwellings not owned
Material deprivation <sup>1</sup>	Quintiles, Ontario Marginalization Index; e.g., percentage unemployed, without a high school degree
Dependency <sup>1</sup>	Quintiles, Ontario Marginalization Index; e.g., percentage of seniors, individuals not participating in the labour force
Ethnic concentration <sup>1</sup>	Quintiles, Ontario Marginalization Index; e.g., percentage of recent immigrants, those who self-identify as a "visible minority"
Rurality	Population <10,000
Diabetes	ICES DIABETES
Pregnancy	ICES MOMBABY <sup>2-5</sup>
Hypertension	ICES HYPER
Acute myocardial infarction	≥1 DAD ICD10: I21, I22, I25.2
Percutaneous coronary intervention	CIHI DAD/SDS: CCP: 48.02, 48.09; CCI: 1IJ50, 1IJ57GQ 1IJ80, 1IJ26, 1IJ54, 1IJ55; OHIP: Z434, Z448, Z449, Z460, Z461
Coronary artery bypass surgery	(CIHI DAD/SDS): CCI 1IJ76; CCP 48.1, 48.2; OHIP R742, R743
Ischemic stroke	One DAD, 2 OHIP, or 1 NACRS&1OHIP: ICD-10 codes I63, I64, H341 (excluding I63.6) as ANY diagnosis type, exclude suspect; ICD-9 code 434, 436 for OHIP
Haemorrhagic stroke	1 record in DAD or NACRS: ICD-10 I60, I61; ICD-9 430, 431
Major bleeding event	ICD 10: GI: I850, I983, K250/252/254/256, K260/262/264/266, K270/272/274/276, K280/282/284/286, K290, K661, K920, K921, K922 ICH: I60, I61, I620, I621, I629 GU: N020-029, R310, R311, R318 Resp: R040, R041, R042, R048, R049 Other: R58, D68.3, H35.6, H45.0, M25.0
Solid cancer, hematologic cancer	OCR: OCR_TOPOG_CD, OCR_DIAG_DATE
Same-day surgery in prior 6 weeks	SDS_ADMDATE, SDS_INCODE1-10, SDS_CACSANETECH
Valvular disease	ICD9: 394, 395, 396 ICD10: I019, I020, I05, I08, I099, I342, I348, I349 ICD10 code Z952, and CCI codes 1HS90LACF, 1HT90LACF, 1HU90DACF, 1HU90LACF, 1HU90PNCF, 1HV90LACF, 1HV90LACFA, 1HV90LACFL, 1HV90LACFN, 1HV90WJCFN
Emphysema	COPD_SPECIFIC in COPD database
Asthma	ASTHMA_SPEC in ASTHMA database
Atrial fibrillation	Any of the following: -history of hospitalization (CIHI DAD) <sup>6</sup> : ICD9 427.3 or ICD10 I48 as any diagnosis type, including suspected -history of ED visit with same codes -4 OHIP claims in 1 year (OHIP) <i>dxcode</i> 427
Heart failure	CHF database
Ischemic heart disease	PCI, CABG, (1 HOSP in DAD with any codes I20-I25) or (2 OHIP billings within a one-year period with dx codes 410-414)
Renal disease	DAD code: ICD10 codes E102, E112, E132, E142, I12, I13, N01.*, N03.*, N05.*, N08.*, N18.*, N19.*, N25.* or Chronic Dialysis (Any 2 codes within 90 days of one another): OHIP: R849, R850, G323, G325, G326, G330, G331, G860, G333, G083, G091, G085, G295, G082, G090, G092, G093, G094, G861, G862, G863, G864, G865, G866, G294, G095, G096 CCP: 51.95, 66.98

	NACRS: CCI: 1PZ21HQBR, 1PZ21HPD4
Pneumonia	(CIHI DAD, NACRS, OHIP) – ICD-10 codes J10.0, J11.0 or J12-J18 as ANY diagnosis type, exclude suspected <sup>7</sup> ; OHIP dxcode 486, excluding claims associated with fee codes G538, G539, G840-G848, G590, G591 or G700 (administration of vaccinations)
Dementia	(1 DAD or 3 OHIP billings separated by 30 days, within a 2-year period or any cholinesterase inhibitor from ODB) ICD10 codes F00-F03, F051, G30, G31, R54 ICD9 code 290, 294, 331, 797 SUBCLNAM= CHOLINESTERASE INHIBITORS ODB: donepezil, galantamine, or rivastigmine (DIN: 02232043, 02232044, 02269457, 02269465, 02244298, 02244299, 02244300, 02244302, 02266717, 02266725) or Tacrine (Cognex) (DIN: 66123288, 66123290, 66123306, 66123318)
Alcohol substance use disorder	ICD10 codes F1094, F1029, F1019, F1099, F10250, F10150, F10950, F10920, F10929, F10251, F10251, F10151, F10951, F1027, F1097, F1026, F1096, F10129, F10120, F1010, F1021, F1020, F10220, F10229 ICD9 codes 291, 303
Aggregated Diagnosis Group category	Aggregated Diagnosis Group (ADG) category group (from OHIP, NACRS-ED, CIHI DAD). The Johns Hopkins ACG© System Version 10.0 was used.
John Hopkins frailty indicator	(OHIP, NACRS-ED CIHI DAD)
Influenza vaccination	2019-20 influenza vaccination (OHIP, ODB) – Received between Sep 1, 2019 and COVID test date. Note: does not capture vaccinations received outside doctor's offices and pharmacies
Venous thromboembolism	CIHI-NACRS, DAD, OHIP ICD-10 I26.*, I26.0, I26.9, I80.* (excluding I80.0), I80.1, I80.2 (excluding I82.0, I82.1, I82.3), I80.3, I80.8, I80.9, I82.*, I82.2, I82.8, I82.9, OHIP DXCODE 415, 451, 452, 453
Mental health ED visit	OHIP <sup>8-10</sup>
MHA Outpatient Services	OHIP <sup>8-10</sup>
1-year baseline total healthcare costs*	See Supplemental Table E1
* variable used for hard matching	

### Supplemental Material References


- Matheson FI, Dunn JR, Smith KL, Moineddin R, Glazier RH. Development of the Canadian Marginalization Index: a new tool for the study of inequality. *Can J Public Health*. 2012;103(8 Suppl 2):S12-16.
- Aoyama K, Ray JG, Pinto R, et al. Temporal Variations in Incidence and Outcomes of Critical Illness Among Pregnant and Postpartum Women in Canada: A Population-Based Observational Study. *J Obstet Gynaecol Can*. 2019;41(5):631-640.
- Metcalfe A, Lix LM, Johnson JA, et al. Validation of an obstetric comorbidity index in an external population. *BJOG*. 2015;122(13):1748-1755.
- Joseph KS, Fahey J, Canadian Perinatal Surveillance S. Validation of perinatal data in the Discharge Abstract Database of the Canadian Institute for Health Information. *Chronic Dis Can*. 2009;29(3):96-100.
- Samiedaluie S, Peterson S, Brant R, Kaczorowski J, Norman WV. Validating abortion procedure coding in Canadian administrative databases. *BMC Health Serv Res*. 2016;16:255.
- Tu K, Nieuwlaet R, Cheng SY, et al. Identifying Patients With Atrial Fibrillation in Administrative Data. *Can J Cardiol*. 2016;32(12):1561-1565.
- Griffin MR, Zhu Y, Moore MR, Whitney CG, Grijalva CG. U.S. hospitalizations for pneumonia after a decade of pneumococcal vaccination. *N Engl J Med*. 2013;369(2):155-163.
- MHASEF Research Team. Mental Health and Addictions System Performance in Ontario: A Baseline Scorecard. Toronto, ON: Institute for Clinical Evaluative Sciences; 2018. Available from: <https://www.ices.on.ca/Publications/Atlases-and-Reports/2018/MHASEF>.
- MHASEF Research Team. The Mental Health of Children and Youth in Ontario: 2017 Scorecard. Toronto, ON: Institute for Clinical Evaluative Sciences; 2017. ISBN: 978-1-926850-72-6. Available from: <https://www.ices.on.ca/Publications/Atlases-and-Reports/2017/MHASEF>.
- ICES Data Dictionary. <https://datadictionary.ices.on.ca/Applications/DataDictionary/Default.aspx>. Last accessed February 11, 2022.

**Supplemental Table E3.** Interaction by sex (tested in matched cohort)

Component cost	P-value
Total	0.5908
Hospital	0.4838
Emergency department	0.5389
Outpatient medication	<b>0.0739*</b>
Outpatient specialists	0.5944
Outpatient cancer	0.2219
Outpatient primary care	<b>0.003*</b>
Outpatient laboratory	<b>&lt;.0001*</b>
Rehabilitation care	<b>0.0008*</b>
Homecare	<b>0.0037*</b>
Mental health	<b>0.0225*</b>
Complex continuing care	0.8936
Long-term care	<b>&lt;.0001*</b>

\* Significant, at P-value threshold of 0.10

Supplemental Figure E1: Cohort construction



Abbreviations: PCR, polymerase chain reaction; OHIP, Ontario Health Insurance Plan; PS, propensity score

**Supplemental Table E4.** Cohort characteristics before propensity score matching

		Total	Test Negative	Test Positive	Standardized Difference
		N=3,631,040	N=3,362,519	N=268,521	
Age (years)	Mean (SD)	46.58 (18.11)	46.77 (18.17)	44.18 (17.18)	0.146
	Median (Q1-Q3)	45 (31-60)	45 (31-60)	43 (29-56)	0.141
Sex	F - n (%)	1,970,075 (54.3%)	1,834,685 (54.6%)	135,390 (50.4%)	0.083
	M - n (%)	1,629,552 (44.9%)	1,498,276 (44.6%)	131,276 (48.9%)	0.087
	P - n (%)	31,413 (0.9%)	29,558 (0.9%)	1,855 (0.7%)	0.021
	Missing Data - n (%)				
Rural	n (%)	9,451 (0.3%)	8,780 (0.3%)	671 (0.2%)	0.002
	Y - n (%)	350,523 (9.7%)	339,985 (10.1%)	10,538 (3.9%)	0.244
Johns Hopkins Frailty Indicator	n (%)	123,999 (3.4%)	116,104 (3.5%)	7,895 (2.9%)	0.029
Adjusted Clinical Group Categories					
Aggregated Diagnosis Group Category 1	n (%)	2,782,083 (76.6%)	2,574,005 (76.5%)	208,078 (77.5%)	0.022
Aggregated Diagnosis Group Category 2	n (%)	2,731,200 (75.2%)	2,529,253 (75.2%)	201,947 (75.2%)	0
Aggregated Diagnosis Group Category 3	n (%)	2,235,629 (61.6%)	2,073,361 (61.7%)	162,268 (60.4%)	0.025
Aggregated Diagnosis Group Category 4	n (%)	195,459 (5.4%)	182,201 (5.4%)	13,258 (4.9%)	0.022
Aggregated Diagnosis Group Category 5	n (%)	1,085,536 (29.9%)	1,020,361 (30.3%)	65,175 (24.3%)	0.137
Aggregated Diagnosis Group Category 6	n (%)	1,523,716 (42.0%)	1,407,922 (41.9%)	115,794 (43.1%)	0.025
Aggregated Diagnosis Group Category 7	n (%)	193,342 (5.3%)	181,715 (5.4%)	11,627 (4.3%)	0.05
Aggregated Diagnosis Group Category 8	n (%)	240,396 (6.6%)	224,682 (6.7%)	15,714 (5.9%)	0.034
Aggregated Diagnosis Group Category 9	n (%)	339,984 (9.4%)	316,552 (9.4%)	23,432 (8.7%)	0.024
Aggregated Diagnosis Group Category 10	n (%)	1,263,741 (34.8%)	1,185,427 (35.3%)	78,314 (29.2%)	0.131
Aggregated Diagnosis Group Category 11	n (%)	1,183,663 (32.6%)	1,103,389 (32.8%)	80,274 (29.9%)	0.063
Aggregated Diagnosis Group Category 12	n (%)	165,525 (4.6%)	152,992 (4.5%)	12,533 (4.7%)	0.006
Recent flu vaccination	n (%)	1,130,869 (31.1%)	1,064,518 (31.7%)	66,351 (24.7%)	0.155
Hypertension	n (%)	877,852 (24.2%)	816,144 (24.3%)	61,708 (23.0%)	0.03
Diabetes	n (%)	433,350 (11.9%)	395,575 (11.8%)	37,775 (14.1%)	0.069
Emphysema	n (%)	97,824 (2.7%)	93,686 (2.8%)	4,138 (1.5%)	0.086
Heart failure	n (%)	93,663 (2.6%)	88,669 (2.6%)	4,994 (1.9%)	0.052
Dementia	n (%)	45,684 (1.3%)	42,410 (1.3%)	3,274 (1.2%)	0.004
Asthma	n (%)	442,882 (12.2%)	414,612 (12.3%)	28,270 (10.5%)	0.057
Cancer	n (%)	116,997 (3.2%)	112,188 (3.3%)	4,809 (1.8%)	0.098
Surgery in previous 6 weeks	n (%)	59,924 (1.7%)	57,709 (1.7%)	2,215 (0.8%)	0.08
Ischemic stroke	n (%)	45,455 (1.3%)	42,908 (1.3%)	2,547 (0.9%)	0.031
Hemorrhagic stroke	n (%)	3,757 (0.1%)	3,533 (0.1%)	224 (0.1%)	0.007
Valvular disease	n (%)	4,844 (0.1%)	4,609 (0.1%)	235 (0.1%)	0.015




Ischemic heart disease	n (%)	177,360 (4.9%)	167,514 (5.0%)	9,846 (3.7%)	0.065	
Major bleeding	n (%)	39,366 (1.1%)	37,210 (1.1%)	2,156 (0.8%)	0.031	
Renal disease	n (%)	46,728 (1.3%)	43,949 (1.3%)	2,779 (1.0%)	0.025	
Pneumonia	n (%)	291,202 (8.0%)	271,036 (8.1%)	20,166 (7.5%)	0.021	
Alcohol use disorder	n (%)	29,427 (0.8%)	27,759 (0.8%)	1,668 (0.6%)	0.024	
Venous thromboembolism	n (%)	420,456 (11.6%)	396,371 (11.8%)	24,085 (9.0%)	0.093	
Missing Data - n (%)		10,824 (0.3%)	10,050 (0.3%)	774 (0.3%)	0.002	
Income Quintile 1	n (%)	697,116 (19.2%)	630,532 (18.8%)	66,584 (24.8%)	0.147	
Income Quintile 2	n (%)	707,319 (19.5%)	649,204 (19.3%)	58,115 (21.6%)	0.058	
Income Quintile 3	n (%)	729,395 (20.1%)	672,162 (20.0%)	57,233 (21.3%)	0.033	
Income Quintile 4	n (%)	731,678 (20.2%)	684,228 (20.3%)	47,450 (17.7%)	0.068	
Income Quintile 5	n (%)	754,708 (20.8%)	716,343 (21.3%)	38,365 (14.3%)	0.184	
Missing Data - n (%)		39,428 (1.1%)	37,365 (1.1%)	2,063 (0.8%)	0.036	
Instability Quintile 1 (Least unstable)	n (%)	763,550 (21.0%)	691,223 (20.6%)	72,327 (26.9%)	0.15	
Instability Quintile 2	n (%)	671,203 (18.5%)	626,597 (18.6%)	44,606 (16.6%)	0.053	
Instability Quintile 3	n (%)	650,977 (17.9%)	610,179 (18.1%)	40,798 (15.2%)	0.079	
Instability Quintile 4	n (%)	654,660 (18.0%)	610,967 (18.2%)	43,693 (16.3%)	0.05	
Instability Quintile 5 (Most unstable)	n (%)	851,222 (23.4%)	786,188 (23.4%)	65,034 (24.2%)	0.02	
Missing Data - n (%)		39,428 (1.1%)	37,365 (1.1%)	2,063 (0.8%)	0.036	
Deprivation Quintile 1 (Least deprived)	n (%)	848,007 (23.4%)	802,901 (23.9%)	45,106 (16.8%)	0.177	
Deprivation Quintile 2	n (%)	755,915 (20.8%)	708,278 (21.1%)	47,637 (17.7%)	0.084	
Deprivation Quintile 3	n (%)	684,713 (18.9%)	632,209 (18.8%)	52,504 (19.6%)	0.019	
Deprivation Quintile 4	n (%)	650,256 (17.9%)	594,666 (17.7%)	55,590 (20.7%)	0.077	
Deprivation Quintile 5 (Most deprived)	n (%)	652,721 (18.0%)	587,100 (17.5%)	65,621 (24.4%)	0.172	
Missing Data - n (%)		39,428 (1.1%)	37,365 (1.1%)	2,063 (0.8%)	0.036	
Dependency Quintile 1 (Least dependent)	n (%)	993,323 (27.4%)	901,394 (26.8%)	91,929 (34.2%)	0.162	
Dependency Quintile 2	n (%)	731,303 (20.1%)	671,373 (20.0%)	59,930 (22.3%)	0.058	
Dependency Quintile 3	n (%)	626,297 (17.2%)	582,059 (17.3%)	44,238 (16.5%)	0.022	
Dependency Quintile 4	n (%)	594,603 (16.4%)	557,097 (16.6%)	37,506 (14.0%)	0.072	
Dependency Quintile 5 (Most dependent)	n (%)	646,086 (17.8%)	613,231 (18.2%)	32,855 (12.2%)	0.168	
Missing Data - n (%)		39,428 (1.1%)	37,365 (1.1%)	2,063 (0.8%)	0.036	
Ethnic Concentration Quintile 1 (Least)	n (%)	552,466 (15.2%)	535,172 (15.9%)	17,294 (6.4%)	0.304	
Ethnic Concentration Quintile 2	n (%)	604,881 (16.7%)	579,980 (17.2%)	24,901 (9.3%)	0.237	
Ethnic Concentration Quintile 3	n (%)	670,672 (18.5%)	635,538 (18.9%)	35,134 (13.1%)	0.159	
Ethnic Concentration Quintile 4	n (%)	783,968 (21.6%)	726,900 (21.6%)	57,068 (21.3%)	0.009	
Ethnic Concentration Quintile 5 (Most)	n (%)	979,625 (27.0%)	847,564 (25.2%)	132,061 (49.2%)	0.512	
Missing Data - n (%)		9,838 (0.3%)	9,146 (0.3%)	692 (0.3%)	0.003	
Public health unit		2226 - n (%)	26,512 (0.7%)	26,287 (0.8%)	225 (0.1%)	0.106
		2227 - n (%)	38,987 (1.1%)	37,168 (1.1%)	1,819 (0.7%)	0.046
		2230 - n (%)	162,703 (4.5%)	151,132 (4.5%)	11,571 (4.3%)	0.009
		2233 - n (%)	36,617 (1.0%)	35,864 (1.1%)	753 (0.3%)	0.096

2234 - n (%)	24,956 (0.7%)	23,709 (0.7%)	1,247 (0.5%)	0.032	
2235 - n (%)	50,556 (1.4%)	49,540 (1.5%)	1,016 (0.4%)	0.115	
2236 - n (%)	145,938 (4.0%)	137,409 (4.1%)	8,529 (3.2%)	0.049	
2237 - n (%)	146,639 (4.0%)	136,426 (4.1%)	10,213 (3.8%)	0.013	
2238 - n (%)	43,705 (1.2%)	43,262 (1.3%)	443 (0.2%)	0.132	
2239 - n (%)	12,439 (0.3%)	12,084 (0.4%)	355 (0.1%)	0.046	
2240 - n (%)	30,100 (0.8%)	28,752 (0.9%)	1,348 (0.5%)	0.043	
224 n (%)	55,293 (1.5%)	54,644 (1.6%)	649 (0.2%)	0.144	
2242 - n (%)	37,243 (1.0%)	34,901 (1.0%)	2,342 (0.9%)	0.017	
2243 - n (%)	45,285 (1.2%)	44,218 (1.3%)	1,067 (0.4%)	0.1	
2244 - n (%)	119,934 (3.3%)	114,752 (3.4%)	5,182 (1.9%)	0.092	
2246 - n (%)	115,348 (3.2%)	108,164 (3.2%)	7,184 (2.7%)	0.032	
2247 - n (%)	31,184 (0.9%)	30,843 (0.9%)	341 (0.1%)	0.11	
2249 - n (%)	24,286 (0.7%)	23,711 (0.7%)	575 (0.2%)	0.073	
225 n (%)	246,887 (6.8%)	234,860 (7.0%)	12,027 (4.5%)	0.108	
2253 - n (%)	371,949 (10.2%)	319,731 (9.5%)	52,218 (19.4%)	0.285	
2254 - n (%)	18,815 (0.5%)	18,016 (0.5%)	799 (0.3%)	0.037	
2255 - n (%)	41,455 (1.1%)	40,826 (1.2%)	629 (0.2%)	0.116	
2256 - n (%)	21,154 (0.6%)	20,864 (0.6%)	290 (0.1%)	0.085	
2257 - n (%)	22,990 (0.6%)	22,621 (0.7%)	369 (0.1%)	0.084	
2258 - n (%)	57,719 (1.6%)	55,263 (1.6%)	2,456 (0.9%)	0.065	
2260 - n (%)	157,395 (4.3%)	151,377 (4.5%)	6,018 (2.2%)	0.125	
226 n (%)	58,624 (1.6%)	57,534 (1.7%)	1,090 (0.4%)	0.128	
2262 - n (%)	45,934 (1.3%)	43,859 (1.3%)	2,075 (0.8%)	0.052	
2263 - n (%)	8,479 (0.2%)	8,348 (0.2%)	131 (0.0%)	0.052	
2265 - n (%)	134,215 (3.7%)	125,166 (3.7%)	9,049 (3.4%)	0.019	
2266 - n (%)	81,639 (2.2%)	77,296 (2.3%)	4,343 (1.6%)	0.049	
2268 - n (%)	100,882 (2.8%)	90,266 (2.7%)	10,616 (4.0%)	0.071	
2270 - n (%)	290,514 (8.0%)	263,253 (7.8%)	27,261 (10.2%)	0.081	
2275 - n (%)	52,650 (1.4%)	50,459 (1.5%)	2,191 (0.8%)	0.064	
3895 - n (%)	762,176 (21.0%)	680,768 (20.2%)	81,408 (30.3%)	0.233	
≥2 COVID vaccine doses	n (%)	72,170 (2.0%)	71,870 (2.1%)	300 (0.1%)	0.193
1-2 COVID vaccine doses	n (%)	79,847 (2.2%)	78,444 (2.3%)	1,403 (0.5%)	0.153
0 COVID vaccine doses	n (%)	3,479,023 (95.8%)	3,212,205 (95.5%)	266,818 (99.4%)	0.245
Study period (bi-weekly, from Jan 2020)	0 - n (%)	10 (0.0%)	*5-9	*1-5	0.001
	1 - n (%)	13 (0.0%)	*8-12	*1-5	0
	2 - n (%)	101 (0.0%)	101 (0.0%)	0 (0.0%)	0.147
	3 - n (%)	131 (0.0%)	*126-130	*1-5	0.079
	4 - n (%)	1,048 (0.0%)	*1006-1010	*38-42	0.162
	5 - n (%)	12,254 (0.3%)	10,887 (0.3%)	1,367 (0.5%)	0.048
	6 - n (%)	16,312 (0.4%)	12,596 (0.4%)	3,716 (1.4%)	0.04
	7 - n (%)	23,678 (0.7%)	18,963 (0.6%)	4,715 (1.8%)	0.09
	8 - n (%)	29,917 (0.8%)	25,799 (0.8%)	4,118 (1.5%)	0.059
	9 - n (%)	30,957 (0.9%)	27,856 (0.8%)	3,101 (1.2%)	0.061
	10 - n (%)	58,964 (1.6%)	55,379 (1.6%)	3,585 (1.3%)	0.008
	11 - n (%)	76,898 (2.1%)	74,742 (2.2%)	2,156 (0.8%)	0.008


	12 - n (%)	84,557 (2.3%)	82,824 (2.5%)	1,733 (0.6%)	0.01
	13 - n (%)	83,925 (2.3%)	82,666 (2.5%)	1,259 (0.5%)	0.029
	14 - n (%)	90,954 (2.5%)	89,383 (2.7%)	1,571 (0.6%)	0.108
	15 - n (%)	91,605 (2.5%)	90,641 (2.7%)	964 (0.4%)	0.111
	16 - n (%)	93,668 (2.6%)	92,593 (2.8%)	1,075 (0.4%)	0.072
	17 - n (%)	97,510 (2.7%)	95,957 (2.9%)	1,553 (0.6%)	0.033
	18 - n (%)	139,751 (3.8%)	135,916 (4.0%)	3,835 (1.4%)	0.026
	19 - n (%)	162,757 (4.5%)	155,950 (4.6%)	6,807 (2.5%)	0.117
	20 - n (%)	138,532 (3.8%)	130,729 (3.9%)	7,803 (2.9%)	0.166
	21 - n (%)	129,192 (3.6%)	119,634 (3.6%)	9,558 (3.6%)	0.165
	22 - n (%)	144,730 (4.0%)	130,629 (3.9%)	14,101 (5.3%)	0.191
	23 - n (%)	168,272 (4.6%)	151,423 (4.5%)	16,849 (6.3%)	0.19
	24 - n (%)	198,286 (5.5%)	177,663 (5.3%)	20,623 (7.7%)	0.176
	25 - n (%)	197,849 (5.4%)	173,237 (5.2%)	24,612 (9.2%)	0.161
	26 - n (%)	215,184 (5.9%)	179,401 (5.3%)	35,783 (13.3%)	0.113
	27 - n (%)	197,754 (5.4%)	172,722 (5.1%)	25,032 (9.3%)	0.054
	28 - n (%)	174,529 (4.8%)	158,981 (4.7%)	15,548 (5.8%)	0
	29 - n (%)	184,128 (5.1%)	172,621 (5.1%)	11,507 (4.3%)	0.065
	30 - n (%)	233,494 (6.4%)	221,366 (6.6%)	12,128 (4.5%)	0.097
	31 - n (%)	274,403 (7.6%)	257,838 (7.7%)	16,565 (6.2%)	0.156
	32 - n (%)	279,677 (7.7%)	262,863 (7.8%)	16,814 (6.3%)	0.277
Prior mental health hospitalization	n (%)	82,415 (2.3%)	77,945 (2.3%)	4,470 (1.7%)	0.047
Prior emergency department visit for mental health condition	n (%)	200,301 (5.5%)	188,832 (5.6%)	11,469 (4.3%)	0.062
Prior outpatient mental health visit	n (%)	796,358 (21.9%)	747,646 (22.2%)	48,712 (18.1%)	0.102
		1726.91	1764.50	1256.21	
1-year Baseline hospital-based costs	Mean (SD)	(10159.99)	(10206.32)	(9548.37)	0.051
	Median (Q1-Q3)	0 (0-336)	0 (0-336)	0 (0-0)	0.212
				240.44	
1-year Baseline emergency department costs	Mean (SD)	312.49 (975.37)	318.25 (971.10)	(1024.54)	0.078
	Median (Q1-Q3)	0 (0-278)	0 (0-289)	0 (0-143)	0.114
				451.54	
1-yr Baseline medication costs	Mean (SD)	610.82 (4140.93)	623.54 (4226.39)	(2857.37)	0.048
	Median (Q1-Q3)	0 (0-50)	0 (0-58)	0 (0-14)	0.152
				669.21	
1-yr Baseline outpatient-specialist costs	Mean (SD)	885.05 (4744.63)	902.28 (4790.92)	(4115.09)	0.052
	Median (Q1-Q3)	140 (0-642)	145 (0-658)	80 (0-457)	0.161
				139.28	
1-yr Baseline outpatient-cancer costs	Mean (SD)	341.02 (5058.17)	357.13 (5174.60)	(3259.14)	0.05
	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.073
1-yr Baseline outpatient-primary care costs	Mean (SD)	380.35 (897.18)	384.18 (913.27)	332.40 (661.60)	0.065
	Median (Q1-Q3)	247 (125-414)	248 (128-416)	222 (99-384)	0.128
1-yr Baseline outpatient-lab costs	Mean (SD)	53.37 (91.71)	53.59 (92.17)	50.53 (85.69)	0.034
	Median (Q1-Q3)	21 (0-71)	21 (0-71)	21 (0-70)	0.019
1-yr Baseline rehabilitation costs	Mean (SD)	103.48 (1967.55)	105.10 (1983.56)	83.17 (1754.51)	0.012
	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.019
				223.09	
1-yr Baseline homecare costs	Mean (SD)	247.50 (2241.14)	249.45 (2226.99)	(2411.07)	0.011

	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.077
1-yr Baseline mental health admission costs	Mean (SD)	203.84 (5604.68)	206.59 (5581.52)	169.40 (5886.90)	0.006
	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.042
1-yr Baseline complex and continuing care costs	Mean (SD)	139.15 (4539.02)	140.46 (4587.99)	122.74 (3873.60)	0.004
	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.003
1-yr Baseline long-term care costs	Mean (SD)	0.08 (57.29)	0.09 (59.51)	0.02 (5.47)	0.002
	Median (Q1-Q3)	0 (0-0)	0 (0-0)	0 (0-0)	0.002

**Supplemental Figure E2:** Difference in annual median health care costs (95% confidence intervals in 2020 starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals and by sex




**Supplemental Figure E3:** Difference in annual 95th percentile health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex




**Supplemental Figure E4: Sensitivity analysis excluding long-term care costs**


**(A)** Differences in annual mean health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex




**(B)** Differences in annual median health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex



**(C)** Differences in annual 95th percentile health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex




(D) Differences in annual 99th percentile health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex






**Supplemental Figure E5:** Differences in 6-month health care costs (95% confidence intervals in 2020 \$CAD, starting 56 days after polymerase chain reaction test) for test-positive versus test-negative matched individuals, overall and by sex, at the following distributions: (A) Mean, (B) Median, (C) 95<sup>th</sup> percentile, (D) 99<sup>th</sup> percentile


**(A) Mean**




**(B) Median**



**(C) 95<sup>th</sup> percentile**



**(D) 99<sup>th</sup> percentile**



**Supplemental Table E5:** Risk differences between test-positive and test-negative people for being a high-cost user within 6 months and 1 year, overall and by sex, defined as (A) >95th percentile and (B) any post-acute health care cost (\$>0)

(A) High-cost defined as cost>95th percentile		6-months						1-year					
		# in T+	# in T-	Δ% T+ vs. T-	RD lower 95%CI	RD upper 95%CI	P-value	# in T+	# in T-	Δ% T+ vs. T-	RD lower 95%CI	RD upper 95%CI	P-value
Overall (n=265,591)	Outpatient cancer	13997	12564	0.0054	0.0042	0.0065	<.0001	13630	12934	0.00	0.00	0.00	<.0001
	Rehabilitation	14369	12726	0.0062	0.0050	0.0073	<.0001	13575	12986	0.00	0.00	0.00	0.0002
	Home care	13756	12825	0.0035	0.0023	0.0047	<.0001	13410	13161	0.00	0.00	0.00	0.1140
	Mental health hospitalization	14084	12482	0.0060	0.0049	0.0072	<.0001	13548	13022	0.00	0.00	0.00	0.0008
	Complex continuing care	14418	12183	0.0084	0.0073	0.0096	<.0001	13998	12595	0.01	0.00	0.01	<.0001
	Long-term care	13794	12943	0.0032	0.0020	0.0044	<.0001	13687	13060	0.00	0.00	0.00	<.0001
Female (n=135,760)	Outpatient cancer	7622	6613	0.0074	0.0058	0.0091	<.0001	7073	6502	0.00	0.00	0.01	<.0001
	Rehabilitation	8353	7108	0.0092	0.0075	0.0109	<.0001	7591	6976	0.00	0.00	0.01	<.0001
	Home care	7530	6984	0.0040	0.0023	0.0057	<.0001	7380	7202	0.00	0.00	0.00	0.1275
	Mental health hospitalization	8269	7179	0.0080	0.0063	0.0097	<.0001	7890	7584	0.00	0.00	0.00	0.0104
	Complex continuing care	8618	7218	0.0103	0.0086	0.0121	<.0001	8384	7477	0.01	0.00	0.01	<.0001
	Long-term care	9185	8463	0.0053	0.0035	0.0072	<.0001	8988	8394	0.00	0.00	0.01	<.0001
Male (n=129,831)	Outpatient cancer	6375	5951	0.0033	0.0017	0.0049	<.0001	6557	6432	0.00	0.00	0.00	0.2495
	Rehabilitation	6016	5618	0.0031	0.0015	0.0046	0.0001	5984	6010	0.00	0.00	0.00	0.8055
	Home care	6226	5841	0.0030	0.0014	0.0046	0.0003	6030	5959	0.00	0.00	0.00	0.5019
	Mental health hospitalization	5815	5303	0.0039	0.0024	0.0055	<.0001	5658	5438	0.00	0.00	0.00	0.0305
	Complex continuing care	5800	4965	0.0064	0.0049	0.0079	<.0001	5614	5118	0.00	0.00	0.01	<.0001
	Long-term care	4609	4480	0.0010	-0.0004	0.0024	0.1671	4699	4666	0.00	0.00	0.00	0.7269

<b>(B) Any post-acute health care cost</b>		<b>6-months</b>						<b>1-year</b>					
	<b>Outcome</b>	<b># in T+</b>	<b># in T-</b>	<b>Δ% T+ vs. T-</b>	<b>RD lower 95%CI</b>	<b>RD upper 95%CI</b>	<b>P-value</b>	<b># in T+</b>	<b># in T-</b>	<b>Δ% T+ vs. T-</b>	<b>RD lower 95%CI</b>	<b>RD upper 95%CI</b>	<b>P-value</b>
Overall (n=265,591)	Outpatient cancer	1739	2369	-	-0.0028	-0.0019	<.0001	2428	3071	0.00	0.00	0.00	<.0001
	Rehabilitation	744	308	0.0016	0.0014	0.0019	<.0001	954	570	0.00	0.00	0.00	<.0001
	Home care	10740	9780	0.0036	0.0026	0.0046	<.0001	13109	12339	0.00	0.00	0.00	<.0001
	Mental health hospitalization	698	717	-	-0.0003	0.0002	0.6127	1008	1127	0.00	0.00	0.00	0.0097
	Complex continuing care	1138	554	0.0022	0.0019	0.0025	<.0001	1327	792	0.00	0.00	0.00	<.0001
	Long-term care	1064	432	0.0024	0.0021	0.0027	<.0001	1390	682	0.00	0.00	0.00	<.0001
Female (n=135,760)	Outpatient cancer	1054	1385	-	-0.0031	-0.0017	<.0001	1468	1788	0.00	0.00	0.00	<.0001
	Rehabilitation	301	136	0.0012	0.0009	0.0015	<.0001	404	280	0.00	0.00	0.00	<.0001
	Home care	5894	5214	0.0050	0.0035	0.0065	<.0001	7165	6545	0.00	0.00	0.01	<.0001
	Mental health hospitalization	299	336	-	-0.0006	0.0001	0.1408	447	531	0.00	0.00	0.00	0.0070
	Complex continuing care	562	257	0.0022	0.0018	0.0027	<.0001	667	367	0.00	0.00	0.00	<.0001
	Long-term care	619	228	0.0029	0.0025	0.0033	<.0001	807	371	0.00	0.00	0.00	<.0001
Male (n=129,831)	Outpatient cancer	685	984	-	-0.0029	-0.0017	<.0001	960	1283	0.00	0.00	0.00	<.0001
	Rehabilitation	443	172	0.0021	0.0017	0.0025	<.0001	550	290	0.00	0.00	0.00	<.0001
	Home care	4846	4566	0.0022	0.0007	0.0036	0.0028	5944	5794	0.00	0.00	0.00	0.1492
	Mental health hospitalization	399	381	0.0001	-0.0003	0.0006	0.5187	561	596	0.00	0.00	0.00	0.3010
	Complex continuing care	576	297	0.0021	0.0017	0.0026	<.0001	660	425	0.00	0.00	0.00	<.0001
	Long-term care	445	204	0.0019	0.0015	0.0022	<.0001	583	311	0.00	0.00	0.00	<.0001