- Transcriptomic Response of Postprandial Blood, Subcutaneous Adipose Tissue
 and Muscle to a combined lifestyle intervention in older adults
- 3 F.A. Bogaards^{1,2,3}, T. Gehrmann^{1,2}, M. Beekman¹, N. Lakenberg¹, E. Suchiman¹, C.P.G.M. de
- 4 Groot³, M.J.T. Reinders^{2,4} and P.E. Slagboom^{1, 4,5}
- ⁵ ¹Molecular Epidemiology, Leiden University Medical Center, Leiden, The Netherlands, ²Leiden
- 6 Computational Biology Center, Leiden, The Netherlands, ³Human Nutrition and Health,
- 7 Wageningen University & Research, Division of Human Nutrition, Wageningen, The
- 8 Netherlands,⁴Delft Bioinformatics Lab, Delft University of Technology, Delft, The
- 9 Netherlands. ⁵Max Planck Institute for Biology of Aging, Cologne, Germany.

10 Abstract

Molecular effects of lifestyle interventions are typically studied in a single tissue. Here, we 11 investigated the sex-specific effects of the Growing Old TOgether (GOTO) study, a moderate 12 13 13-week combined lifestyle intervention on the transcriptomes of postprandial blood, 14 subcutaneous adipose tissue (SAT) and muscle tissue in healthy older adults, the overlap in 15 effect between tissues and their relation to whole-body parameters of metabolic health. The 16 GOTO intervention had virtually no effect on the postprandial blood transcriptome, while the 17 SAT and muscle transcriptomes responded significantly. In SAT, pathways involved in HDL 18 remodeling, O₂/CO₂ exchange and signaling were overrepresented, while in muscle, collagen 19 and extracellular matrix pathways were significantly overexpressed. Additionally, we found that the effects of the SAT transcriptome closest associated with gains in metabolic health. 20 21 Lastly, in males, we identified a shared variation between the transcriptomes of the three 22 tissues. We conclude that the GOTO intervention had a significant effect on metabolic and 23 muscle fibre pathways in the SAT and muscle transcriptome, respectively. Aligning the NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

response in the three tissues revealed a blood transcriptome component which may act as an
integrated health marker for metabolic intervention effects across tissues.

26 Main

Worldwide, the proportion of older people in our society is increasing, and with it the burden of age-related diseases on the quality of life and health economics ^{1,2}. Improving metabolic health is considered key to improving vitality at older ages ^{3,4}. Dietary, activity-based, or combined lifestyle interventions are known to improve culprit factors in human ageing such as metabolic health and immunity. Although effects are usually heterogeneous, the metabolic health can significantly be improved, even in middle and older age ^{5–10}.

33 The effect of lifestyle interventions can be studied by traditional blood-based markers of metabolic health (including: BMI, fasting insulin, systolic blood pressure (SBP), HDL 34 35 cholesterol, diastolic blood pressure (DBP), dual-energy X-ray absorptiometry (DEXA)-based 36 whole-body fat%, trunk fat%)^{6,11} and a range of omics based assays such as the metabolome $^{6,12-14}$, proteome 15,16 , epigenome 8 and the transcriptome $^{17-27}$, of blood $^{6,12-17,20,28,29}$, gut 15 , 37 adipose tissue ^{18,21,26,30,31}, and muscle tissue ^{19,22,23,27}. Human intervention studies are usually 38 small (n = 120 - 250) ^{6,29,32-34} and focus on one tissue only, mainly blood. Mostly, a strong 39 effect is observed on the fasting blood proteome ¹⁶ and metabolome ⁶, but typically not on 40 the fasting transcriptome, where the effects are milder and more variable ^{17,24,35}, except for 41 rigorous lifestyle interventions ²⁹. The transcriptomes of adipose tissue and muscle do react 42 respectively on separate dietary ²⁵ and physical activity ^{19,22,23,27} lifestyle interventions. In 43 mice, combined lifestyle interventions were shown to affect the transcriptome of brown 44 45 adipose tissue and skeletal muscle ³⁶.

In humans, however, the simultaneous effect of combined lifestyle interventions on 46 the transcriptome of blood, subcutaneous adipose tissue (SAT) and muscle was not 47 investigated as yet. To understand how differences in omics profiles relate to changes over 48 49 the years in health parameters one often relies on population-based omics studies. However, such studies mainly rely on blood samples ^{37,38}. We speculate that some of the transcriptomic 50 51 changes in different tissues in response to interventions align due to the systemic nature of 52 lifestyle interventions and that a part of changes in the blood transcriptome reflect responses 53 in SAT and muscle and related health gains. Hereto, an integrative analysis of the SAT, muscle 54 and blood transcriptomic response to such intervention is necessary.

55 Integrative data analysis methods pool data from different datasets to jointly analyze them ³⁹. Joint and Independent Variation Explained (JIVE) ⁴⁰ is one such integrative data 56 analysis method. JIVE decomposes the variation across the different datasets in three 57 58 components: 1) the joint variation between the datasets, 2) dataset-specific variations, and 59 3) the residual noise for each dataset separately. The blood transcriptome response in mild 60 interventions is usually minimal, JIVE however may reveal a part that represents the joint 61 response variation across tissues. We are interested whether the joint response captures the intervention effects on metabolic health stronger than the blood transcriptome alone. 62

We applied the tissue-specific as well as the integrative data analysis approach on a 13-week combined physical activity and dietary lifestyle intervention study in older adults; the Growing Old TOgether (GOTO) study. Within this study the transcriptome of postprandial blood, SAT and muscle tissue before and after intervention has been measured. We report on genes whose expressions within each of the different tissues associated with whole body health improvements across the intervention, and report on the observed joint transcriptional effect. We conclude that especially the SAT and muscle transcriptome respond

to the intervention by HDL remodeling, cell signaling and CO₂/O₂ exchange for SAT, and collagen/extracellular matrix pathways for muscle. Especially the SAT transcriptome in both sexes associates to beneficial changes in whole-body composition. The blood transcriptome does not generally respond to intervention or reflect health gain. Our integrative analysis of blood, however, revealed that the joint effect of the three tissues associated with the parameters of whole-body composition parameters. This blood transcriptome component may act as an integrated health marker for metabolic intervention effects across tissues.

77 Results

78 Health markers improve but improvement varies among subsets

79 Participants compliant to the intervention overall improved their metabolic health (Table 1) 80 with for some parameters slight sex-differences in magnitude of effect. We have 81 transcriptome data available in subsets of the complete GOTO study group: 1) Overlapping 82 Blood, SAT and Muscle 57, which consists of 33 males and 24 females, 2) Blood Postprandial 83 Samples 88, which contain 45 males and 43 females, 3) SAT Postprandial samples 78, which contains 38 males and 40 females, and 4) Muscle Postprandial samples 82, which contains, 84 85 48 males and 34 females. The direction of effect on the health parameters in the subsets of participants for which data in different tissues is available was similar (Table 1), though there 86 were some differences in significance level. For example, in males, the intervention only had 87 88 a significant effect on HDL cholesterol in the group with muscle transcriptome measurements. 89 While in females, HDL cholesterol size was significantly different in only the participants with 90 SAT transcriptome measurements, and the FRS was significantly different in the complete set 91 of measurements. These differences in effect could be partially explained by slight baseline 92 classical metabolic health marker level differences between the groups (Supplementary Table

1). To circumvent the slight differences in intervention effect on the classical metabolic health
markers between the different groups, the associations between the transcriptome and the
classical metabolic health markers will be only be calculated on the overlapping postprandial
blood, SAT and muscle samples.

97 Strong sex-dependent transcriptional response in subcutaneous adipose tissue and

98 muscle.

99 We investigated the effect of the GOTO intervention by measuring transcriptional differences 100 in blood, SAT and muscle (differential gene expression analysis, see Materials and Methods). 101 In males, the number of significantly differentially expressed genes (DEGs) across the 102 intervention varied considerably between blood (one), SAT (89), and muscle (251) (Figure 1, 103 Table 2, Supplementary Table 2, 3 & 4). The ratio between up and down regulated DEGs also 104 differed between the affected tissues: a slightly larger number of downregulated genes in SAT 105 whereas over 80% of the DEGs were upregulated in muscle. Noteworthy, the muscle DEGs 106 contained 14 collagen genes, all of which were upregulated (Supplementary Table 4).

For females, the number of DEGs across the different tissues were somewhat similar as in males, but the effect sizes were smaller (Figure 1, Table 2): zero DEGs in blood, 168 in SAT, and 168 in muscle (Supplementary Table 5 & 6). Like the male samples, more than half of the DEGs in SAT were downregulated, while in muscle over two-thirds of significant DEGs were upregulated. The female muscle DEGs contained 7 collagen genes, all of them upregulated (Supplementary Table 6).

Overall, in both males and females, the intervention had hardly any effect on the blood transcriptome while in SAT both up- and downregulated genes were observed, whereas in muscle the majority of the DEGs were upregulated.

116 Lipid and collagen related pathways enriched in SAT and muscle response, respectively

117 We then investigated which pathways were associated with the found DEGs (pathway 118 overrepresentation analysis, see Materials and Methods). In male SAT, five pathways were significantly overrepresented (Figure 2, Supplementary Table 7). The three upregulated 119 pathways were involved with HDL remodeling, nuclear receptor signaling and cholesterol 120 121 transport, while the downregulated pathway was involved in fertilization, the remaining 122 pathway (equally down - and upregulated) was involved in ligand clearance. In male muscle, all 18 enriched pathways were upregulated. The majority of these were involved in collagen 123 formation or extracellular matrix formation/degradation (Figure 2, Supplementary Table 7). 124 125 The remaining pathways were involved in myogenesis (NCAM1 pathways), mesenchymal 126 stem cell differentiation (PDGF pathway), scavenger receptors (binding and uptake, class A 127 receptors) and protein O-glycosylation.

128 In female SAT, seven pathways were significantly overrepresented (Figure 2, 129 Supplementary Table 7). The three upregulated pathways were involved in signaling 130 pathways and HDL remodeling. The four downregulated pathways were related to MAPK signaling and CO_2/O_2 exchange. However, the three CO_2/O_2 pathways were only 131 overrepresented due to the same three genes (Supplementary Table 7). In female muscle, all 132 133 eight overrepresented pathways were upregulated (Figure 2, Supplementary Table 7), six of 134 which were involved in collagen formation and extracellular matrix organization/degradation. 135 The remaining pathways were involved in in netrin-1 signaling and osteoblast differentiation. Taken together, both in males and females, the overrepresented pathways in muscle 136

137 were largely involved in collagen and extracellular matrix remodeling. In SAT, there were 138 fewer enriched pathways which were less consistent in terms of direction (both -up and

downregulated). The overrepresented pathways in SAT were involved in signaling, O₂/CO₂
 exchange, HDL remodeling and cellular signaling.

141 Response of tissue-specific DEGs reproduced across SAT and muscle, not in blood

142 Next, we questioned whether there was a transcription response common to the three tissues. Hereto, we examined the response of the eigengene of a set of tissue-specific DEGs 143 144 in the other tissues (see Materials and Methods). For example, the eigengene of the male-145 upregulated DEGs in SAT tissue was also positively upregulated in muscle, whereas it was non-146 significantly downregulated in blood (second row in Figure 3). Similarly, in females, we noted 147 correspondence between SAT and muscle, although the eigengene of the upregulated DEGs in SAT was stronger than in males, in blood the effect was less significant (sixth row, Figure 148 149 3). Since there was only one significant DEG in postprandial blood (Figure 1, Table 2), no blood 150 down- or upregulated DEGs eigengenes were calculated.

151 In general, we noted a similar behavior of the male/female eigengenes across the 152 tissues (Figure 3). Remarkably, the eigengene of SAT-downregulated DEGs was upregulated 153 in muscle (in both male and female) and the eigengene of muscle-upregulated DEGs was 154 downregulated in SAT (although not significantly). Furthermore, the effect of the muscle and 155 SAT eigengenes of DEGs were not significantly changed in blood, suggesting that these 156 SAT/muscle genes were not affected by the intervention in blood. In summary, when 157 investigating the intervention effect of the genes strongest affected by the intervention, no 158 overlap of transcriptional response was observed between blood and SAT or muscle. 159 Relationships did exist within the transcriptional response between SAT and muscle, although 160 not always consistent in direction of effect.

161 *Especially SAT- and muscle-specific DEGs associate with health markers*

To better understand the differential expressed transcriptome, we calculated the association 162 163 between the eigengene values and the change in ten clinical metabolic health markers: /) two DEXA scan body composition measurements: whole-body fat%, trunk fat%; II) seven 164 traditional metabolic health markers: BMI, Waist Circumference (WC) fasting insulin, fasting 165 HDL cholesterol, fasting HDL cholesterol size, SBP, DBP; and III) the Framingham Risk Score 166 167 (FRS), which is a cardiovascular risk score (Figure 4). Since the intervention effects on some of the health markers differed between the subgroups (Table 1), which thus can cause 168 169 differences in effects due to differences in subsets, the relation between the expression data 170 and metabolic health markers was performed on the overlapping blood, SAT and muscle 171 samples (57; 33 males, 24 females). In blood, there were no significant associations between any of the eigengenes and classical metabolic health markers. The SAT eigengenes were 172 173 significantly associated to the change in DEXA body composition parameters (whole-body 174 fat%, trunk fat%), four of the seven traditional metabolic health markers (BMI, WC and fasting HDL cholesterol size) in the SAT transcriptome of both sexes, and with the fasting HDL 175 176 cholesterol level and FRS only in men. Interestingly, also the muscle eigengenes expressed in 177 SAT were significantly associated to the change in all of these parameters in men, except for fasting HDL cholesterol level, with the downregulated muscle eigengenes showing the 178 179 strongest associations overall. In the transcriptome of muscle, the downregulated SAT 180 eigengene associated negatively to DBP and the downregulated muscle eigengene associated to the change in DEXA body composition markers; whole-body fat%, trunk fat%, and 181 182 metabolic health markers; BMI in men, BMI and WC in women.

Taken together, especially in the SAT transcriptome, the SAT and muscle eigengenes
reflected changes in DEXA body composition markers (whole-body and trunk fat%) and

185 classical metabolic health markers (BMI, WC, fasting HDL cholesterol level and size), and to

186 the former four also did the downregulated muscle eigengene in the muscle transcriptome.

187 Many effects were similar in males and females, although more pronounced in males.

188 Integrated analysis reveals male postprandial blood, SAT and muscle transcriptome

189 share a joint variation

190 To explore the tissue dependency in a more intricate way, we captured the joint transcriptional reaction to the intervention across the three different tissues using Joint and 191 192 Individual Variation Explained (JIVE) analysis (Materials and Methods, Supplementary Figure 193 1). The joint representation is a linear subspace spanned by the genes, analogous to a 194 principal component analysis (PCA), but that now also explains the joint variation in 195 expression in all three tissues across the intervention (Materials and Methods). Hence, although informed by all three tissues, the blood expression of a participant (before and after 196 197 the intervention) can be mapped to this joint subspace without needing information about the muscle or SAT transcriptome. In other words, the blood transcriptome of a participant is 198 199 decomposed in three (independent) transcriptome terms: 1) a term that reflects the joint transcriptomic variation in blood, SAT and muscle, 2) a term reflecting the remaining 200 201 structured variation in blood only (also referred to as the individual variation), and 3) a term 202 that represents the residual noise in the blood transcriptome, i.e. the signal that is not 203 explained by the joint and individual terms. Subsequently, we can measure the differential 204 expression across the intervention in either the joint subspace (of blood, SAT and muscle) or 205 the individual subspaces (for each tissue types). The number of joint and individual components are determined through permutation testing (Materials and Methods). 206

In the male samples, JIVE identified 2 joint components for the shared transcriptomic
variation in blood, SAT and muscle, and 11, 15 and 15 individual components for the blood,

SAT and muscle, respectively (Supplementary Table 8). In postprandial blood and SAT, the joint components together explained 8% and 7.2% of the total variation in those tissues, respectively. In the postprandial muscle transcriptome, both joint components explained 31.3% of the variation. In female samples, JIVE did not identify a joint term that was independent from the individual terms (Supplementary Figure 2).

- 214 In male samples, the drivers of the joint components were involved in phospholipid
- 215 metabolism and the immune system

216 To investigate what genes expressed in postprandial blood were the drivers of the joint effect, 217 the 250 genes that had the largest loading factors for each of the two joint components were 218 selected, respectively (Materials and Methods). Both sets of genes were subsequently used 219 in a pathway enrichment analysis. In postprandial blood, the genes driving the first joint 220 component (JC1) that explains most of the joint variance across the tissues were involved 221 phospholipid metabolism, while the genes driving the second joint component (JC2) were 222 involved in interferon signaling, inflammation and other aspects of the immune system 223 (Figure 5).

224 Phospholipid metabolism genes that drive the joint effect also capture the intervention

225 effect individually

Next we wondered whether the genes driving the significant effects of JC1 and JC2 also showed an intervention effect by themselves. Within the both sets of 250 genes that had the highest loading factors for each of the joint components, we selected those genes that overlapped with the found enriched pathways. Interestingly, we found a remarkable difference between the intervention effect of the genes involved in phospholipid metabolism and those involved in immunity (Figure 6): the joint effect of the genes involved in

phospholipid metabolism was more strongly influenced by the intervention than the individual effect, however, neither effect crossed the significant threshold after adjustment for multiple testing (Figure 6). Compared to the phospholipid metabolism genes, the effect sizes in both the joint nor individual effect of genes involved in the immune system were considerably weaker. These effects were also non-significantly associated with the intervention effect.

238 Immune genes that relate to the joint effect capture metabolic health effects of the

239 intervention

240 Next we investigated whether changes in the genes involved in phospholipid metabolism 241 and immunity associated with changes in the metabolic health markers. Here we found that 242 neither the joint nor individual effect of the phospholipid genes associated to these health 243 marker effects (Figure 6). In contrast, the joint effect of the genes involved in the immune 244 system did strongly associate with markers of body composition (whole-body and trunk 245 fat%) while, except for RAB44 and RAPGEF3 (which was associated to the fasting HDL 246 cholesterol level), the individual effect did not. Hence, it is possible to detect effects of the 247 intervention in the postprandial blood transcriptome, but only when the individual blood 248 effect has been removed.

249 Variance in the muscle and SAT transcriptome explain the selection of contributing

250 genes to the joint components of blood differently

To investigate which of the two tissues, SAT or muscle, correlated most with the joint components in blood, we calculated the correlations of the top 250 genes with the highest loading factors for each of the joint components (JC1 and JC2) in blood with those in muscle and SAT (Supplementary Figure 3). The JC1 blood genes showed increasingly higher absolute

correlations with the JC1 and JC2 muscle genes, whereas the JC2 blood genes showed an opposite behavior and were merely not correlated with the JC1 or JC2 muscle genes. On the other hand, the JC1 and JC2 SAT genes showed a more strong absolute correlation with the JC2 blood genes as opposed to the JC1 blood genes. These results indicate that the JC1 blood genes (dominated by genes involved in phospholipid metabolism) are largely correlated with the variance in the muscle transcriptome, the JC2 blood genes (dominated by genes involved in the immune system) are more correlated with the variances in the SAT transcriptome.

262 **Discussion**

263 We investigated the effects in men and women of a mild combined lifestyle intervention in 264 older adults on the postprandial transcriptome of blood, subcutaneous adipose tissue (SAT) 265 and muscle tissue, and related the response to changes in parameters of metabolic health. 266 Furthermore, using JIVE, we explored the shared variation between the three transcriptomes. 267 In both sexes, the 13-week combined lifestyle intervention had virtually no effect on the 268 postprandial blood transcriptome, while the postprandial SAT and muscle transcriptomes 269 responded significantly. In postprandial SAT, pathways involved in HDL remodeling, cell 270 signaling and CO2/O2 exchange (both sexes), ligand binding by scavenger receptors, 271 NR1H3/NR1H2 cholesterol related gene expression and fertilization (male only), and signal 272 transduction and MAPK signaling cascades (female only) were overrepresented, while in the 273 postprandial muscle, pathways involved in collagen, extracellular matrix remodeling, protein O-glycolysation (male only) and osteoblast differentiation (female only) were significantly 274 275 overexpressed.

The response of the SAT transcriptome in both sexes associated most strongly with changes in the Framing Risk Score (FRS), and metabolic health, including HDL cholesterol level

and size, and specifically in body composition markers (whole-body and trunk fat%, BMI and 278 279 Waist Circumference (WC)), and so did a part of the changes in the muscle transcriptome (whole-body and trunk fat%, BMI, WC and diastolic blood pressure (DBP)). We did observe 280 sex differences in the association effect sizes. Lastly, we identified shared variation between 281 282 the three transcriptomes, but only in the tissues of men. We identified in the blood transcriptome a joint intervention effect dominated by genes involved in phospholipid 283 metabolism, largely influenced by genes overlapping with changes in the muscle 284 285 transcriptome, and a second joint intervention effect dominated by genes involved in the immune system, overlapping with changes in the SAT transcriptome. The latter associated 286 287 with changes in body composition (whole-body and trunk fat%). These effects were not 288 detected in the original transcriptome of postprandial blood.

The effect of lifestyle interventions on the human transcriptome has mainly been 289 290 reported for dietary and physical activity regimes separately, typically in metabolically unhealthy or frail participants ^{22,24–26,30}. This makes a direct comparison with the response of 291 292 the healthy, non-diabetic, older adults ⁶ to the GOTO combined lifestyle intervention difficult. 293 Nevertheless, we do find similarities to dietary interventions. Like in GOTO, nutritional 294 interventions with a mild energy restriction (20-25%, for 4-12 weeks) hardly affected the 295 blood transcriptome ²⁴. In adipose tissue, caloric restriction interventions changed the 296 expression of genes involved in glucose and lipid metabolism, signal transduction, stem cell maintenance and vascularization ^{18,21,25,26,30}, the former three of which were also 297 298 overrepresented in the SAT transcriptome of the GOTO study (Figure 2). Lastly, in overweight 299 or obese participants of a weight-loss intervention study (6 months, diet, physical activity 300 separate and combined), genes involved in signaling pathways, translation, cell life span and ATP binding were significantly overrepresented ³¹, part of which we also identified in GOTO. 301

302 The effects of the muscle transcriptome to physical activity changes were reported in both 303 anabolic and catabolic interventions, the latter being more similar to the regimen in GOTO. In line with the results found in GOTO, a systematic review investigating the effects of 21 activity 304 305 interventions on the ECM in muscle, found that both types of activity interventions had a 306 positive effect on collagen level and identified ECM adaptations related to muscle remodeling in both young and older participants ²⁷. Not all gene expression effects are common to all 307 308 interventions. A 6-months aerobic, resistance, or aerobic + resistance training regimen in 309 obese older adults changed expression of genes involved in myogenesis, autophagy, mitochondrial biogenesis and inflammation, in all three training regimens ²³, which was not 310 311 observed for the muscle transcriptome in the GOTO study. This difference may be ascribed to 312 the better health at baseline of GOTO individuals and/or the duration and intensity of this in 313 essence anabolic intervention. Another anabolic 6-month resistance training intervention in 314 frail and healthy older adults, showed effects similar to GOTO; an increase in expression levels of collagen and extracellular matrix related genes ²². Finally, in mice a combined caloric 315 316 restriction and physical activity intervention reported DEGs involved in MAPK signaling, 317 metabolic processes, cholesterol metabolism, extracellular matrix organization ³⁶, which we also found in the GOTO study (Figure 2). 318

There is a number of other differences in study designs as compared to reported interventions. GOTO was a personalized intervention, to mimic a realistic situation when people change their lifestyle. Consequently, there was a considerable variation in the instructed lifestyle change between the participants; some changed their diet more, while other focused more in a change in physical activity. Furthermore the transcriptome in GOTO was measured in tissues taken after a standardized meal ¹⁷, which is usually not done.

325 A direct association between the expression levels in adipose tissue and muscle, and 326 classical metabolic health markers is generally not looked into, making it difficult to relate the 327 effects identified in GOTO (Figure 4). However, one cohort study of BMI discordant monozygotic twins ($\Delta BMI \ge 2.5 \text{ kg/m2}$) reported on adipose tissue and skeletal muscle 328 329 transcriptome differences ⁴¹. This study identified DEGs involved in ECM remodeling, 330 DNA/RNA metabolism, mitochondrial metabolism (both adipose tissue and skeletal muscle), 331 fatty acid metabolism, MAPK and AMPK signaling (adipose tissue only), and amino acid 332 metabolism (skeletal muscle only) ⁴¹, which largely overlap with the overrepresented 333 pathways in GOTO (Figure 2). Additionally, it was emphasized that combined differential gene 334 expression scores of the adipose tissue transcriptome were stronger associated with classical 335 metabolic health markers (including BMI, whole-body fat%, HDL cholesterol) than those of muscle transcriptome ⁴¹, which we also observed in GOTO (Figure 4). 336

When investigating the effects of the DEGs of one tissue in the other two tissues, we identified that the intervention had an effect on the SAT DEGs in the muscle transcriptome, but not the other way around (Figure 3). This indicated that the muscle transcriptome captured some of the effects on the SAT transcriptome, possibly by a SAT component in the muscle tissue.

Using the JIVE data integration approach ⁴⁰, we have identified a shared variation between the three tissues in the male samples represented by two joint components (Supplementary Figures 1 & 2, Supplementary Table 8). The genes that drove these two joint components were involved in phospholipid metabolism and the immune system, respectively (Figure 5). Like the eigengene effects in the postprandial muscle transcriptome (Figure 3 and 4), the genes driving the first joint component (JC1) in blood showed a strong intervention effect but did not capture metabolic health effects (Figure 6). Moreover, these genes were also more

349 correlated to variances in the muscle transcriptome than the SAT transcriptome 350 (Supplementary Figure 3). Conversely, similarly to the effects in postprandial SAT, the genes 351 driving the second joint component (JC2) in blood strongly captured the metabolic health 352 effects of the intervention but had a weaker association to the intervention itself. This was 353 also confirmed by a higher correlation with variances in the SAT transcriptome than the 354 muscle transcriptome of the genes belonging to this second joint blood component 355 (Supplementary Figure 3). From these results, we postulated that JC1 in blood was largely 356 driven by the muscle transcriptome, while JC2 in blood was primarily driven by the SAT transcriptome. 357

358 This study faced several limitations, which need to be considered when interpreting the 359 results. Firstly the number of participants with transcriptomics measurements of postprandial blood (88; 45 males, 24 females), postprandial SAT (78; 38 males, 40 females) and 360 361 postprandial muscle (82; 48 males, 34 females), is relatively low, compared to the complete 362 number of participants of the GOTO study (153; 75 males, 78 females). The variation in the 363 instructed lifestyle changes in combination with the low number of samples and slight 364 differences at baseline (Supplementary Table 1), could also have led to a large variation in outcomes when comparing participants with postprandial blood, SAT and muscle 365 transcriptome measurements (Table 1). To circumvent part of this variation, the effects 366 367 between the transcriptome and metabolic health markers were only studied within the 368 participants with transcriptomic measurements in postprandial blood, SAT and muscle (57; 369 33 males, 24 females). Secondly, the group of participants with transcriptomic measurements 370 of all three tissues was also used for the JIVE analysis. In females, the JIVE model did not 371 identify a shared variation, this could be because of the shared effects between the tissues 372 were too weak, which was likely attenuated by the low number of participants (24). In males,

the JIVE analysis did identify two joint components between the transcriptomes of the threetissues.

375 Overall, we were able to identify considerable effects of a 13-week moderate nutritional 376 and physical activity intervention on the transcriptome of SAT and muscle of healthy older 377 adults. We demonstrated that the changes in expression of genes in metabolism pathways in SAT, and collagen and muscle remodeling pathways in muscle associated with beneficial 378 379 features of overall metabolic health. These health effects were best captured by the 380 postprandial SAT transcriptome. We identified a joint space of three tissues in postprandial 381 blood, which was able to capture some of the effects in the other two tissues, even though 382 postprandial blood alone did not capture this. We propose that this method could be used to 383 estimate part of the effect of an intervention on SAT and muscle transcriptomes, when only 384 blood expression data is available.

385 Materials and Methods

386 *Study Design*

387 The Growing Old tOgether (GOTO) study was a 13-week lifestyle intervention study with 164 388 healthy older adult participants (83 males, 81 females, mean age 62.9 +- 5.7, mean BMI 26.9 389 +- 2.5). The participants underwent a 25% reduction in energy expenditure, equally divided over decreased nutrient intake and increased physical activity, as described in ⁶. Both before 390 391 and after the intervention, the participants were subjected to a nutrient challenge, consisting 392 of an overnight fast and an intake of standardized nutrient shake, as described in ¹⁷. Thirty 393 minutes after the nutrient challenge, samples of postprandial whole blood, postprandial 394 subcutaneous adipose tissue (SAT) (subcutaneous abdominal fat pad) and postprandial 395 muscle (vastus vascularis) were taken. Out of 164 participants, 153 (75 men, 78 women) were

compliant to the intervention according to thresholds described in ¹². Out of 153 compliant
participants, postprandial transcriptomics were measured in three tissues: 88 (45 males, 43
females) in blood, 78 (38 males, 40 females) in SAT, 82 (48 males, 34 females) in muscle. Fiftyseven participants (33 males, 24 females) had postprandial transcriptomic measurements in
all three tissues.

401 Diagnostic Measurements

All measurements were performed in fasting serum collected through venipuncture. Insulin 402 403 was measured using a Immulite 2000 xPi (Siemens, Eschborn, Germany). HOMA2-IR was 404 calculated the publicly available HOMA calculator using (https://www.dtu.ox.ac.uk/homacalculator/)⁴². Fasting HDL cholesterol and HDL cholesterol 405 406 size were measured using the previously described Nightingale platform ⁴³. Complete methods of diagnostic measurements are described in ⁶. 407

408 DEXA measurements

409 Method for the DEXA measurements is described in ¹⁴. In short: whole-body DEXA

410 (Discovery A, Hologic Inc., Bedford, MA,USA) was used to measure eleven body composition

- 411 components, including whole-body fat mass and trunk fat mass. Whole-body fat% was
- 412 calculated by dividing whole-body fat mass by whole-body mass. Trunk fat% was calculated
- 413 by dividing the trunk fat mass by the total trunk weight.
- 414 Investigation of Health Marker Effects

The intervention effect of each health marker was calculated using a linear mixed model, adjusted for age (fixed effects) and individual (random effects). The effect for males and females were calculated separately. The significance was adjusted for multiple testing using the Bonferroni correction method.

419 RNA isolation and Sequencing

The method for RNA isolation and sequencing is described in ¹⁷. In short: libraries were 420 prepared using Illumina TruSeq version 2 library preparation kits. Data processing was 421 performed the in-house BIOPET Gentrap pipeline, described in ⁴⁴. The following steps were 422 part of the data processing: low quality trimming using sickle version 12.00. Cutadapt 423 424 version 1.1 was used to perform the adapter clipping. The reads were aligned to GRCh37, while masking for SNPs common in the Dutch population (GoNL ⁴⁵ MAF > 0.01), using STAR 425 426 version 2.3.0e. Picard version 2.4.1. was used to perform sam to bam conversion and 427 sorting. Read quantification was performed using htseq-count version 0.6.1.p1 using 428 Ensemble gene annotations version 86 for gene definitions. In postprandial blood, the 429 sequencing resulted in an average of 37.2 million reads per sample, 97% (+- 0.4%) of which 430 were mapped. In postprandial SAT, samples had an average of 11.4 million sequenced 431 reads, 95% (+- 1.6%) of which were mapped. In postprandial muscle, an average of 36.9 million sequence reads per sample, 98% (+- 0.4%) of which were mapped. 432 433 Differential Gene Expression

In all tissues, the normalizations and differential gene expression analysis were performed in
a sex stratified manner. Genes that had less than 2 counts on average, were removed from
the dataset. Tissue specific confounders were identified by associationing them with the
principal components that represent 90% of variation of TMM-CPM normalized gene
expression.

439 Postprandial Blood. There were 88 participants (45 male, 43 females) with RNA-seq
440 measurements pre -and post intervention. We adjusted for the following confounders: 1)
441 technical effects: median 5' bias, median 3' bias, and flowcell, mean insert size in female

442 samples only, 2) cell type percentages: eosinophiles, monocytes, lymphocytes, basophiles),

443 and 3) personal details (age at baseline, and person ID as a random effect).

444 *Postprandial SAT.* 78 participants (38 male, 40 female) had RNA-seq measurements pre-and
445 post-intervention. We adjusted for the following confounders: 1) technical effects: molarity
446 of sample, isolation date, and plate number in female samples only; and 2) personal
447 measures: age at baseline, and person ID as a random effect.

448 Postprandial Muscle. There were 82 participants (48 male, 34 female) with RNA-seq 449 measurements pre- and post-intervention. In male samples we adjusted for the following 450 confouders: 1) technical effects: total yield, flow cell, isolations series, 28S/18S ratio, plate 451 number, mean insert size, total number of aligned bases, RNA integrity number, median 5' 452 bias, median 3' bias; and 2) personal details: age at baseline, and person ID as a random effect. 453 In female samples, we adjusted for the following confounders: 1) technical effects: RNA 454 260/280 ratio, 28S/18S ratio, biopsy number, hand grip strength, RNA integrity number, mean 455 insert size, isolation series, sequence % that passed quality control, total yield, total number 456 of aligned bases, median 3' bias; and 2) personal details: age at baseline, and person ID as a random effect. 457

The differential gene expression analysis was performed using linear mixed models analysis (limma version 3.52.2) in combination with VOOM normalization ⁴⁶. We adjusted for multiple testing using the false discovery rate. An fdr-adjusted p-value < 0.01 was considered significant. Genes with an absolute log2 fold change \geq 0.25 were considered differentially expressed.

463 Functional Enrichment

464 The pathway overrepresentation analysis was performed using the enricher function of 465 clusterProfiler (version 4.4.4) ⁴⁷. Up-and downregulated significant DEGs were used as an

input, all expressed genes that passed the quality control were used as a background. The
Reactome database (version 81) ⁴⁸ was used for the functional enrichment. The tests were
adjusted for multiple testing using the Bonferroni correction method. An adjusted p-value
below 0.05 was considered significant.

470 *Eigengene Analysis*

471 The input for the eigengene calculation were the up- or downregulated significant DEGs. The eigengenes were calculated using the prcomp function of R package stats (version 4.2.1)⁴⁹, 472 473 the first principal component was considered the eigengene. Eigengene values were scaled 474 to mean 0 and standard deviation (sd) 1. The associations between the intervention (0 for 475 baseline, 1 for post intervention) and the eigengenes was calculated with logistic regression using the function *qlmer* of package *qlmnet* (version 4.1-6) ⁵⁰, adjusting for the same 476 477 covariates as in the differential gene expression analysis. The significance level of the association was adjusted for multiple testing, using the false discovery rate. Similarly, the 478 479 association between eigengenes and metabolic health markers are calculated. Metabolic 480 health marker measurements were scaled to mean 0 and sd 1. Outliers more than 3 sd values 481 from the mean were removed. Only participants with transcriptome measurements of all 482 three tissues were used in the health marker association analysis.

483 Joint and Individual Variation Explained (JIVE) Analysis

The Joint and Individual Variation Explained (JIVE) was calculated using the R package *r.jive* (version 2.4) ^{40,51}. Analysis was performed on all genes that passed the quality control across 57 participants (33 Male, 24 Female). Prior to JIVE, the gene expression levels were scaled to a mean of 0 and a sd of 1. For each tissue matrix $X^t \in \mathbb{R}^{p_t \times n}$ (where *t* represents the tissue, p_t represents the number of genes measured for tissue *t*, and *n* represents the number of

samples), the JIVE analysis linearly decomposes the input matrix (X^{t}) into three matrices 489 representing: 1) the shared data across the tissue: $J^t \in \mathbb{R}^{p_t \times n}$ explaining the joint variation 490 across the three tissue types, 2) a data matrix representing the variation unique to a tissue: 491 $A^t \in \mathbb{R}^{p_t \times n}$, and 3) a data matrix representing the residual noise, i.e the remaining 492 $R^t \in \mathbb{R}^{p_t \times n}$. tissue: with: $X^t = I^t + A^t +$ unexplained variation in а 493 R^{t} (Supplementary Figure 1). I^{t} is further decomposed into sub-matrices $I^{t} = U^{t}S$, 494 where $U^t \in \mathbb{R}^{p_t x r}$ represents a matrix containing the p_t gene loading vectors for the *r* joint 495 496 components, and S is a sample scoring matrix of joint effects across all tissues (S is the same for all tissues). The number of joint components, r, is selected using permutation testing as 497 described in ⁴⁰. Likewise, A^t is decomposed, and for each tissue, $A^t = W^t S^t$, where $W^t \in$ 498 $\mathbb{R}^{p_t x r_t}$ represents a matrix containing the p_t gene loading vectors for the r_t individual tissue 499 components (selected by permutation testing), and $S^t \in \mathbb{R}^{r_t \times n}$ is a sample scoring matrix of 500 individual components. For each tissue, we can identify the genes that contribute most to the 501 joint and individual effects based on the ranking of genes by each column of U^t and W^t , 502 503 respectively.

504 Data Availability Statement

505 The data that support the findings of this study are accessible upon request to the 506 corresponding author. The data are not publicly available due to privacy or ethical 507 restrictions.

508 Conflict of Interest Statement

The authors have stated explicitly that there are no conflicts of interest in connection withthis article.

511 Author Contributions

- 512 P.E. Slagboom and C.P.G.M. de Groot designed the study. M. Beekman collected and
- 513 curated the health data. N. Lakenberg and E. Suchiman generated the transcriptome data.
- 514 F.A. Bogaards T. Gehrmann, M.J.T. Reinders and P.E. Slagboom designed the data analysis
- approach, F.A. Bogaards, performed data analysis and F.A. Bogaards, T. Gehrmann, M.J.T.
- 516 Reinders, C.P.G.M. de Groot and P.E. Slagboom performed the research and interpreted the
- 517 data. All authors were involved in drafting and revising the manuscript.

518 Ethics Statement

- 519 The Medical Ethical Committee of the Leiden University Medical Center approved the study
- 520 (P11.187) and all participants signed a written informed consent. All experiments were
- 521 performed in accordance with relevant and approved guidelines and regulations. This trial
- 522 was registered at the Dutch Trial Register (http://www.trialregister.nl) as NTR3499.

523 Funding and Acknowledgements

- 524 This work was funded by the Horizon 2020 ERC Advanced grant: GEROPROTECT, the
- 525 Netherlands Consortium for Healthy Ageing (NWO grant 050-060-810), the framework of
- 526 the BBMRI Metabolomics Consortium funded by BBMRI-NL (NWO 184.021.007 and
- 527 184.033.111), ZonMw Project VOILA (ZonMW 457001001) and X-omics (NWO 184.034.019).
- 528 The funding agencies had no role in the design and conduct of the study; collection,
- 529 management, analysis, and interpretation of the data; and preparation, review, or approval
- of the manuscript. The authors would like to express their gratitude to all participants of the
- 531 GOTO study who did their very best to adhere to the intervention guidelines and underwent
- all measurements.

533 References

- Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of ageing.
 Nature 561, 45–56 (2018).
- 536 2. Sander, M. *et al.* The challenges of human population ageing. *Age Ageing* **44**, 185–
- 537 187 (2015).
- 538 3. Finkel, T. The metabolic regulation of aging. *Nat. Med. 2015 2112* 21, 1416–1423
 539 (2015).
- 540 4. Salvioli, S. *et al.* Immune System, Cell Senescence, Aging and Longevity Inflamm-
- 541 Aging Reappraised. *Curr. Pharm. Des.* **19**, 1675–1679 (2013).
- 542 5. Ravussin, E. *et al.* A 2-year randomized controlled trial of human caloric restriction:
- 543 Feasibility and effects on predictors of health span and longevity. Journals Gerontol. -
- 544 Ser. A Biol. Sci. Med. Sci. **70**, 1097–1104 (2015).
- 545 6. van de Rest, O. *et al.* Metabolic effects of a 13-weeks lifestyle intervention in older
- 546 adults: The Growing Old Together Study. *Aging (Albany. NY).* **8**, 111–126 (2016).
- 547 7. Marseglia, A. *et al.* Effect of the NU-AGE diet on cognitive functioning in older adults:
- 548 A randomized controlled trial. *Front. Physiol.* **9**, (2018).
- 549 8. Fitzgerald, K. N. et al. Potential reversal of epigenetic age using a diet and lifestyle
- 550 intervention: a pilot randomized clinical trial. *Aging (Albany NY)* **13**, 9419 (2021).
- 551 9. Villareal, D. T. *et al.* Effect of lifestyle intervention on metabolic coronary heart
- 552 disease risk factors in obese older adults. *Am. J. Clin. Nutr.* 84, 1317–1323 (2006).
- 553 10. Campbell, K. L., Landells, C. E., Fan, J. & Brenner, D. R. A Systematic Review of the
- 554 Effect of Lifestyle Interventions on Adipose Tissue Gene Expression: Implications for
- 555 Carcinogenesis. *Obesity* **25**, S40–S51 (2017).

- 556 11. Das, S. K. et al. Body-composition changes in the Comprehensive Assessment of Long-
- 557 term Effects of Reducing Intake of Energy (CALERIE)-2 study: A 2-y randomized
- 558 controlled trial of calorie restriction in nonobese humans. Am. J. Clin. Nutr. 105, 913–
- 559 927 (2017).
- 560 12. Bogaards, F. A. *et al.* PLIS: A metabolomic response monitor to a lifestyle intervention
 561 study in older adults. *FASEB J.* **36**, e22578 (2022).
- 562 13. Almanza-Aguilera, E. *et al.* Impact in Plasma Metabolome as Effect of Lifestyle
- 563 Intervention for Weight-Loss Reveals Metabolic Benefits in Metabolically Healthy
- 564 Obese Women. J. Proteome Res. 17, 2600–2610 (2018).
- 565 14. Beekman, M. *et al.* Lifestyle-Intervention-Induced Reduction of Abdominal Fat Is
- 566 Reflected by a Decreased Circulating Glycerol Level and an Increased HDL Diameter.
 567 *Mol. Nutr. Food Res.* 64, (2020).
- 568 15. Diener, C. *et al.* Baseline Gut Metagenomic Functional Gene Signature Associated
- 569with Variable Weight Loss Responses following a Healthy Lifestyle Intervention in
- 570 Humans. *mSystems* **6**, (2021).
- 571 16. Franzke, B. *et al.* The plasma proteome is favorably modified by a high protein diet
- 572 but not by additional resistance training in older adults: A 17-week randomized
- 573 controlled trial. *Front. Nutr.* **9**, (2022).
- 574 17. Gehrmann, T. et al. A combined lifestyle intervention induces a sensitization of the
- 575 blood transcriptomic response to a nutrient challenge. *bioRxiv* 2021.05.18.444591
- 576 (2021) doi:10.1101/2021.05.18.444591.
- 577 18. Swindell, W. R., List, E. O., Berryman, D. E. & Kopchick, J. J. Transcriptional profiling
- 578 identifies strain-specific effects of caloric restriction and opposite responses in human
- and mouse white adipose tissue. *Aging (Albany. NY).* **10**, 701–746 (2018).

- 580 19. Rubenstein, A. B. *et al.* Skeletal muscle transcriptome response to a bout of
- 581 endurance exercise in physically active and sedentary older adults. Am. J. Physiol. -
- 582 *Endocrinol. Metab.* **322**, E260–E277 (2022).
- 583 20. van Bussel, I. P. G., Fazelzadeh, P., Frost, G. S., Rundle, M. & Afman, L. A. Measuring
- 584 phenotypic flexibility by transcriptome time-course analyses during challenge tests
- 585 before and after energy restriction. *FASEB J.* **33**, 10280–10290 (2019).
- 586 21. Capel, F. *et al.* Macrophages and adipocytes in human obesity: adipose tissue gene
- 587 expression and insulin sensitivity during calorie restriction and weight stabilization.
- 588 Diabetes 58, 1558–1567 (2009).
- 589 22. Hangelbroek, R. W. J. et al. Expression of protocadherin gamma in skeletal muscle
- tissue is associated with age and muscle weakness. *J. Cachexia. Sarcopenia Muscle* 7,
 604–614 (2016).
- 592 23. Colleluori, G. *et al.* Aerobic Plus Resistance Exercise in Obese Older Adults Improves
- 593 Muscle Protein Synthesis and Preserves Myocellular Quality Despite Weight Loss. *Cell*

594 *Metab.* **30**, 261-273.e6 (2019).

- 595 24. Day, K. *et al.* Transcriptomic changes in peripheral blood mononuclear cells with
- 596 weight loss: systematic literature review and primary data synthesis. (2021)
- 597 doi:10.1186/s12263-021-00692-6.
- 598 25. Wahl, D. & LaRocca, T. J. Transcriptomic Effects of Healthspan-Promoting Dietary
- 599 Interventions: Current Evidence and Future Directions. *Front. Nutr.* **8**, 515 (2021).
- 600 26. Fachim, H. A. *et al.* Lifestyle intervention in individuals with impaired glucose
- 601 regulation affects Caveolin-1 expression and DNA methylation. *Adipocyte* **9**, 96–107
- 602 (2020).
- 603 27. Kritikaki, E. et al. Exercise training-induced extracellular matrix protein adaptation in

604 locomotor muscles: A systematic review. *Cells* **10**, (2021).

- 605 28. Matsuo, Y. et al. Impaired HDL function in obese adolescents: Impact of lifestyle
- 606 intervention and bariatric surgery. *Obesity* **21**, E687–E695 (2013).
- 607 29. Ellsworth, D. L. *et al.* Intensive cardiovascular risk reduction induces sustainable
- 608 changes in expression of genes and pathways important to vascular function. *Circ.*
- 609 *Cardiovasc. Genet.* **7**, 151–160 (2014).
- 610 30. Armenise, C. et al. Transcriptome profiling from adipose tissue during a low-calorie
- 611 diet reveals predictors of weight and glycemic outcomes in obese, nondiabetic
- 612 subjects. Am. J. Clin. Nutr. **106**, 736–746 (2017).
- 613 31. Campbell, K. L. et al. Gene expression changes in adipose tissue with diet-and/or

614 exercise-induced weight loss. *Cancer Prev. Res.* 6, 217–231 (2013).

- 615 32. Vroege, D. P. et al. Dose-response effects of a Web-based physical activity program
- on body composition and metabolic health in inactive older adults: Additional
- 617 analyses of a randomized controlled trial. J. Med. Internet Res. **16**, (2014).
- 618 33. Rickman, A. D. *et al.* The CALERIE Study: Design and methods of an innovative 25%
- 619 caloric restriction intervention. *Contemp. Clin. Trials* **32**, 874–881 (2011).
- 620 34. Verberne, L. D. M. et al. Evaluation of a combined lifestyle intervention for
- 621 overweight and obese patients in primary health care: a quasi-experimental design.
- 622 *Fam. Pract.* **33**, 671–677 (2016).
- 623 35. Fazelzadeh, P. et al. Weight loss moderately affects the mixed meal challenge
- 624 response of the plasma metabolome and transcriptome of peripheral blood
- 625 mononuclear cells in abdominally obese subjects. *Metabolomics* **14**, (2018).
- 626 36. Feng, Y. et al. Transcriptomics Dissection of Calorie Restriction and Exercise Training
- 627 in Brown Adipose Tissue and Skeletal Muscle. *Nutrients* **15**, 1047 (2023).

| 628 | 37. | Rowland, B. et al. Transcriptome-wide association study in UK Biobank Europeans |
|-----|-----|------------------------------------------------------------------------------------------------|
| 629 | | identifies associations with blood cell traits. Hum. Mol. Genet. 31 , 2333–2347 (2022). |
| 630 | 38. | Harris, S. E. et al. Age-related gene expression changes, and transcriptome wide |
| 631 | | association study of physical and cognitive aging traits, in the Lothian Birth Cohort |
| 632 | | 1936. Aging (Albany NY) 9 , 2489 (2017). |
| 633 | 39. | Reel, P. S., Reel, S., Pearson, E., Trucco, E. & Jefferson, E. Using machine learning |
| 634 | | approaches for multi-omics data analysis: A review. Biotechnol. Adv. 49, 107739 |
| 635 | | (2021). |
| 636 | 40. | Lock, E. F., Hoadley, K. A., Marron, J. S. & Nobel, A. B. Joint and individual variation |
| 637 | | explained (JIVE) for integrated analysis of multiple data types. Ann. Appl. Stat. 7, 523- |
| 638 | | 542 (2013). |
| 639 | 41. | van der Kolk, B. W. et al. Molecular pathways behind acquired obesity: Adipose tissue |
| 640 | | and skeletal muscle multiomics in monozygotic twin pairs discordant for BMI. (2021) |
| 641 | | doi:10.1016/j.xcrm.2021.100226. |
| 642 | 42. | Wallace, T. M., Levy, J. C. & Matthews, D. R. Use and abuse of HOMA modeling. |
| 643 | | Diabetes Care 27 , 1487–1495 (2004). |
| 644 | 43. | P, S. et al. High-throughput serum NMR metabonomics for cost-effective holistic |
| 645 | | studies on systemic metabolism. Analyst 134 , 1781–1785 (2009). |
| 646 | 44. | Zhernakova, D. V et al. Identification of context-dependent expression quantitative |
| 647 | | trait loci in whole blood. <i>Nat. Genet. 2016 491</i> 49 , 139–145 (2016). |
| 648 | 45. | Francioli, L. C. et al. Whole-genome sequence variation, population structure and |
| 649 | | demographic history of the Dutch population. Nat. Genet. 2014 468 46, 818–825 |
| 650 | | (2014). |
| 651 | 46. | Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. Voom: Precision weights unlock linear |

- model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, 29 (2014).
- 47. Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing
- biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).
- 655 48. Gillespie, M. et al. The reactome pathway knowledgebase 2022. Nucleic Acids Res. 50,
- 656 D687–D692 (2022).
- 657 49. R Core Team. R Core Team 2021 R: A language and environment for statistical
- 658 computing. R foundation for statistical computing. https://www.R-project.org/. R
- 659 Found. Stat. Comput. **2**, 2019 (2022).
- 660 50. Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear
- 661 Models via Coordinate Descent. (2009).
- 662 51. O'Connell, M. J. & Lock, E. F. R.JIVE for exploration of multi-source molecular data.
- 663 *Bioinformatics* **32**, 2877–2879 (2016).

Table 1: Effect of the 13-weeks lifestyle intervention on metabolic health marker measurements differs per subset selection.

665

| | | Complete GOTO dataset 153 (75 Male, 78 Female) | | Overlapping Blood, SAT and Muscle Samples 57 (33 Male 24 Female) | | Blood Postprandial Samples 88 (45 Male, 43 Female) | | SAT Postprandial Samples 78 (38 Male, 40 Female) | | Muscle Postprandial Samples 82 (48 Male, 34 Female) | |
|----------------------------------|--------|------------------------------------------------------|----------|---------------------------------------------------------------------------|----------|-------------------------------------------------------|----------|-----------------------------------------------------|----------|-----------------------------------------------------------|----------|
| Health Marker, mean (SD) | Sex | Delta (sd) | P-adjust | Delta (sd) | P-adjust | Delta (sd) | P-adjust | Delta (sd) | P-adjust | Delta (sd) | P-adjust |
| BMI, kg/m ² | Male | -1.09 (0.78) | 2.4e-18* | -1.02 (0.78) | 1.8e-07* | -1.15 (0.79) | 1.4e-11* | -1.10 (0.77) | 1.3e-09* | -1.18 (0.78) | 6.9e-13* |
| Waist Circumference | Male | -4.49 (5.26) | 1.8e-09* | -3.82 (4.97) | 1.1e-03* | -4.47 (5.29) | 1.1e-05* | -4.42 (5.10) | 4.8e-05* | -4.94 (5.41) | 8.7e-07* |
| Systolic Blood Pressure, mm Hg | Male | -3.25 (12.54) | 2.8e-01 | -3.80 (14.36) | 1.0e+00 | -2.52 (13.42) | 1.0e+00 | -3.91 (14.14) | 9.7e-01 | -3.05 (13.53) | 1.0e+00 |
| Diastolic Blood Pressure, mm Hg | Male | -0.75 (6.98) | 1.0e+00 | -3.04 (7.09) | 1.9e-01 | -2.27 (6.67) | 2.7e-01 | -3.09 (7.10) | 1.1e-01 | -2.32 (6.86) | 2.3e-01 |
| Fasting Insulin, mU/L | Male | -0.31 (3.30) | 1.0e+00 | 0.14 (3.63) | 1.0e+00 | 0.04 (3.62) | 1.0e+00 | -0.11 (3.71) | 1.0e+00 | -0.23 (3.72) | 1.0e+00 |
| Whole-Body Fat, % | Male | -1.60 (1.65) | 1.2e-09* | -1.17 (1.55) | 1.5e-03* | -1.43 (1.71) | 8.2e-06* | -0.92 (1.52) | 9.3e-05* | -1.77 (1.66) | 4.8e-07* |
| Trunk Fat, % | Male | -2.47 (2.13) | 3.2e-12* | -1.96 (1.81) | 3.3e-05* | -2.26 (2.13) | 2.3e-07* | -1.48 (1.74) | 1.1e-06* | -2.59 (2.04) | 8.7e-09* |
| Fasting HDL Cholesterol, mmol/L | Male | 0.04 (0.14) | 8.2e-02 | 0.04 (0.13) | 1.0e+00 | 0.06 (0.14) | 7.7e-02 | 0.05 (0.13) | 3.4e-01 | 0.07 (0.14) | 1.7e-02* |
| Fasting HDL Cholesterol size, nm | Male | 0.06 (0.09) | 2.0e-06* | 0.06 (0.08) | 9.9e-04* | 0.08 (0.08) | 2.0e-06* | 0.06 (0.09) | 4.6e-04* | 0.08 (0.09) | 1.3e-06* |
| FRS | Male | -0.40 (1.22) | 5.8e-02 | -0.61 (1.37) | 1.6e-01 | -0.53 (1.25) | 6.6e-02 | -0.58 (1.31) | 9.7e-02 | -0.54 (1.24) | 3.9e-02* |
| BMI, kg/m ² | Female | -1.23 (0.82) | 1.4e-20* | -1.26 (0.72) | 1.4e-07* | -1.27 (0.64) | 2.7e-15* | -1.35 (0.73) | 3.0e-13* | -1.36 (0.74) | 2.5e-11* |
| Waist Circumference | Female | -4.33 (5.57) | 1.4e-08* | -4.71 (5.74) | 5.4e-03* | -4.70 (5.30) | 7.5e-06* | -4.58 (5.31) | 3.0e-05* | -5.21 (5.32) | 2.3e-05* |
| Systolic Blood Pressure, mm Hg | Female | -3.01 (11.69) | 2.6e-01 | -1.67 (11.69) | 1.0e+00 | -2.85 (11.30) | 1.0e+00 | -2.46 (11.73) | 1.0e+00 | -1.35 (12.07) | 1.0e+00 |
| Diastolic Blood Pressure, mm Hg | Female | -1.25 (6.52) | 9.5e-01 | -1.19 (5.75) | 1.0e+00 | -1.66 (5.64) | 6.1e-01 | -2.29 (6.27) | 2.6e-01 | -1.43 (6.54) | 1.0e+00 |
| Fasting Insulin, mU/L | Female | -0.36 (3.25) | 1.0e+00 | -0.30 (2.35) | 1.0e+00 | -0.02 (3.01) | 1.0e+00 | -0.33 (3.63) | 1.0e+00 | -0.90 (2.84) | 7.3e-01 |
| Whole-Body Fat, % | Female | -1.57 (1.70) | 9.8e-09* | -1.33 (2.22) | 1.1e-02* | -1.64 (1.85) | 5.2e-05* | -1.58 (1.98) | 7.2e-05* | -1.98 (2.07) | 2.5e-04* |
| Trunk Fat, % | Female | -2.22 (2.15) | 3.1e-10* | -1.76 (2.78) | 7.1e-03* | -2.32 (2.36) | 1.2e-05* | -2.31 (2.48) | 2.6e-05* | -2.60 (2.57) | 1.1e-04* |
| Fasting HDL Cholesterol, mmol/L | Female | -0.04 (0.15) | 2.7e-01 | -0.03 (0.14) | 1.0e+00 | -0.05 (0.16) | 4.7e-01 | -0.05 (0.15) | 5.9e-01 | 0.00 (0.14) | 1.0e+00 |
| Fasting HDL Cholesterol size, nm | Female | 0.03 (0.08) | 1.7e-02* | 0.05 (0.09) | 1.7e-01 | 0.02 (0.08) | 6.0e-01 | 0.04 (0.08) | 2.6e-02* | 0.04 (0.09) | 9.1e-02 |
| FRS | Female | -0.65 (1.60) | 5.5e-03* | -0.46 (1.59) | 1.0e+00 | -0.42 (1.47) | 6.8e-01 | -0.62 (1.76) | 3.1e-01 | -0.68 (1.79) | 3.4e-01 |

Abbreviations: SD; standard deviation, SAT; subcutaneous adipose tissue, BMI; body mass index, FRS; Framingham Risk Score. Associations were calculated using a liner mixed model, adjusting

for age at baseline and person ID (as a random effect). P-values were adjusted using the Bonferroni correction method, an adjusted p-value < 0.05 was indicated with an *.





Figure 1: Intervention effect on postprandial blood, SAT and muscle gene expression levels in men and women. The log2 fold change is plotted on the x-axis, the -log10 FDR adjusted p-value is plotted on the y-axis. The dashed horizontal line represents the significance threshold of FDR 0.01, the two vertical lines represent the log2 fold change thresholds of -0.25 and 0.25. Genes with an absolute log2 fold change \geq 0.25 and an FDR adjusted p-value < 0.01 were considered significantly differentially expressed. Each dot represents a gene. Yellow dots represent the significantly differentially expressed genes (DEGs). Grey dots represent genes with an absolute log2 fold change < 0.25 and/or FDR \geq 0.01. Significantly differentially expressed genes with the strongest log2 fold changes were labeled with their gene symbol. Abbreviations: SAT; subcutaneous adipose tissue.

669

Table 2: The 13-weeks lifestyle intervention had the strongest effect on postprandial SAT and Muscle

671

| Sex | Tissue | State | Direction of Effect | Expressed Genes | Genes with Log2FC ≥ 0.25 | Genes with FDR < 0.01 | Genes with Log2FC ≥ 0.25 & FDR < 0.01 |
|--------|--------|--------------|------------------------|--------------------|-----------------------------|--------------------------|---------------------------------------------|
| Male | Blood | Postprandial | Downregulated | 8602 | 52 | 1 | 1 |
| Male | Blood | Postprandial | Upregulated | 10093 | 45 | 1 | 0 |
| Male | SAT | Postprandial | Downregulated | 10099 | 402 | 123 | 48 |
| Male | SAT | Postprandial | Upregulated | 9614 | 173 | 180 | 41 |
| Male | Muscle | Postprandial | Downregulated | 9413 | 228 | 139 | 46 |
| Male | Muscle | Postprandial | Upregulated | 7166 | 674 | 239 | 205 |
| Female | Blood | Postprandial | Downregulated | 9619 | 63 | 0 | 0 |
| Female | Blood | Postprandial | Upregulated | 9143 | 32 | 0 | 0 |
| Female | SAT | Postprandial | Downregulated | 9860 | 254 | 331 | 92 |
| Female | SAT | Postprandial | Upregulated | 9599 | 143 | 267 | 76 |
| Female | Muscle | Postprandial | Downregulated | 9464 | 292 | 127 | 46 |
| Female | Muscle | Postprandial | Upregulated | 7570 | 827 | 132 | 122 |

672 Abbreviations: SAT; subcutaneous adipose tissue

674

| | Male Samples | Female Samples | | | | | | |
|--------------------------------------------------------------------------------------------------|--------------|----------------|---------------|--|--|--|--|--|
| Signaling by Nuclear Receptors | * | ** | | | | | | |
| HDL remodeling | * | ** | | | | | | |
| Signal Transduction | | * | | | | | | |
| MAPK family signaling cascades | | * | | | | | | |
| O2/CO2 exchange in erythrocytes | | *** | Postprandial | | | | | |
| Erythrocytes take up carbon dioxide and release oxygen | | *** | Transcriptome | | | | | |
| Erythrocytes take up oxygen and release carbon dioxide NR1H3 & NR1H2 regulate gene expression | * | *** | | | | | | |
| linked to cholesterol transport and efflux Fertilization | ** | | | | | | | |
| Binding and Uptake of Ligands by Scavenger Receptors | ** | | | | | | | |
| Degradation of the extracellular matrix | *** | ** | | | | | | |
| Extracellular matrix organization | *** | *** | | | | | | |
| Collagen formation | *** | *** | | | | | | |
| Collagen biosynthesis and modifying enzymes | *** | *** | | | | | | |
| Assembly of collagen fibrils and other multimeric structures | *** | *** | | | | | | |
| Collagen chain trimerization | *** | *** | | | | | | |
| Netrin-1 signaling | | * | | | | | | |
| RUNX2 regulates osteoblast differentiation | | * | | | | | | |
| O-glycosylation of TSR domain-containing proteins | * | | | | | | | |
| NCAM signaling for neurite out-growth | *** | | Postprandial | | | | | |
| Defective B3GALTL causes PpS | * | | Transcriptome | | | | | |
| Collagen degradation | * | | | | | | | |
| Elastic fibre formation | ** | | | | | | | |
| Signaling by PDGF | *** | | | | | | | |
| Laminin interactions | * | | | | | | | |
| ECM proteoglycans | *** | | | | | | | |
| NCAM1 interactions | *** | | | | | | | |
| Scavenging by Class A Receptors | *** | | | | | | | |
| Crosslinking of collagen fibrils | *** | | | | | | | |
| Binding and Uptake of Ligands by Scavenger Receptors | *** | | | | | | | |
| Normalized Intervention Effect_1.0 -0.5 0.0 0.5 1.0 | | | | | | | | |

Figure 2: Significantly Overrepresented pathways in Postprandial SAT and Muscle Transcriptome. The

overrepresented pathways are plotted on the y-axis. Overrepresented pathways in the male samples are plotted on the left, overrepresented samples in female samples, on the right. The fill color represents the normalized direction of effect: dark blue (-1) all significantly differentially expressed genes in the pathway were underexpressed, dark red (1) all significantly differentially expressed genes in the pathway were overexpressed. P- values were adjusted for multiple testing using the Bonferroni correction method. Significance is indicated by the asterisks (* = p- adjust <0.05, ** = padjust <0.01, *** = p- adjust <0.001). Abbreviations: SAT; subcutaneous adipose tissue.



Figure 3: Effect of the 13-weeks lifestyle intervention on Eigengenes calculated using up- or downregulated significantly differentially expressed genes. Genes used for input of the eigengenes were plotted on the y-axis. Light blue squares represent male samples, dark red circles represent female samples. The estimated intervention effect is plotted on the x-axis. Brackets indicate the 95% confidence interval. The p-values were adjusted for multiple testing using the false discovery rate. Significance is indicated by the asterisks (* = p.adjust < 0.05; ** = p.adjust < 0.01, *** = *p.adjust < 0.001). Abbreviations: SAT; subcutaneous adipose tissue.*

676

678



Figure 4: The SAT postprandial transcriptome captures the metabolic health changes stronger than the blood and muscle transcriptome. Genes used for input of the eigengenes were plotted on the y-axis. The metabolic health markers were plotted on the x-axis. Colors indicate the direction of the effect: blue, a negative effect, white; no effect, red; a positive effect, the darker the color, the stronger the effect. The p-values were adjusted for multiple testing using the false discovery rate. Significance is indicated by the asterisks (* = p.adjust < 0.05; ** = p.adjust < 0.01, *** = 0.001). Abbreviations: SAT; subcutaneous adipose tissue.



681 Figure 5: Significantly Overrepresented pathways in the top 250 drivers of the postprandial blood joint effect. The overrepresented pathways are plotted on the y-axis. Y-axis is ordered by significance, strongest significantly overrepresenting pathways first. The x-axis represents -log10 of the Bonferroni adjusted significance p-value. 682



Figure 6: In postprandial blood, the joint component 1 drivers strongly captured the intervention effect while joint component 2 drivers was significantly associated to whole-body fat% and trunk fat% levels. The genes were plotted on the y-axis, the intervention effect and metabolic health markers were plotted on the x-axis. Colors indicate the direction of the effect: blue, a negative effect, white; no effect, red; a positive effect, the darker the color, the stronger the effect. The p-values were adjusted using the false discovery rate. Significance is indicated by the asterisks (* = p.adjust < 0.05).