

SUPPLEMENTARY MATERIAL

Supplementary table 1. Baseline characteristics by gender. HbA1c, blood glycosylated haemoglobin. Values are mean \pm SEM (n). P values are from unpaired t-test.

Characteristic	Males (n)	Females (n)	p value
Age (years)	58.3 \pm 2.7 (9)	63.6 \pm 2.8 (8)	0.20
BMI (Kg/m ²)	31.4 \pm 1.0 (9)	29.8 \pm 1.0 (8)	0.28
HbA1c (mmol/mol)	65.6 \pm 5.6 (9)	63.1 \pm 4.9 (8)	0.74
HbA1c (%)	8.1 \pm 0.5 (9)	7.9 \pm 0.4 (8)	0.71
Time since T2D diagnosed (years)	3.6 \pm 0.9 (5)	6.6 \pm 1.6 (7)	0.26
Dose of Metformin per day (mg)	1500.0 \pm 288.7(3)	1250 \pm 155.9(9)	0.45
Time of Metformin (years)	4.0\pm1.0 (5)	11.4\pm2.4 (8)	0.04
Number of steps	7114 \pm 984.4 (9)	6481 \pm 713.2 (9)	0.60

Supplementary table 2. Baseline characteristics by season. HbA1c, blood glycosylated haemoglobin. Values are mean \pm SEM (n). P values are from unpaired t-test.

Characteristic	Winter (n)	Summer (n)	p value
Age (years)	60.5 \pm 2.3 (10)	61.29 \pm 3.87 (7)	0.86
BMI (Kg/m ²)	30.4 \pm 0.9 (10)	31.0 \pm 1.3 (7)	0.69
HbA1c (mmol/mol)	70.7\pm4.9 (10)	55.6\pm3.5 (7)	0.04
HbA1c (%)	8.7\pm0.4 (10)	7.3\pm0.3 (7)	0.03
Time since T2D diagnosed (years)	4.6 \pm 2.2 (5)	5.8 \pm 1.0 (7)	0.59
Dose of Metformin per day (mg)	1042.0\pm100.3(6)	1583.0\pm200.7(6)	0.04
Time of Metformin (years)	10.3 \pm 3.5 (6)	7.0 \pm 1.6 (7)	0.38
Number of steps	6184 \pm 655.9 (11)	7761 \pm 1094 (7)	0.20

Supplementary table 3. Baseline characteristics by metformin timing during baseline period. HbA1c, blood glycosylated haemoglobin. Values are mean \pm SEM (n). P values are from unpaired t-test.


Baseline period			
Characteristic	Metformin before breakfast (n)	Metformin after breakfast (n)	p value
Age (years)	59.1 \pm 2.6 (9)	64.3 \pm 4.0 (4)	0.30
BMI (Kg/m ²)	30.6 \pm 1.3 (9)	30.8 \pm 1.1 (5)	0.91
HbA1c (mmol/mol)	56.6 \pm 4.6 (9)	68.8 \pm 6.6 (5)	0.26
HbA1c (%)	7.6 \pm 0.4 (9)	8.4 \pm 0.6 (5)	0.27
Time since T2D diagnosed (years)	8.4 \pm 2.5 (9)	10.7 \pm 0.9 (3)	0.64
Dose of Metformin per day (mg)	1918.0 \pm 511.3(8)	1333.0 \pm 333.3(3)	0.52
Time of Metformin (years)	6.4 \pm 1.2 (8)	3.7 \pm 2.7 (3)	0.29
Number of steps	7396 \pm 1025 (9)	6133 \pm 801.5 (5)	0.42

Supplementary table 4. Baseline characteristics by metformin timing in the morning exercise period. HbA1c, blood glycosylated haemoglobin. Values are mean \pm SEM (n). P values are from unpaired t-test.


Morning exercise			
Characteristic	Metformin before breakfast (n)	Metformin after breakfast (n)	p value
Age (years)	59.1 \pm 2.9 (8)	63.4 \pm 3.5 (5)	0.86
BMI (Kg/m ²)	30.8 \pm 1.4 (8)	30.8 \pm 1.1 (5)	0.99
HbA1c (mmol/mol)	56.9\pm4.3 (8)	75.2\pm4.7 (5)	0.01
HbA1c (%)	7.3\pm0.4 (6)	9.0\pm0.4 (5)	0.02
Time since T2D diagnosed (years)	5.6 \pm 1.0 (7)	4.7 \pm 3.7 (3)	0.74
Dose of Metformin per day (mg)	1357.0 \pm 179.8(7)	1167 \pm 166.7(3)	0.54
Time of Metformin (years)	10.3\pm3.5 (8)	7.0\pm1.6 (3)	0.02
Number of steps	7056 \pm 1096 (8)	6752 \pm 1141 (5)	0.86

Supplementary table 5. Baseline characteristics by metformin timing in the evening exercise period. HbA1c, blood glycosylated haemoglobin. Values are mean \pm SEM (n). P values are from unpaired t-test.


Evening exercise			
Characteristic	Metformin before breakfast (n)	Metformin after breakfast (n)	p value
Age (years)	60.9 \pm 1.8 (8)	61.0 \pm 8.7 (3)	0.98
BMI (Kg/m ²)	30.5 \pm 1.3 (8)	32.6 \pm 1.5 (3)	0.41
HbA1c (mmol/mol)	66.7 \pm 5.8 (8)	60.3 \pm 7.6 (3)	0.55
HbA1c (%)	8.2 \pm 0.5 (8)	7.7 \pm 0.7 (3)	0.55
Time since T2D diagnosed (years)	11.0 \pm 3.4 (6)	9.0 \pm 2.5 (3)	0.71
Dose of Metformin per day (mg)	1333.0 \pm 166.7(6)	1333.0 \pm 333.3(3)	0.99
Time of Metformin (years)	8.2\pm1.0 (8)	2.0\pm1.0 (3)	0.008
Number of steps	6264 \pm 758.3 (8)	5603 \pm 1119 (3)	0.65




Supplementary Figure 1. Number of exercise sessions completed during trial in the morning and evening exercise periods (n=18), blue lines are mean \pm SEM, the dotted line represents the number of sessions participants were asked to complete. Data were analysed using paired t-test.




Supplementary Figure 2. Total hours of sleep from Garmin devices at baseline (Garmin) and Munich Chronotype Questionnaires (MCTQ) (n=14), blue lines are mean±SEM, data were analysed using paired t-test. *= $p < 0.05$.




Supplementary Figure 3. Sleep architecture in the baseline, and morning and evening exercise periods of the trial. The values are shown (A) Deep sleep (n=15), (B) Light sleep (n=15), (C) REM sleep (n=15) and (D) Time awake (n=15), blue lines are mean±SEM. Data were analysed using one-way ANOVA followed up by Holm-Šidák's multiple comparisons test.



Supplementary figure 4. Mean area under the curve (AUC) glucose concentrations over 24-hours hourly during baseline, and morning and evening exercise split by sex. Glucose levels in Males weeks 1-2 (n=9), weeks 5-6 (n=8); Females weeks 1-2 (n=9), weeks 5-6 (n=8). Values are mean±SEM. Data were analysed using two-way mixed-model ANOVA followed up by Holm-Šídák's multiple comparisons test.



Supplementary Figure 5. Mean area under the curve (AUC) values over 24-hours hourly in baseline, and morning and evening exercise periods split by season. Winter weeks 1-2 (n=11), weeks 5-6 (n=10). Summer weeks 1-2 (n=7), weeks 5-6 (n=6). Values are mean±SEM. Data were analysed using two-way mixed-model ANOVA followed up by Holm-Šídák's multiple comparisons test.



Supplementary Figure 6. Meal and metformin timing. Mean area under the curve (AUC) glucose values for the first 24 hours of exercise day comparing Metformin pre and post dinner. (A) Baseline, metformin pre-dinner (n=10), metformin post-dinner (n=3); (B) Morning exercise, metformin pre-dinner (n=9), metformin post-dinner (n=4); (C) Evening exercise, metformin pre-dinner (n=7), metformin post-dinner (n=6). Values are mean \pm SEM. Data were analysed using unpaired t-test.