

Supplemental Table 6. Associations between aging constructs and mortality in cancer survivors stratified by sex, HRS (2016-2020)

Sample A^a: 946 cancer survivors			
	HR (95% CI)^b per 1 SD increase in age acceleration		
Sex group	KDM-BA-Accel	PhenoAgeAccel	SA-Accel
Female (59 deaths out of 511 survivors)	1.26 (0.96, 1.65)	1.50 (1.12, 2.01)	0.82 (0.52, 1.31)
Male (63 deaths out of 435 survivors)	1.06 (0.77, 1.50)	1.54 (0.99, 2.40)	0.92 (0.62, 1.38)
P-interaction	0.44	0.87	0.60
Sample B^a: 582 cancer survivors			
	HR (95% CI)^b per 1 SD increase in age acceleration		
Sex group	HannumAccel	HorvathAccel	LevineAccel
Female (48 deaths out of 314 survivors)	1.51 (1.26, 1.81)	1.37 (1.01, 1.87)	1.65 (1.35, 2.01)
Male (55 deaths out of 268 survivors)	1.12 (0.86, 1.45)	1.09 (0.77, 1.53)	1.55 (1.18, 2.03)
P-interaction	0.27	0.55	0.99
Sex group	GrimAgeAccel	POA	
Female (48 deaths out of 314 survivors)	2.97 (2.21, 4.00)	1.44 (1.00, 2.07)	
Male (55 deaths out of 268 survivors)	1.55 (1.03, 2.33)	1.09 (0.77, 1.54)	
P-interaction	0.0017	0.06	

^aSample A includes cancer survivors who had measures of KDM-BA, PhenoAge, and SA. Sample B includes cancer survivors who had data on ECs.

^bThe model was adjusted for chronological age, race/ethnicity, BMI, smoking status, comorbidity index, and CMV infection.