

Supplemental Table 8. Association between aging constructs and risk of total cancer, HRS (2016-2020)

Sample A: 4,413 participants without a history of cancer in 2016^a				
Aging constructs	No. of cancer cases	Total person-years	HR (95% CI) per 1 SD increase in aging constructs	
			Model 1 ^b	Model 2 ^c
KDM-BA-Accel (SD = 3.76 years)			1.05 (0.89, 1.23)	1.00 (0.84, 1.20)
PhenoAgeAccel (SD = 7.07 years)	182	17,405	1.06 (0.88, 1.28)	0.95 (0.76, 1.18)
SA-Accel (SD = 10.35 years)			1.00 (0.82, 1.22)	0.98 (0.82, 1.16)
Sample B: 2,688 respondents without a history of cancer in 2016^a				
Aging constructs	No. of cancer cases	Total person-years	HR (95% CI) per 1 SD increase in aging constructs	
			Model 1 ^b	Model 2 ^c
HannumAccel (SD = 5.13 years)			1.15 (0.95, 1.40)	1.12 (0.90, 1.40)
HorvathAccel (SD = 6.26 years)			1.06 (0.88, 1.29)	1.04 (0.87, 1.26)
LevineAccel (SD = 6.77 years)			1.07 (0.86, 1.33)	1.05 (0.83, 1.32)
GrimAgeAccel (SD = 4.64 years)	122	10,366	1.19 (0.98, 1.45)	1.09 (0.81, 1.45)
POA (SD = 0.09 years of physiological decline per one chronological year)			1.23 (0.99, 1.03)	1.14 (0.81, 1.59)

^aSample A includes 4,413 participants who had measures of KDM-BA, PhenoAge, and SA. Sample B includes 2,688 participants who had data on ECs. Participants with a prevalent cancer in 2016 and with missing cancer status after 2016 or missing cancer diagnosis time were excluded.

^bModel 1 was adjusted for chronological age.

^cModel 2 was adjusted for chronological age, sex, race/ethnicity, BMI, smoking status, comorbidity index, and CMV infection.