

Supplementary Table S1. Associations between isolated spina bifida and placental characteristics and pathology in cases and controls.

	PT controls (n=12)	SB controls (n=10)	Cases (n=12)	Unadjusted p value		Adjusted p value †	
Placental anthropometry							
Weight (g)	455 (378, 560) ^a	352 (304, 399) ^b	263 (175, 370) ^b	0.006		0.001	
Missing (n [%])	0	8 (80)	2 (16.7)				
zWeight *	1.3 (0.7, 2.2) ^a	-2.7 (-3.1, -2.4) ^b	-0.4 (-1.6, 0.2) ^b	0.0003		0.0003	
Missing (n [%])	0	8 (80)	3 (25)				
Birthweight: placental weight ratio	4.4 (4, 4.7) ^a	8.1 (7.8, 8.4) ^{ab}	5.5 (3.9, 7.4) ^b	0.1		0.002	
Missing (n [%])	0	8 (80)	2 (16.7)				
Placental histopathology							
% Intervillous fibrinoid	0 (0, 4.3) ^a	5 (5, 15) ^b	5 (5, 5) ^{ab}	0.0004		0.003	
% Stellate reticulum	25 (10, 30)	10 (8.8, 35)	30 (16.3, 57.5)	0.13		0.45	
% Terminal villi	30 (30, 35)	30 (20, 36.3)	22.5 (11.3, 43.8)	0.7		0.32	
% Stem villi	15 (11.3, 20) ^a	10 (8.8, 10) ^b	5 (5, 8.8) ^b	0.0008		0.0003	
% Mature intermediate villi	15 (11.3, 15)	10 (8.8, 11.3)	10 (5, 10)	0.02		0.09	
% Immature intermediate villi	10 (5, 13.8) ^a	10 (5, 28.8) ^b	32.5 (6.3, 56.3) ^b	0.1		0.01	
Branching maturity score							
Branching maturity score	0.05 (0.03, 0.1)	0.04 (0.01, 0.1)	0.01 (0.003, 0.1)	0.18		0.58	
Villous stromal maturity score							
Villous stromal maturity score	0.1 (0.06, 0.3)	0.18 (0.1, 0.3)	0.06 (0.03, 0.1)	0.11		0.66	
Syncytiotrophoblast maturity score							
Syncytiotrophoblast maturity score	3 (2, 3)	3.5 (3, 4)	2 (1, 3)	0.03		0.75	
Odds of placental pathology in cases (n=12)							
	PT controls (n=12)	SB controls (n=10)	Cases (n=12)	(vs. PT cons)		(vs. SB cons)	
Placental pathology				OR (95% CI)	aOR (95% CI) †	OR (95% CI)	aOR (95% CI) †
Hofbauer cells (n [%])				10 (1.7, 89.8)*	16.2 (1.4, 580)*	8 (1.3, 73.1)*	0.75 (0.03, 14.9)
Many	2 (16.7)	2 (20)	8 (66.7)				
Few	10 (83.3)	8 (80)	4 (33.3)				
Cytotrophoblast density (n [%])				11 (1.42, 235)*	14.9 (1.12, 609)*	-	-
Thick	1 (8.33)	0	6 (50)				
Thin/None	11 (91.7)	10 (100)	6 (50)				
Syncytiotrophoblast density (n [%])				15.4 (2, 332)**	146 (2.95, 3.46e5)**	12.6 (1.6, 274)*	0.14 (0.0005, 11)
Thick	1 (8.3)	1 (10)	7 (58.3)				
Thin	11 (91.7)	9 (90)	5 (41.7)				
Vasculosyncytial membrane (n [%])				0.3 (0.03, 1.7)	0.09 (0.003, 1.2)	0.47 (0.05, 3.5)	22.5 (0.8, 1304)
Many	5 (41.7)	3 (30)	2 (16.7)				
Few/None	7 (58.3)	7 (70)	10 (83.3)				
Villi capillarisation (n [%])				-	-	-	-
Many	0	2 (20)	0				
Few/None	12 (100)	8 (80)	12 (100)				
Syncytial knots (n [%])				2.2 (0.18, 51.7)	1.69 (0.1, 51.1)	0.2 (0.02, 1.3)	6 (0.2, 369)
Many	1 (8.3)	5 (50)	2 (16.7)				
Few/None	11 (91.7)	5 (50)	10 (83.3)				

White et al. (2023) *Altered placental phenotype and increased risk of placental pathology in fetal spina bifida: a matched case-control study*

Collagen in stem villi (n [%])				-	-	-	-
Lots	6 (50)	4 (40)	0				
Some/None	6 (50)	6 (60)	12 (100)				
Smooth muscle around stem vessels (n [%])				0.13 (0.006, 1)	0.1 (0.003, 0.9)*	0.1 (0.004, 0.8)*	0.8 (0.02, 18)
Lots	5 (41.7)	5 (50)	1 (8.3)				
Some/None	7 (58.3)	5 (50)	11 (91.7)				
Accelerated maturation (n [%])				0.1 (0.02, 0.9)*	0.2 (0.02, 1.1)	-	-
Yes	7 (58.3)	0	2 (16.7)				
No	5 (41.7)	10 (100)	10 (83.3)				
Distal villous hypoplasia (n [%])				0.2 (0.02, 1.2)	0.2 (0.02, 1.5)	-	-
Any	6 (50)	0	2 (16.7)				
None	6 (50)	10 (100)	10 (83.3)				
Delayed villous maturation (n [%])				-	-	0.8 (0.1, 7)	7 (0.11, 457)
Yes	0	2 (20)	2 (16.7)				
No	12 (100)	8 (80)	10 (83.3)				
Maturation (n [%]) [‡]							
Immature	0	4 (40)	2 (16.7)	-	-	0.5 (0.05, 3.7)	89.3 (0.3, 9.6e5)
Appropriate	4 (33.3)	6 (60)	6 (50)				
Hypermature	8 (66.7)	0	4 (33.3)	0.3 (0.1, 1.8)	0.9 (0.1, 9.5)	-	-

[†] Adjusted for infant sex and gestational age at delivery.

[‡] Reference group: appropriate maturity.

Continuous data are median (IQR) with p value for one-way analysis of variance (Unadjusted: ANOVA [normal distribution/equal variance] or Wilcoxon test [non-parametric data or outliers]; Adjusted: Multiple linear regression models). Categorical data are n (%) and odds ratios or adjusted odds ratios (95% CI) for presence of pathology in case placentae (reference groups: PT controls and SB controls) from Nominal Logistic regression models with p value from Likelihood Ratio Chi Square test. Statistical significance is denoted as *p<.05, **p<.01, ***p<.001. CI = confidence interval. OR = odds ratio. aOR = adjusted odds ratio.

Supplementary Table S22. Associations between isolated spina bifida and infant birth outcomes and placental phenotype in male cases and controls.

Infant outcomes at delivery	Preterm controls (n=8)	SB controls (n=2)	Cases (n=4)	Unadjusted p value	Adjusted p value †		
Stillborn/TOP (n [%])	-	-	2 (50)	-	-		
GA at birth (weeks) ‡	33.2 (31.3, 34.6)	39 (37.9, 40)	34.8 (33.5, 36.5)	0.03	-		
<37 weeks GA at birth (n [%])	8 (100)	0	3 (75)	0.007	-		
Sub-categories of PT birth (n [%])				0.06	-		
Moderate/late PT (32-36 weeks GA)	5 (62.5)	0	3 (75)				
Very PT (28-32 weeks GA)	2 (25)	0	0				
Extremely PT (<28 weeks GA)	1 (12.5)	0	0				
Birthweight z score	0.32 (-0.35, 0.46)	-1.81 (-2.07, -1.54)	-0.1 (-0.24, 1.03)	0.007	-		
Apgar at one minute (/10)	9 (4, 9)	6.5 (4, 9)	8 (7, 9)	0.94	0.24		
Apgar score <7 (n [%])	3 (37.5)	1 (50)	0	0.23	0.08		
Missing (n [%])	1 (12.5)	0	1 (25)				
Apgar at five minutes (/10)	9 (8, 9)	9 (9, 9)	9 (9, 9)	0.27	0.41		
Apgar score <7 (n [%])	0	0	0	-	-		
Missing (n [%])	1 (12.5)	0	1 (25)				
Placental anthropometry							
Weight (g)	450 (378, 500)	304	363 (265, 461)	0.37	0.24		
Missing (n [%])	0	1 (50)	1 (25)				
zWeight *	1.12 (0.66, 2.23)	-3.08	-0.34 (-2.58, 0.41)	0.047	-		
Missing (n [%])	0	1 (50)	1 (25)				
Birthweight: placental weight ratio	4.58 (4.1, 5.25)	7.83	6.94 (5.79, 9.4)	0.06	0.18		
Missing (n [%])	0	1 (50)	1 (25)				
Placental pathology							
% Intervillous fibrinoid	0 (0, 3.75) ^a	12.5 (10, 15) ^b	5 (5, 5) ^c	0.01	0.0005		
% Stellate reticulum	17.5 (10, 30)	30 (10, 50)	20 (7.5, 36.3)	0.85	0.55		
% Terminal villi	32.5 (30, 38.8)	25 (20, 30)	35 (21.3, 60)	0.5	0.66		
% Stem villi	15 (10, 18.8) ^a	12.5 (10, 15) ^{ab}	5 (5, 12.5) ^b	0.18	0.04		
% Mature intermediate villi	15 (15, 18.8)	12.5 (5, 20)	12.5 (6.25, 18.8)	0.63	0.58		
% Immature intermediate villi	7.5 (5, 13.8)	22.5 (5, 40)	15 (3.5, 40)	0.71	0.81		
Branching maturity score	0.07 (0.03, 0.11)	0.04 (0.006, 0.08)	0.08 (0.008, 0.15)	0.58	0.82		
Villous stromal maturity score	0.18 (0.06, 0.27)	0.16 (0.04, 0.27)	0.1 (0.06, 0.28)	0.9	0.48		
Syncytiotrophoblast maturity score	3 (2.25, 3)	3 (3, 3)	2.5 (2, 3.75)	0.85	0.2		
Odds of placental pathology in cases (n=4)							
	PT controls (n=8)	SB controls (n=2)	Cases (n=4)	(vs. PT cons) OR (95% CI)	aOR (95% CI) †	(vs. SB cons) OR (95% CI)	aOR (95% CI) †
Placental pathology							
Hofbauer cells (n [%])							
Many	1 (12.5)	0	2 (50)	7 (0.45, 211)	10.9 (0.43, 1120)	-	-

White et al. (2023) *Altered placental phenotype and increased risk of placental pathology in fetal spina bifida: a matched case-control study*

Few	7 (87.5)	2 (100)	2 (50)				
Cytotrophoblast density (n [%])							
Thick	0	0	2 (50)	-	-	-	-
Thin/None	8 (100)	2 (100)	2 (50)				
Syncytiotrophoblast density (n [%])							
Thick	0	0	2 (50)	-	-	-	-
Thin	8 (100)	2 (100)	2 (50)				
Vasculosyncytial membrane (n [%])							
Many	4 (50)	0	0	-	-	-	-
Few/None	4 (50)	2 (100)	4 (100)				
Villi capillarisation (n [%])							
Many	0	0	0	-	-	-	-
Few/None	8 (100)	2 (100)	4 (100)				
Syncytial knots (n [%])							
Many	1 (12.5)	0	1 (25)	-	-	-	-
Few/None	7 (87.5)	2 (100)	3 (75)				
Collagen in stem villi (n [%])							
Lots	4 (50)	1 (50)	0	-	-	-	-
Some/None	4 (50)	1 (50)	4 (100)				
Smooth muscle around stem vessels (n [%])							
Lots	3 (37.5)	1 (50)	0	-	-	-	-
Some/None	5 (62.5)	1 (50)	4 (100)				
Accelerated maturation (n [%])							
Yes	5 (62.5)	0	1 (25)	0.2 (0.008, 2.43)	0.59 (0.02, 13.5)	-	-
No	3 (37.5)	2 (100)	3 (75)				
Distal villous hyperplasia (n [%])							
Any	4 (50)	0	1 (25)	0.33 (0.01, 4.05)	0.93 (0.03, 23.1)	-	-
None	4 (50)	2 (100)	3 (75)				
Delayed villous maturation (n [%])							
Yes	0	0	2 (50)	-	-	-	-
No	8 (100)	2 (100)	2 (50)				
Maturation (n [%])[‡]							
Immature	0	1 (50)	2 (50)	-	-	-	-
Appropriate	2 (25)	1 (50)	0				
Hypermaturation	6 (75)	0	2 (50)	-	-	-	-

[†] Adjusted for gestational age at delivery.

[‡] Reference group: appropriate maturity.

Continuous data are median (IQR) with p value for one-way analysis of variance (Unadjusted: ANOVA [normal distribution/equal variance] or Wilcoxon test [non-parametric data or outliers]; Adjusted: Standard Least Squares models). Group differences from adjusted post-hoc tests (Tukey HSD) are indicated by different subscript letters. No letters = no group differences. Categorical data are n (%) and odds ratios or adjusted odds ratios (95% CI) for presence of pathology in case placentae (reference groups: PT controls and SB controls) from Nominal Logistic regression models with p value from Likelihood Ratio Chi Square test. A ‘-’ indicates that data were unavailable. Statistical significance is denoted as *p<.05, **p<.01, ***p<.001. CI = confidence interval. PT = preterm. SB = spina bifida. TOP = termination of pregnancy. OR = odds ratio. aOR = adjusted odds ratio.

Supplementary Table S23. Associations between isolated spina bifida and infant birth outcomes and placental phenotype in female cases and controls.

Infant outcomes at delivery	Preterm controls (n=4)	SB controls (n=8)	Cases (n=8)	Unadjusted p value		Adjusted p value †	
Stillborn/TOP (n [%])	-	1 (12.5)	1 (12.5)	-	-	-	-
GA at birth (weeks) ‡	33.7 (26.8, 35.9)	39.7 (38.9, 40.2)	27.2 (22.5, 35)	0.0009	-	-	-
<37 weeks GA at birth (n [%])	4 (100)	0	7 (87.5)	0.0001	-	-	-
Sub-categories of PT birth (n [%])				0.0007	-	-	-
Moderate/late PT (32-36 weeks GA)	2 (50)	0	1 (12.5)				
Very PT (28-32 weeks GA)	1 (25)	0	2 (25)				
Extremely PT (<28 weeks GA)	1 (25)	0	4 (50)				
Birthweight z score	0.1 (-0.73, 0.99)	0.16 (-0.5, 0.45)	-0.58 (-0.71, 0.17)	0.3	-	-	-
Apgar at one minute (/10)	7.5 (5.5, 8)	9 (9, 9)	8 (4, 9)	0.08	-	-	0.16
Apgar score <7 (n [%])	1 (25)	1 (12.5)	1 (12.5)	0.72	-	-	0.27
Missing (n [%])	0	0	5 (62.5)				
Apgar at five minutes (/10)	9 (6.75, 9)	9 (9, 9)	9 (9, 9)	0.25	-	-	0.1
Apgar score <7 (n [%])	1 (25)	0	0	0.24	-	-	1
Missing (n [%])	0	0	5 (62.5)				
Placental anthropometry							
Weight (g)	495 (305, 633) ^a	399 ^{ab}	221 (147, 283) ^b	0.03	-	-	0.01
Missing (n [%])	0	7 (87.5)	1 (12.5)				
zWeight [*]	1.46 (0.79, 2.88)	-2.35	-0.62 (-1.24, 0.03)	0.007	-	-	-
Missing (n [%])	0	7 (87.5)	2 (25)				
Birthweight: placental weight ratio	4.18 (3.23, 4.58)	8.37	4.1 (3.53, 6.25)	0.39	-	-	0.03
Missing (n [%])	0	7 (87.5)	1 (12.5)				
Placental pathology							
% Intervillous fibrinoid	1 (0, 4.25)	5 (5, 12.5)	5 (1.25, 8.75)	0.06	-	-	0.19
% Stellate reticulum	27.5 (13.8, 41.3)	10 (6.25, 26.3)	40 (22.5, 67.5)	0.05	-	-	0.81
% Terminal villi	30 (18.8, 30)	32.5 (22.5, 38.8)	15 (10, 36.3)	0.39	-	-	0.47
% Stem villi	17.5 (15, 20) ^a	10 (6.25, 10) ^b	5 (5, 8.75) ^b	0.004	-	-	0.0002
% Mature intermediate villi	12.5 (6.25, 15)	10 (10, 10)	7.5 (5, 10)	0.13	-	-	0.23
% Immature intermediate villi	10 (6.25, 32.5)	10 (5, 21.3)	37.5 (15, 63.8)	0.08	-	-	0.15
Branching maturity score	0.04 (0.01, 0.07)	0.04 (0.02, 0.1)	0.007 (0.002, 0.04)	0.14	-	-	0.81
Villous stromal maturity score	0.09 (0.05, 0.23)	0.18 (0.08, 0.44)	0.05 (0.02, 0.09)	0.04	-	-	0.59
Syncytiotrophoblast maturity score	2.5 (1.25, 3)	4 (3, 4)	1.5 (1, 3)	0.03	-	-	0.75
Odds of placental pathology in cases (n=8)							
	Preterm controls (n=4)	SB controls (n=8)	Cases (n=8)	(vs. PT cons) OR (95% CI)		(vs. SB cons) OR (95% CI)	
Placental pathology				aOR (95% CI)†	aOR (95% CI)†	aOR (95% CI)†	aOR (95% CI)†
Hofbauer cells (n [%])							
Many	1 (25)	2 (25)	6 (75)	9 (0.7, 258)	15.9 (0.21, 40640)	9 (1.09, 115)*	0.12 (0.0002, 6.57)

White et al. (2023) *Altered placental phenotype and increased risk of placental pathology in fetal spina bifida: a matched case-control study*

Few	3 (75)	6 (75)	2 (25)				
Cytotrophoblast density (n [%])							
Thick	1 (25)	0	4 (50)	3 (0.25, 76.6)	0.79 (0.006, 46.9)	-	-
Thin/None	3 (75)	8 (100)	4 (50)				
Syncytiotrophoblast density (n [%])							
Thick	1 (25)	1 (12.5)	5 (62.5)	5 (0.41, 131)	5.23 (2.78, 1.44e8)	11.7 (1.21, 283)*	1.18e-5 (5.71e-17, 0.8)
Thin	3 (75)	7 (87.5)	3 (37.5)				
Vasculosyncytial membrane (n [%])							
Many	1 (25)	3 (37.5)	2 (25)	1 (0.06, 26.8)	8.07 (0.68, 28792)	0.56 (0.05, 4.72)	470 (2, 4.08e8)
Few/None	3 (75)	5 (62.5)	6 (75)				
Villi capillarisation (n [%])							
Many	0	2 (25)	0	-	-	-	-
Few/None	4 (100)	6 (75)	8 (100)				
Syncytial knots (n [%])							
Many	0	5 (63.5)	1 (12.5)	-	-	0.09 (0.004, 0.83)*	1.5 (0.02, 165)
Few/None	4 (100)	3 (37.5)	7 (87.5)				
Collagen in stem villi (n [%])							
Lots	2 (50)	3 (37.5)	0	-	-	-	-
Some/None	2 (50)	5 (62.5)	8 (100)				
Smooth muscle around stem vessels (n [%])							
Lots	2 (50)	4 (50)	1 (12.5)	0.14 (0.005, 2.24)	0.16 (0.002, 4.79)	0.14 (0.006, 1.39)	2.43 (0.05, 141)
Some/None	2 (50)	4 (50)	7 (87.5)				
Accelerated maturation (n [%])							
Yes	2 (50)	0	1 (12.5)	0.14 (0.005, 2.24)	0.18 (0.003, 4.67)	-	-
No	2 (50)	8 (100)	7 (87.5)				
Distal villous hyperplasia (n [%])							
Any	2 (50)	0	1 (12.5)	0.14 (0.005, 2.24)	0.18 (0.003, 4.67)	-	-
None	2 (50)	8 (100)	7 (87.5)				
Delayed villous maturation (n [%])							
Yes	0	2 (25)	0	-	-	-	-
No	4 (100)	6 (75)	8 (100)				
Maturation (n [%]) [‡]							
Immature	0	3 (37.5)	0	-	-	-	-
Appropriate	2 (50)	5 (62.5)	6 (75)				
Hyper mature	2 (50)	0	2 (25)	0.33 (0.02, 4.31)	0.46 (0.03, 7.3)	-	-

[†] Adjusted for gestational age at delivery.

[‡] Reference group: appropriate maturity.

Continuous data are median (IQR) with p value for one-way analysis of variance (Unadjusted: ANOVA [normal distribution/equal variance] or Wilcoxon test [non-parametric data or outliers]; Adjusted: Standard Least Squares models). Group differences from adjusted post-hoc tests (Tukey HSD) are indicated by different subscript letters. No letters = no group differences. Categorical data are n (%) and odds ratios or adjusted odds ratios (95% CI) for presence of pathology in case placentae (reference groups: PT controls and SB controls) from Nominal Logistic regression models with p value from Likelihood Ratio Chi Square test. A '-' indicates that data were unavailable. Statistical significance is denoted as *p<.05, **p<.01, ***p<.001. CI = confidence interval. PT = preterm. SB = spina bifida. TOP = termination of pregnancy. OR = odds ratio. aOR = adjusted odds ratio.