

Text S1 – PRIMETIME data inputs

1. Population numbers	2
2. All-cause mortality rates	3
3. Background trends in overweight and obesity	4
4. Disease incidence rates.....	5
5. Disease case fatality rates.....	7
6. Disease prevalence	9
7. Background trends in disease incidence and case fatality.....	11
8. Relative risks of disease	13
9. Utility weights	15
10. Disease costs.....	16

1. Population numbers

Table S1 UK population by age and sex from the Human Mortality Database[1] (all ages in 2015) and population projections from the Office for National Statistics[2] (future 0-4 year olds)

Age group (years)	Intervention analyses			Model validation		
	Year0*	Male	Female	Year0*	Male	Female
0-4	2050	2,088,536	1,990,411	–	–	–
0-4	2045	2,069,673	1,972,502	–	–	–
0-4	2040	1,999,663	1,905,857	–	–	–
0-4	2035	1,953,494	1,861,924	–	–	–
0-4	2030	1,968,036	1,875,882	2020	2,006,190	1,910,568
0-4	2025	1,994,735	1,901,451	2015	2,063,050	1,963,632
0-4	2020	2,006,190	1,910,568	2010	1,960,543	1,871,193
0-4	2015	2,063,050	1,963,632	2005	1,755,352	1,669,413
5-9	2015	1,999,632	1,907,222	2005	1,860,676	1,771,490
10-14	2015	1,809,767	1,725,856	2005	1,979,193	1,886,300
15-19	2015	1,970,469	1,866,308	2005	1,973,931	1,911,380
20-24	2015	2,191,121	2,113,257	2005	1,953,411	1,936,362
25-29	2015	2,213,208	2,203,621	2005	1,879,089	1,892,516
30-34	2015	2,171,167	2,197,785	2005	2,109,298	2,139,208
35-39	2015	2,009,195	2,027,365	2005	2,314,153	2,352,797
40-44	2015	2,149,871	2,195,049	2005	2,256,961	2,304,185
45-49	2015	2,292,536	2,359,142	2005	1,986,295	2,015,247
50-54	2015	2,227,216	2,284,486	2005	1,824,379	1,859,121
55-59	2015	1,924,116	1,972,675	2005	1,925,996	1,968,812
60-64	2015	1,716,700	1,789,959	2005	1,505,366	1,568,625
65-69	2015	1,742,943	1,845,273	2005	1,297,881	1,402,969
70-74	2015	1,273,261	1,406,103	2005	1,075,857	1,257,363
75-79	2015	985,818	1,165,295	2005	830,104	1,104,028
80-84	2015	671,986	903,785	2005	561,950	927,136
85-89	2015	366,304	604,483	2005	217,373	472,565
90-94	2015	134,860	309,232	2005	78,864	240,969
95-100	2015	22,305	82,158	2005	13,891	67,683

* Year0 reflects the year in which future cohorts enter the PRIMETIME model simulation.

2. All-cause mortality rates




Figure S1 UK all-cause mortality from the Human Mortality Database[1] for 2005 (model validation) and 2015 (intervention analyses)

3. Background trends in overweight and obesity




Figure S2 Predicted trends in overweight and obesity with non-linear (base case) and linear (sensitivity) models from Cobiac et al.[3]

4. Disease incidence rates




Figure S3 Incidence rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2005 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)




Figure S4 Incidence rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2015 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)

5. Disease case fatality rates




Figure S5 Case fatality rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2005 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)




Figure S6 Case fatality rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2015 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)

6. Disease prevalence




Figure S7 Starting prevalence rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2005 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)




Figure S8 Starting prevalence rates, derived from Global Burden of Disease[4] estimates using disbayes,[5] – 2015 (Note: graphs are presented on different scales to illustrate each disease in sufficient detail, but this does have the effect of exaggerating variability in rates for some diseases)

7. Background trends in disease incidence and case fatality

Table S2 Annual trends in incidence rates, by sex and age group

Cause	Female			Male		
	0-34	35-64	65+	0-34	35-64	65+
1995-2005						
Ischaemic heart disease	-0.0123	-0.0256	-0.0302	-0.0223	-0.0223	-0.0284
Ischaemic stroke	-0.0083	-0.0279	-0.0300	-0.0099	-0.0248	-0.0334
Intracerebral haemorrhage	-0.0080	-0.0328	-0.0074	-0.0134	-0.0231	-0.0147
Hypertensive heart disease	0.0514	0.0289	0.1270	0.0548	0.0757	0.1290
Atrial fibrillation and flutter	-0.0043	-0.0008	-0.0050	0.0091	-0.0010	-0.0072
Diabetes mellitus type 2	0.0945	0.0022	0.0144	0.0901	0.0022	0.0472
Breast cancer	-0.0192	-0.0065	0.0008	NA	NA	NA
Colon and rectum cancer	0.0063	-0.0103	-0.0059	0.0073	-0.0080	-0.0039
Oesophageal cancer	-0.0026	-0.0013	0.0023	-0.0014	0.0061	0.0085
Kidney cancer	0.0093	0.0025	0.0218	0.0178	0.0058	0.0140
Liver cancer	0.0231	0.0294	0.0419	0.0389	0.0371	0.0428
Multiple myeloma	0.0123	0.0011	0.0074	0.0130	0.0036	0.0063
Pancreatic cancer	-0.0027	-0.0006	0.0076	-0.0025	-0.0043	0.0006
Uterine cancer	0.0422	0.0292	0.0303	NA	NA	NA
2005-2015						
Ischaemic heart disease	0.0052	0.0013	-0.0205	-0.0030	0.0007	-0.0156
Ischaemic stroke	0.0033	-0.0172	-0.0320	-0.0007	-0.0180	-0.0318
Intracerebral haemorrhage	-0.0051	-0.0202	-0.0041	-0.0091	-0.0203	-0.0121
Hypertensive heart disease	0.0185	-0.0027	0.0034	0.0086	0.0028	0.0037
Atrial fibrillation and flutter	0.0069	0.0142	0.0023	0.0069	0.0120	-0.0019
Diabetes mellitus type 2	0.0379	0.0261	0.0220	0.0376	0.0303	0.0131
Breast cancer	0.0112	-0.0073	0.0042	NA	NA	NA
Colon and rectum cancer	0.0359	0.0013	0.0008	0.0295	0.0003	-0.0021
Oesophageal cancer	0.0105	-0.0066	-0.0078	0.0166	-0.0046	-0.0028
Kidney cancer	0.0147	0.0015	0.0136	0.0214	0.0057	0.0093
Liver cancer	0.0291	0.0372	0.0418	0.0359	0.0372	0.0398
Multiple myeloma	0.0020	-0.0056	0.0056	0.0028	0.0009	0.0079
Pancreatic cancer	0.0102	0.0024	0.0076	0.0126	0.0030	0.0068
Uterine cancer	0.0395	0.0209	0.0269	NA	NA	NA

Table S3 Annual trends in case fatality rates, by sex and age group

Cause	Female			Male		
	0-34	35-64	65+	0-34	35-64	65+
1995-2005						
Ischaemic heart disease	-0.0403	-0.0433	-0.0498	-0.0179	-0.0340	-0.0491
Ischaemic stroke	-0.0335	-0.0489	-0.0085	-0.0362	-0.0505	-0.0248
Intracerebral haemorrhage	-0.0328	-0.0284	0.0013	-0.0164	-0.0173	-0.0020
Hypertensive heart disease	0.2510	0.1150	-0.1930	0.0831	0.0732	-0.0849
Atrial fibrillation and flutter	-0.0072	-0.0059	0.0147	0.0285	0.0066	0.0102
Diabetes mellitus type 2	-0.0147	-0.0156	-0.0138	-0.0128	-0.0181	-0.0183
Breast cancer	-0.0238	-0.0256	-0.0141	NA	NA	NA
Colon and rectum cancer	-0.0162	-0.0207	-0.0163	-0.0184	-0.0228	-0.0215
Oesophageal cancer	-0.0189	-0.0151	-0.0105	-0.0154	-0.0112	-0.0097
Kidney cancer	-0.0030	-0.0165	-0.0167	-0.0016	-0.0223	-0.0196
Liver cancer	-0.0238	-0.0178	-0.0140	-0.0243	-0.0240	-0.0206
Multiple myeloma	-0.0239	-0.0265	-0.0160	-0.0290	-0.0296	-0.0216
Pancreatic cancer	-0.0082	-0.0058	-0.0036	-0.0056	-0.0037	-0.0023
Uterine cancer	0.0269	-0.0130	-0.0059	NA	NA	NA
2005-2015						
Ischaemic heart disease	-0.0023	-0.0348	-0.0503	0.0127	-0.0325	-0.0467
Ischaemic stroke	-0.0647	-0.0424	-0.0292	-0.0547	-0.0320	-0.0285
Intracerebral haemorrhage	-0.0234	-0.0178	-0.0055	-0.0142	-0.0137	-0.0052
Hypertensive heart disease	-0.0668	-0.0200	0.0170	0.0257	0.0088	0.0184
Atrial fibrillation and flutter	0.0091	-0.0061	0.0098	-0.0174	-0.0126	0.0081
Diabetes mellitus type 2	-0.0084	-0.0017	-0.0048	-0.0122	-0.0013	-0.0038
Breast cancer	-0.0055	-0.0198	-0.0182	NA	NA	NA
Colon and rectum cancer	-0.0112	-0.0094	-0.0186	-0.0108	-0.0117	-0.0206
Oesophageal cancer	-0.0117	-0.0140	-0.0104	-0.0058	-0.0084	-0.0094
Kidney cancer	-0.0060	-0.0086	-0.0175	-0.0046	-0.0096	-0.0134
Liver cancer	-0.0206	-0.0110	-0.0099	-0.0218	-0.0201	-0.0195
Multiple myeloma	-0.0132	-0.0078	-0.0129	-0.0041	-0.0083	-0.0144
Pancreatic cancer	-0.0133	-0.0054	-0.0061	-0.0068	-0.0039	-0.0047
Uterine cancer	-0.0197	-0.0096	-0.0163	NA	NA	NA

8. Relative risks of disease

Table S4 Relative risks of modelled obesity-related diseases

Disease	Population subgroup	Units/Category	Mean relative risk (95% CI)	Source
Ischaemic heart disease	35-44	per 5kg/m ³²	1.66 (1.51 to 1.84)	Singh et al 2013[6]
	45-54	per 5kg/m ³¹	1.55 (1.46 to 1.64)	
	55-64	per 5kg/m ³⁰	1.44 (1.4 to 1.48)	
	65-74	per 5kg/m ²⁹	1.35 (1.32 to 1.38)	
	75-84	per 5kg/m ²⁸	1.26 (1.2 to 1.32)	
	85+	per 5kg/m ²⁷	1.14 (1.04 to 1.26)	
Ischaemic stroke	35-44	per 5kg/m ²⁶	1.86 (1.67 to 2.08)	Singh et al 2013[6]
	45-54	per 5kg/m ²⁵	1.67 (1.53 to 1.81)	
	55-64	per 5kg/m ²⁴	1.5 (1.4 to 1.6)	
	65-74	per 5kg/m ²³	1.35 (1.28 to 1.41)	
	75-84	per 5kg/m ²²	1.21 (1.16 to 1.26)	
	85+	per 5kg/m ²¹	1.04 (0.96 to 1.12)	
Intracerebral haemorrhage	35-44	per 5kg/m ²⁰	2.54 (1.96 to 3.28)	Singh et al 2013[6]
	45-54	per 5kg/m ¹⁹	2.1 (1.66 to 2.66)	
	55-64	per 5kg/m ¹⁸	1.75 (1.44 to 2.13)	
	65-74	per 5kg/m ¹⁷	1.48 (1.29 to 1.71)	
	75-84	per 5kg/m ¹⁶	1.3 (1.21 to 1.4)	
	85+	per 5kg/m ¹⁵	1.05 (0.92 to 1.2)	
Hypertensive heart disease	35-44	per 5kg/m ¹⁴	2.15 (0.8 to 5.78)	Singh et al 2013[6]
	45-54	per 5kg/m ¹³	2.02 (0.97 to 4.21)	
	55-64	per 5kg/m ¹²	1.9 (1.17 to 3.07)	
	65-74	per 5kg/m ¹¹	1.81 (1.45 to 2.26)	
	75-84	per 5kg/m ¹⁰	1.63 (1.53 to 1.74)	
	85+	per 5kg/m ⁹	1.45 (1.05 to 2.01)	
Diabetes mellitus 2	35-44	per 5kg/m ⁸	3.07 (2.28 to 4.15)	Singh et al 2013[6]
	45-54	per 5kg/m ⁷	2.66 (2.15 to 3.3)	
	55-64	per 5kg/m ⁶	2.32 (2.04 to 2.63)	
	65-74	per 5kg/m ⁵	2.03 (1.95 to 2.11)	
	75-84	per 5kg/m ⁴	1.7 (1.61 to 1.79)	
	85+	per 5kg/m ³	1.38 (1.23 to 1.56)	

Atrial fibrillation and flutter	–	per 5kg/m ²	1.28 (1.2 to 1.38)	Aune et al 2017[7]
Gallbladder and biliary diseases	–	per 5kg/m ¹	1.63 (1.49 to 1.78)	Aune et al 2015[8]
Colon and rectum cancer	–	per 5kg/m ⁰	1.05 (1.03 to 1.07)	WCRF 2018[9]
Breast cancer	women	per 5kg/m ¹	1.12 (1.09 to 1.15)	WCRF 2018[9]
Uterine cancer	women	per 5kg/m ²	1.54 (1.47 to 1.61)	Kyrgiou et al 2017[10]
Oesophageal cancer	adeno-carcinoma	per 5kg/m ³	1.54 (1.41 to 1.67)	Kyrgiou et al 2017[10]
	squamous cell carcinoma	per 5kg/m ⁴	0.63 (0.53 to 0.75)	
Kidney cancer	men	per 5kg/m ⁵	1.24 (1.17 to 1.32)	Kyrgiou et al 2017[10]
	women	per 5kg/m ⁶	1.33 (1.25 to 1.42)	
Pancreatic cancer	–	per 5kg/m ²	1.1 (1.06 to 1.14)	Kyrgiou et al 2017[10]
Multiple myeloma	–	per 5kg/m ²	1.12 (1.09 to 1.15)	Kyrgiou et al 2017[10]
Liver cancer	–	per 5kg/m ²	1.3 (1.16 to 1.46)	WCRF 2018[9]
Asthma*	<18 years	Healthy	1	Azizpour et al 2018[11]
		Overweight	1.64 (1.13 to 2.38)	
		Obese	1.92 (1.39 to 2.65)	
	18+ years	Healthy	1	Beuther et al 2007[12]
		Overweight	1.38 (1.17 to 1.62)	
		Obese	1.92 (1.43 to 2.59)	
Low back pain*	–	Healthy	1	Shiri et al 2010[13]
		Overweight	1.08 (0.9 to 1.29)	
		Obese	1.42 (1.11 to 0.181)	
Osteoarthritis knee*	50+ years	Healthy	1	Silverwood et al 2015[14]
		Overweight	1.98 (1.57 to 2.2)	
		Obese	2.66 (2.15 to 3.28)	
Osteoarthritis hip*	–	per 5kg/m ²	1.11 (1.07 to 1.16)	Jiang et al 2011[15]
Depressive disorders*	–	Healthy	1	Amiri et al 2018[16]
	–	Overweight	1.04 (0.99 to 1.11)	
	–	Obese	1.15 (1.06 to 1.25)	

* Disease included in sensitivity analyses only

Table S5 Relative risk of ischaemic heart disease and ischaemic stroke in people with type 2 diabetes

Disease	Population subgroup	Units/Category	Mean relative risk (95% CI)	Source
Ischaemic heart disease	men	Diabetes	1.85 (1.64 to 2.1)	Peters et al 2014[17]
	women	Diabetes	2.63 (2.27 to 3.06)	
Ischaemic stroke	men	Diabetes	1.83 (1.6 to 2.08)	Peters et al 2014[18]
	women	Diabetes	2.28 (1.93 to 2.69)	

9. Utility weights

Table S6 Disease-specific utility weights, estimated for the UK by Sullivan et al[19]

Disease	ICD9 code	Mean utility (SD)
Ischaemic heart disease incidence	icd410	-0.063 (0.013)
Ischaemic heart disease prevalence	icd412	-0.037 (0.026)
Ischaemic stroke incidence	icd436	-0.117 (0.012)
Ischaemic stroke prevalence	icd438	-0.073 (0.024)
Intracerebral haemorrhage incidence	icd436	-0.117 (0.012)
Intracerebral haemorrhage prevalence	icd438	-0.073 (0.024)
Hypertensive heart disease	icd401	-0.046 (0.004)
Diabetes mellitus type 2	icd250	-0.071 (0.005)
Atrial fibrillation and flutter	icd427	-0.038 (0.007)
Colon and rectum cancer	icd153	-0.067 (0.017)
Breast cancer	icd174	-0.019 (0.014)
Uterine cancer	icd202	-0.010 (0.026)
Oesophageal cancer	icd202	-0.010 (0.026)
Kidney cancer	icd189	-0.048 (0.041)
Pancreatic cancer	icd202	-0.010 (0.026)
Multiple myeloma	icd195	-0.086 (0.027)
Liver cancer	icd155	-0.093 (0.044)
Asthma	icd493	-0.046 (0.006)
Low back pain	icd724	-0.087 (0.006)
Osteoarthritis hip	icd715	-0.114 (0.008)
Osteoarthritis knee	icd715	-0.114 (0.008)
Depressive disorders	icd296	-0.127 (0.010)
Gallbladder and biliary diseases	icd574-76*	-0.062 (0.018)

* weighted by Hospital Episode Statistics admissions primary diagnosis for ICD10 K80-83.[20]

Table S7 Parameters for estimating background utility weight, , estimated for the UK by Sullivan et al[19]

Parameter	Mean utility (SD)
Age (continuous years)	-0.00027 (0.00017)
Male	0.0010 (0.00063)
Age 10-19	0.913 (0.0045)
Age 20-29	0.905 (0.0021)
Age 30-39	0.879 (0.0021)
Age 40-49	0.837 (0.0028)
Age 50-59	0.798 (0.0035)
Age 60-69	0.774 (0.0039)
Age 70-79	0.723 (0.0049)
Age 80-89	0.657 (0.0075)

10. Disease costs

Table S8 Costs of treatment in the National Health Service (see Text S2)

Condition	Units	Mean cost (SD)*
Ischaemic heart disease	prevalent case	£606 (£121)
Ischaemic stroke	prevalent case	£1,950 (£390)
Intracerebral haemorrhage	prevalent case	£2,563 (£513)
Hypertensive heart disease	prevalent case	£103 (£21)
Diabetes mellitus type 2	prevalent case	£187 (£37)
Atrial fibrillation and flutter	prevalent case	£195 (£39)
Colon and rectum cancer	incident case	£9,204 (£1,841)
Breast cancer	incident case	£12,433 (£2,487)
Uterine cancer	incident case	£2,060 (£412)
Oesophageal cancer	incident case	£2,421 (£484)
Kidney cancer	incident case	£4,979 (£996)
Pancreatic cancer	incident case	£2,695 (£539)
Multiple myeloma	incident case	£22,915 (£4,583)
Liver cancer	incident case	£2,172 (£434)
Asthma	incident case	£4,186 (£837)
Low back pain	incident case	£425 (£85)
Osteoarthritis hip	incident case	£13,951 (£2,790)
Osteoarthritis knee	incident case	£1,799 (£360)
Depressive disorders	incident case	£410 (£82)
Gallbladder and biliary diseases	incident case	£372 (£74)
Total non-modelled diseases	person	£1,099 (£220)

* Standard deviation estimated as 20% of point estimate.

** Decayed missing and filled deciduous (dmft) and permanent (DMFT) teeth.

References

1. University of California Berkeley (USA), Max Planck Institute for Demographic Research (Germany). Human Mortality Database [5 May 2021]. Available from: <https://www.mortality.org>.
2. Office for National Statistics. 2016-based National Population Projections 2017 [1 September 2020]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/datasets/z3zippedpopulationprojectionsdatafilesengland>.
3. Cobiac LJ, Scarborough P. Modelling future trajectories of obesity and body mass index in England. PLoS ONE. 2021;16(6):e0252072. doi: 10.1371/journal.pone.0252072.
4. Global Burden of Disease Study 2015 (GBD 2015) Data Resources: Institute for Health Metrics and Evaluation, University of Washington; 2016 [cited 2016 12 December 2016]. Available from: <http://ghdx.healthdata.org/gbd-2015>.
5. Jackson C. disbayes [5 May 2021]. Available from: <https://github.com/chjackson/disbayes>.
6. Singh GM, Danaei G, Farzadfar F, Stevens GA, Woodward M, Wormser D, et al. The age-specific quantitative effects of metabolic risk factors on cardiovascular diseases and diabetes: a pooled analysis. PLoS ONE. 2013;8(7):e65174.
7. Aune D, Sen A, Schlesinger S, Norat T, Janszky I, Romundstad P, et al. Body mass index, abdominal fatness, fat mass and the risk of atrial fibrillation: a systematic review and dose–response meta-analysis of prospective studies. Eur J Epidemiol. 2017;32(3):181-92.
8. Aune D, Norat T, Vatten LJ. Body mass index, abdominal fatness and the risk of gallbladder disease. Eur J Epidemiol. 2015;30(9):1009-19.
9. World Cancer Research Fund. Diet, Nutrition, Physical Activity and Cancer: a Global Perspective, Third Expert Report. World Cancer Research Fund; American Institute for Cancer Research, 2018.
10. Kyrgiou M, Kalliala I, Markozannes G, Gunter MJ, Paraskevidis E, Gabra H, et al. Adiposity and cancer at major anatomical sites: umbrella review of the literature. Bmj. 2017;356.
11. Azizpour Y, Delpisheh A, Montazeri Z, Sayehmiri K, Darabi B. Effect of childhood BMI on asthma: a systematic review and meta-analysis of case-control studies. BMC Pediatr. 2018;18(1):1-13.
12. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. Am J Respir Crit Care Med. 2007;175(7):661-6.
13. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity and low back pain: a meta-analysis. Am J Epidemiol. 2010;171(2):135-54.
14. Silverwood V, Blagojevic-Bucknall M, Jinks C, Jordan J, Protheroe J, Jordan K. Current evidence on risk factors for knee osteoarthritis in older adults: a systematic review and meta-analysis. Osteoarthritis Cartilage. 2015;23(4):507-15.
15. Jiang L, Rong J, Wang Y, Hu F, Bao C, Li X, et al. The relationship between body mass index and hip osteoarthritis: a systematic review and meta-analysis. Joint Bone Spine. 2011;78(2):150-5.

16. Amiri S, Behnezhad S, Nadinlui KB. Body Mass Index (BMI) and risk of depression in adults: A systematic review and meta-analysis of longitudinal studies. *Obesity Medicine*. 2018;12:1-12.
17. Peters SA, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia*. 2014;57(8):1542-51.
18. Peters SA, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775 385 individuals and 12 539 strokes. *Lancet*. 2014;383(9933):1973-80.
19. Sullivan PW, Slejko JF, Sculpher MJ, Ghushchyan V. Catalogue of EQ-5D scores for the United Kingdom. *Med Decis Making*. 2011;31(6):800-4.
20. NHS Digital. Hospital Episode Statistics for England: Admitted Patient Care statistics, 2019-20 2020 [21 April 2021]. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2019-20>.