



Supplementary Tables and Figures

Supplementary Figure 1: Directed acyclic graph representing the associations between tobacco smoking and serum total bilirubin with dotted lines representing the associations evaluated in this study



Supplementary Figure 2: Flowchart of participant inclusion and exclusion



Supplementary Table 1: ICD Codes for Head and Neck Cancer and Lung Cancer

Type of Cancer	ICD Codes
Head and Neck Cancer	'140', '140.1', '140.3', '140.4', '140.5', '140.6', '140.8', '140.9', '141', '141.1', '141.2', '141.3', '141.4', '141.5', '141.6', '141.8', '141.9', '142', '142.1', '142.2', '142.8', '142.9', '143', '143.1', '143.8', '143.9', '144', '144.1', '144.8', '144.9', '145', '145.1', '145.2', '145.3', '145.4', '145.5', '145.6', '145.8', '145.9', '146', '146.1', '146.2', '146.3', '146.4', '146.5', '146.6', '146.7', '146.8', '146.9', '147', '147.1', '147.2', '147.3', '147.8', '147.9', '148', '148.1', '148.2', '148.3', '148.8', '148.9', '149', '149.1', '149.8', '149.9', '160', '160.1', '160.2', '160.3', '160.4', '160.5', '160.8', '160.9', '161', '161.1', '161.2', '161.3', '161.8', '161.9', 'C00.0', 'C00.1', 'C00.2', 'C00.3', 'C00.4', 'C00.5', 'C00.6', 'C00.8', 'C00.9', 'C02.0', 'C02.1', 'C02.2', 'C02.3', 'C02.4', 'C02.8', 'C02.9', 'C06.0', 'C06.1', 'C06.2', 'C06.8', 'C06.80', 'C06.89', 'C06.9', 'C08.0', 'C08.1', 'C08.9', 'C09.0', 'C09.1', 'C09.8', 'C09.9', 'C10.0', 'C10.1', 'C10.2', 'C10.3', 'C10.4', 'C10.8', 'C10.9', 'C11.0', 'C11.1', 'C11.2', 'C11.3', 'C11.8', 'C11.9', 'C13.0', 'C13.1', 'C13.2', 'C13.8', 'C13.9', 'C14.0', 'C14.2', 'C14.8', 'C30.0'.

Lung Cancer	'C34.0', 'C34.1', 'C34.2', 'C34.3', 'C34.8', 'C34.9', '162.0', '162.2', '162.3', '162.4', '162.5', '162.8', '162.9'
-------------	---

Supplementary Table 2: Baseline characteristics of participants, stratified by case: control status

		Cancer Controls	LC Cases	HNC Cases
n		388793	2378	2039
Patient Age, mean (SD)		57.0 (18.7)	75.8 (10.9)	69.7 (13.5)
SB mg/dL, mean (SD)		0.6 (0.3)	0.5 (0.2)	0.5 (0.2)
Sex, n (%)	Female	213979 (55.0)	1259 (52.9)	589 (28.9)
	Male	174814 (45.0)	1119 (47.1)	1450 (71.1)
Smoking History, n (%)	Never Smokers	270277 (69.6)	831 (34.9)	1010 (49.5)
	Ever Smokers	118516 (30.4)	1547 (65.1)	1029 (50.5)
Self-reported Race, n (%)	American Indian or Alaska Native	2045 (0.7)	10 (0.5)	6 (0.4)
	Asian	36632 (13.4)	395 (19.6)	208 (12.8)
	Black or African American	15860 (5.8)	111 (5.5)	59 (3.6)
	Native Hawaiian or Other Pacific Islander	530 (0.2)	9 (0.4)	2 (0.1)

	Other Race	32185 (11.8)	203 (10.1)	171 (10.5)
	White or Caucasian	161596 (59.1)	1111 (55.2)	1078 (66.2)
Self-Reported Ethnicity, n (%)	Hispanic or Latin	33631 (14.3)	136 (7.0)	129 (9.0)
	Not Hispanic or Latin	211951 (82.2)	1681 (88.3)	1380 (87.9)

Supplementary Table 3: Effects of HNC and LC on SB

Results of linear regressions adjusted for age, sex, SIRE to evaluate the associations between HNC/LC and SB levels(dependent variable).

Variables	Effect on SB		
	mg/dL	95% CI	P-Value
HNC	-0.11	-0.13, -0.09	< 0.0001
HNC (adjusted for smoking)	-0.11	-0.13, -0.09	< 0.0001
LC	-0.09	-0.1, -0.07	< 0.0001
LC (adjusted for smoking)	-0.08	-0.1, -0.07	< 0.0001

Supplementary Table 4a: Propensity matched group baseline characteristics stratified by HNC

		Overall	HNC Controls	HNC Cases
n		4074	2037	2037
Patient Age, mean (SD)		69.7 (13.5)	69.7 (13.5)	69.7 (13.5)
Serum Total Bilirubin, [Q1,Q3]		0.5 [0.4,0.7]	0.6 [0.4,0.8]	0.5 [0.4,0.6]
Sex, n (%)	Female	1174 (28.8)	587 (28.8)	587 (28.8)
	Male	2900 (71.2)	1450 (71.2)	1450 (71.2)

Smoking History, n (%)	0	2018 (49.5)	1009 (49.5)	1009 (49.5)
	1	2056 (50.5)	1028 (50.5)	1028 (50.5)
Self-reported Race, n (%)	American Indian or Alaska Native	11 (0.3)	6 (0.4)	5 (0.3)
	Asian	414 (12.7)	207 (12.7)	207 (12.7)
	Black or African American	118 (3.6)	59 (3.6)	59 (3.6)
	Native Hawaiian or Other Pacific Islander	4 (0.1)	2 (0.1)	2 (0.1)
	Other Race	342 (10.5)	171 (10.5)	171 (10.5)
	Unknown Race	210 (6.5)	105 (6.4)	105 (6.5)
	White or Caucasian	2156 (66.2)	1078 (66.2)	1078 (66.3)
Self Reported Ethnicity, n (%)	Hispanic or Latin	343 (9.1)	173 (9.2)	170 (9.0)
	Not Hispanic or Latin	3310 (87.9)	1655 (87.8)	1655 (88.0)
	Unknown Ethnicity	112 (3.0)	56 (3.0)	56 (3.0)

Supplementary Table 4b: Propensity matched group baseline characteristics stratified by LC

		Overall	Lung Cancer Controls	Lung Cancer Cases
n		4746	2373	2373
Patient Age, mean (SD)		75.8 (10.9)	75.8 (10.9)	75.8 (10.9)
Serum Total Bilirubin, [Q1,Q3]		0.5 [0.4,0.7]	0.5 [0.4,0.7]	0.5 [0.4,0.7]
Sex, n (%)	Female	2512 (52.9)	1256 (52.9)	1256 (52.9)
	Male	2234 (47.1)	1117 (47.1)	1117 (47.1)

Smoking History, n (%)	Never Smokers	1656 (34.9)	828 (34.9)	828 (34.9)
	Ever Smokers	3090 (65.1)	1545 (65.1)	1545 (65.1)
Self Reported Race, n (%)	American Indian or Alaska Native	15 (0.4)	5 (0.2)	10 (0.5)
	Asian	786 (19.6)	393 (19.6)	393 (19.6)
	Black or African American	222 (5.5)	111 (5.5)	111 (5.5)
	Native Hawaiian or Other Pacific Islander	12 (0.3)	6 (0.3)	6 (0.3)
	Other Race	406 (10.1)	203 (10.1)	203 (10.1)
	Unknown Race	348 (8.7)	174 (8.7)	174 (8.7)
	White or Caucasian	2222 (55.4)	1111 (55.5)	1111 (55.3)
Self Reported Ethnicity, n (%)	Hispanic or Latin	326 (7.3)	171 (7.7)	155 (7.0)
	Not Hispanic or Latin	3918 (88.1)	1959 (87.8)	1959 (88.4)
	Unknown Ethnicity	202 (4.5)	101 (4.5)	101 (4.6)

Supplementary Table 5: Baseline characteristics of European GIA UCLA ATLAS Biobank participants included in SB-PGS analysis

		Cancer Controls	LC Cases	HNC Cases
n		14871	124	152
Patient Age, mean (SD)		63.2	73.8	70.1
SB mg/dL, mean (SD)		0.6	0.5	0.5
Sex, n (%)	Female	7432 (50)	63 (50)	30 (20)

	Male	7439 (50)	61 (50)	122 (80)
Smoking History, n (%)	Never Smokers	9421 (64)	35 (28)	82 (54)
	Ever Smokers	5450 (36)	89 (72)	70 (46)