

Text S1 – PRIMETIME data inputs

1. Population numbers	2
2. All-cause mortality rates	3
3. Background trends in overweight and obesity	4
4. Dental caries rates	5
5. Disease incidence rates.....	6
6. Disease case fatality rates.....	7
7. Disease prevalence	8
8. Background trends in disease incidence and case fatality.....	9
9. Relative risks of disease	10
10. Utility weights	12
11. Disease costs.....	13

1. Population numbers

Table S1 UK population by age and sex from the Human Mortality Database¹ (all ages in 2015) and population projections from the Office for National Statistics² (future 0-4 year olds)

Age group (years)	Year0*	Starting population	
		Male	Female
0-4	2050	2,088,536	1,990,411
0-4	2045	2,069,673	1,972,502
0-4	2040	1,999,663	1,905,857
0-4	2035	1,953,494	1,861,924
0-4	2030	1,968,036	1,875,882
0-4	2025	1,994,735	1,901,451
0-4	2020	2,006,190	1,910,568
0-4	2015	2,063,050	1,963,632
5-9	2015	1,999,632	1,907,222
10-14	2015	1,809,767	1,725,856
15-19	2015	1,970,469	1,866,308
20-24	2015	2,191,121	2,113,257
25-29	2015	2,213,208	2,203,621
30-34	2015	2,171,167	2,197,785
35-39	2015	2,009,195	2,027,365
40-44	2015	2,149,871	2,195,049
45-49	2015	2,292,536	2,359,142
50-54	2015	2,227,216	2,284,486
55-59	2015	1,924,116	1,972,675
60-64	2015	1,716,700	1,789,959
65-69	2015	1,742,943	1,845,273
70-74	2015	1,273,261	1,406,103
75-79	2015	985,818	1,165,295
80-84	2015	671,986	903,785
85-89	2015	366,304	604,483
90-94	2015	134,860	309,232
95-100	2015	22,305	82,158

* Year0 reflects the year in which future cohorts enter the PRIMETIME model simulation.

2. All-cause mortality rates




Figure S1 UK all-cause mortality, by single year of age and sex, from the Human Mortality Database.¹

3. Background trends in overweight and obesity




Figure S2 Predicted trends in overweight and obesity with non-linear (base case) and linear (sensitivity) models from Cobiac et al.³

4. Dental caries rates




Figure S3 (a) prevalence of and (b) mean number of decayed, missing and filled teeth that are deciduous (dmft) and permanent (DMFT); derived from data collected in the Child and Adult Dental Health Surveys,^{4,5} adjusted for congenital tooth absence,⁶ and causes of tooth extraction.^{7,8}

5. Disease incidence rates




Figure S4 Incidence rates, derived from Global Burden of Disease⁹ estimates using disbayes.¹⁰ (NB. Graphs are presented on different scales to show detail, but in some cases this may give an exaggerated appearance of variability across ages.

6. Disease case fatality rates




Figure S5 Case fatality rates, derived from Global Burden of Disease⁹ estimates using disbayes.¹⁰

7. Disease prevalence




Figure S6 Starting prevalence rates, derived from Global Burden of Disease⁹ estimates using disbayes.¹⁰

8. Background trends in disease incidence and case fatality

Table S2 Annual trends in incidence rates, by sex and age group

Cause	Male			Female		
	0-34	35-64	65+	0-34	35-64	65+
Ischaemic heart disease	0.0052	0.0013	-0.0205	-0.0030	0.0007	-0.0156
Ischaemic stroke	0.0033	-0.0172	-0.0320	-0.0007	-0.0180	-0.0318
Intracerebral haemorrhage	-0.0051	-0.0202	-0.0041	-0.0091	-0.0203	-0.0121
Hypertensive heart disease	0.0185	-0.0027	0.0034	0.0086	0.0028	0.0037
Atrial fibrillation and flutter	0.0069	0.0142	0.0023	0.0069	0.0120	-0.0019
Diabetes mellitus type 2	0.0379	0.0261	0.0220	0.0376	0.0303	0.0131
Breast cancer	0.0112	-0.0073	0.0042	NA	NA	NA
Colon and rectum cancer	0.0359	0.0013	0.0008	0.0295	0.0003	-0.0021
Oesophageal cancer	0.0105	-0.0066	-0.0078	0.0166	-0.0046	-0.0028
Kidney cancer	0.0147	0.0015	0.0136	0.0214	0.0057	0.0093
Liver cancer	0.0291	0.0372	0.0418	0.0359	0.0372	0.0398
Multiple myeloma	0.0020	-0.0056	0.0056	0.0028	0.0009	0.0079
Pancreatic cancer	0.0102	0.0024	0.0076	0.0126	0.0030	0.0068
Uterine cancer	0.0395	0.0209	0.0269	NA	NA	NA

Table S3 Annual trends in case fatality rates, by sex and age group

Cause	Male			Female		
	0-34	35-64	65+	0-34	35-64	65+
Ischaemic heart disease	-0.0023	-0.0348	-0.0503	0.0127	-0.0325	-0.0467
Ischaemic stroke	-0.0647	-0.0424	-0.0292	-0.0547	-0.0320	-0.0285
Intracerebral haemorrhage	-0.0234	-0.0178	-0.0055	-0.0142	-0.0137	-0.0052
Hypertensive heart disease	-0.0668	-0.0200	0.0170	0.0257	0.0088	0.0184
Atrial fibrillation and flutter	0.0091	-0.0061	0.0098	-0.0174	-0.0126	0.0081
Diabetes mellitus type 2	-0.0084	-0.0017	-0.0048	-0.0122	-0.0013	-0.0038
Breast cancer	-0.0055	-0.0198	-0.0182	NA	NA	NA
Colon and rectum cancer	-0.0112	-0.0094	-0.0186	-0.0108	-0.0117	-0.0206
Oesophageal cancer	-0.0117	-0.0140	-0.0104	-0.0058	-0.0084	-0.0094
Kidney cancer	-0.0060	-0.0086	-0.0175	-0.0046	-0.0096	-0.0134
Liver cancer	-0.0206	-0.0110	-0.0099	-0.0218	-0.0201	-0.0195
Multiple myeloma	-0.0132	-0.0078	-0.0129	-0.0041	-0.0083	-0.0144
Pancreatic cancer	-0.0133	-0.0054	-0.0061	-0.0068	-0.0039	-0.0047
Uterine cancer	-0.0197	-0.0096	-0.0163	NA	NA	NA

9. Relative risks of disease

Table S4 Relative risks of modelled obesity-related diseases

Disease	Population subgroup	Units/Category	Mean relative risk (95% CI)	Source
Ischaemic heart disease	35-44	per 5kg/m ³²	1.66 (1.51 to 1.84)	Singh et al 2013 ¹¹
	45-54	per 5kg/m ³¹	1.55 (1.46 to 1.64)	
	55-64	per 5kg/m ³⁰	1.44 (1.4 to 1.48)	
	65-74	per 5kg/m ²⁹	1.35 (1.32 to 1.38)	
	75-84	per 5kg/m ²⁸	1.26 (1.2 to 1.32)	
	85+	per 5kg/m ²⁷	1.14 (1.04 to 1.26)	
Ischaemic stroke	35-44	per 5kg/m ²⁶	1.86 (1.67 to 2.08)	Singh et al 2013 ¹¹
	45-54	per 5kg/m ²⁵	1.67 (1.53 to 1.81)	
	55-64	per 5kg/m ²⁴	1.5 (1.4 to 1.6)	
	65-74	per 5kg/m ²³	1.35 (1.28 to 1.41)	
	75-84	per 5kg/m ²²	1.21 (1.16 to 1.26)	
	85+	per 5kg/m ²¹	1.04 (0.96 to 1.12)	
Intracerebral haemorrhage	35-44	per 5kg/m ²⁰	2.54 (1.96 to 3.28)	Singh et al 2013 ¹¹
	45-54	per 5kg/m ¹⁹	2.1 (1.66 to 2.66)	
	55-64	per 5kg/m ¹⁸	1.75 (1.44 to 2.13)	
	65-74	per 5kg/m ¹⁷	1.48 (1.29 to 1.71)	
	75-84	per 5kg/m ¹⁶	1.3 (1.21 to 1.4)	
	85+	per 5kg/m ¹⁵	1.05 (0.92 to 1.2)	
Hypertensive heart disease	35-44	per 5kg/m ¹⁴	2.15 (0.8 to 5.78)	Singh et al 2013 ¹¹
	45-54	per 5kg/m ¹³	2.02 (0.97 to 4.21)	
	55-64	per 5kg/m ¹²	1.9 (1.17 to 3.07)	
	65-74	per 5kg/m ¹¹	1.81 (1.45 to 2.26)	
	75-84	per 5kg/m ¹⁰	1.63 (1.53 to 1.74)	
	85+	per 5kg/m ⁹	1.45 (1.05 to 2.01)	
Diabetes mellitus 2	35-44	per 5kg/m ⁸	3.07 (2.28 to 4.15)	Singh et al 2013 ¹¹
	45-54	per 5kg/m ⁷	2.66 (2.15 to 3.3)	
	55-64	per 5kg/m ⁶	2.32 (2.04 to 2.63)	
	65-74	per 5kg/m ⁵	2.03 (1.95 to 2.11)	

	75-84	per 5kg/m ⁴	1.7 (1.61 to 1.79)		
	85+	per 5kg/m ³	1.38 (1.23 to 1.56)		
Atrial fibrillation and flutter	–	per 5kg/m ²	1.28 (1.2 to 1.38)	Aune et al 2017 ¹²	
Gallbladder and biliary diseases	–	per 5kg/m ¹	1.63 (1.49 to 1.78)	Aune et al 2015 ¹³	
Colon and rectum cancer	–	per 5kg/m ⁰	1.05 (1.03 to 1.07)	WCRF 2018 ¹⁴	
Breast cancer	women	per 5kg/m ¹	1.12 (1.09 to 1.15)	WCRF 2018 ¹⁴	
Uterine cancer	women	per 5kg/m ²	1.54 (1.47 to 1.61)	Kyrgiou et al 2017 ¹⁵	
Oesophageal cancer	adeno-carcinoma	per 5kg/m ³	1.54 (1.41 to 1.67)	Kyrgiou et al 2017 ¹⁵	
		squamous cell carcinoma	per 5kg/m ⁴		0.63 (0.53 to 0.75)
Kidney cancer	men	per 5kg/m ⁵	1.24 (1.17 to 1.32)	Kyrgiou et al 2017 ¹⁵	
	women	per 5kg/m ⁶	1.33 (1.25 to 1.42)		
Pancreatic cancer	–	per 5kg/m ²	1.1 (1.06 to 1.14)	Kyrgiou et al 2017 ¹⁵	
Multiple myeloma	–	per 5kg/m ²	1.12 (1.09 to 1.15)	Kyrgiou et al 2017 ¹⁵	
Liver cancer	–	per 5kg/m ²	1.3 (1.16 to 1.46)	WCRF 2018 ¹⁴	
Asthma*	<18 years	Healthy	1	Azizpour et al 2018 ¹⁶	
		Overweight	1.64 (1.13 to 2.38)		
		Obese	1.92 (1.39 to 2.65)		
	18+ years	Healthy	1		Beuther et al 2007 ¹⁷
		Overweight	1.38 (1.17 to 1.62)		
		Obese	1.92 (1.43 to 2.59)		
Low back pain*	–	Healthy	1	Shiri et al 2010 ¹⁸	
		Overweight	1.08 (0.9 to 1.29)		
		Obese	1.42 (1.11 to 0.181)		
Osteoarthritis knee*	50+ years	Healthy	1	Silverwood et al 2015 ¹⁹	
		Overweight	1.98 (1.57 to 2.2)		
		Obese	2.66 (2.15 to 3.28)		
Osteoarthritis hip*	–	per 5kg/m ²	1.11 (1.07 to 1.16)	Jiang et al 2011 ²⁰	
Depressive disorders*	–	Healthy	1	Amiri et al 2018 ²¹	
		Overweight	1.04 (0.99 to 1.11)		
		Obese	1.15 (1.06 to 1.25)		

* Disease included in sensitivity analyses only

Table S5 Relative risk of ischaemic heart disease and ischaemic stroke in people with type 2 diabetes

Disease	Population subgroup	Units/Category	Mean relative risk (95% CI)	Source
Ischaemic heart disease	men	Diabetes	1.85 (1.64 to 2.1)	Peters et al 2014 ²²
	women	Diabetes	2.63 (2.27 to 3.06)	
Ischaemic stroke	men	Diabetes	1.83 (1.6 to 2.08)	Peters et al 2014 ²³
	women	Diabetes	2.28 (1.93 to 2.69)	

10. Utility weights

Table S6 Disease-specific utility weights, estimated for the UK by Sullivan et al²⁴

Disease	ICD9 code	Mean utility (SD)
Ischaemic heart disease incidence	icd410	-0.063 (0.013)
Ischaemic heart disease prevalence	icd412	-0.037 (0.026)
Ischaemic stroke incidence	icd436	-0.117 (0.012)
Ischaemic stroke prevalence	icd438	-0.073 (0.024)
Intracerebral haemorrhage incidence	icd436	-0.117 (0.012)
Intracerebral haemorrhage prevalence	icd438	-0.073 (0.024)
Hypertensive heart disease	icd401	-0.046 (0.004)
Diabetes mellitus type 2	icd250	-0.071 (0.005)
Atrial fibrillation and flutter	icd427	-0.038 (0.007)
Colon and rectum cancer	icd153	-0.067 (0.017)
Breast cancer	icd174	-0.019 (0.014)
Uterine cancer	icd202	-0.010 (0.026)
Oesophageal cancer	icd202	-0.010 (0.026)
Kidney cancer	icd189	-0.048 (0.041)
Pancreatic cancer	icd202	-0.010 (0.026)
Multiple myeloma	icd195	-0.086 (0.027)
Liver cancer	icd155	-0.093 (0.044)
Asthma	icd493	-0.046 (0.006)
Low back pain	icd724	-0.087 (0.006)
Osteoarthritis hip	icd715	-0.114 (0.008)
Osteoarthritis knee	icd715	-0.114 (0.008)
Depressive disorders	icd296	-0.127 (0.010)
Dental caries	icd521	-0.002 (0.012)
Gallbladder and biliary diseases	icd574-76*	-0.062 (0.018)

* weighted by Hospital Episode Statistics admissions primary diagnosis for ICD10 K80-83.²⁵

Table S7 Parameters for estimating background utility weight, , estimated for the UK by Sullivan et al²⁴

Parameter	Mean utility (SD)
Age (continuous years)	-0.00027 (0.00017)
Male	0.0010 (0.00063)
Age 10-19	0.913 (0.0045)
Age 20-29	0.905 (0.0021)
Age 30-39	0.879 (0.0021)
Age 40-49	0.837 (0.0028)
Age 50-59	0.798 (0.0035)
Age 60-69	0.774 (0.0039)
Age 70-79	0.723 (0.0049)
Age 80-89	0.657 (0.0075)

11. Disease costs

Table S8 Costs of treatment in the National Health Service

Condition	Units	Mean cost (SD)*
Dental caries	dmft/DMFT**	£85 (£17)
Ischaemic heart disease	prevalent case	£606 (£121)
Ischaemic stroke	prevalent case	£1,950 (£390)
Intracerebral haemorrhage	prevalent case	£2,563 (£513)
Hypertensive heart disease	prevalent case	£103 (£21)
Diabetes mellitus type 2	prevalent case	£187 (£37)
Atrial fibrillation and flutter	prevalent case	£195 (£39)
Colon and rectum cancer	incident case	£9,204 (£1,841)
Breast cancer	incident case	£12,433 (£2,487)
Uterine cancer	incident case	£2,060 (£412)
Oesophageal cancer	incident case	£2,421 (£484)
Kidney cancer	incident case	£4,979 (£996)
Pancreatic cancer	incident case	£2,695 (£539)
Multiple myeloma	incident case	£22,915 (£4,583)
Liver cancer	incident case	£2,172 (£434)
Asthma	incident case	£4,186 (£837)
Low back pain	incident case	£425 (£85)
Osteoarthritis hip	incident case	£13,951 (£2,790)
Osteoarthritis knee	incident case	£1,799 (£360)
Depressive disorders	incident case	£410 (£82)
Gallbladder and biliary diseases	incident case	£372 (£74)
Total non-modelled diseases	person	£1,099 (£220)

* Standard deviation estimated as 20% of point estimate.

** Decayed missing and filled deciduous (dmft) and permanent (DMFT) teeth.

References

1. University of California Berkeley (USA), Max Planck Institute for Demographic Research (Germany). Human Mortality Database [Available from: <https://www.mortality.org> accessed 5 May 2021.
2. Office for National Statistics. 2016-based National Population Projections 2017 [Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationprojections/datasets/z3zippedpopulationprojectionsdatafilesengland> accessed 1 September 2020.
3. Cobiac LJ, Scarborough P. Modelling future trajectories of obesity and body mass index in England. *PLoS ONE* 2021;16(6):e0252072. doi: 10.1371/journal.pone.0252072
4. Adult Dental Health Survey, 2009. UK Data Service. Available from: <http://doi.org/10.5255/UKDA-SN-6884-22012>.
5. Children's Dental Health Survey, 2013. UK Data Service. Available from: <http://doi.org/10.5255/UKDA-SN-7774-12015>.
6. Khalaf K, Miskelly J, Voge E, et al. Prevalence of hypodontia and associated factors: a systematic review and meta-analysis. *J Orthod* 2014;41(4):299-316.
7. McCaul L, Jenkins W, Kay E. The reasons for the extraction of various tooth types in Scotland: a 15-year follow up. *J Dent* 2001;29(6):401-07.
8. Richards W, Ameen J, Coll A, et al. Reasons for tooth extraction in four general dental practices in South Wales. *Br Dent J* 2005;198(5):275-78.
9. Global Burden of Disease Study 2015 (GBD 2015) Data Resources: Institute for Health Metrics and Evaluation, University of Washington; 2016 [Available from: <http://ghdx.healthdata.org/gbd-2015> accessed 12 December 2016 2016.
10. Jackson C. disbayes [Available from: <https://github.com/chjackson/disbayes> accessed 5 May 2021.
11. Singh GM, Danaei G, Farzadfar F, et al. The age-specific quantitative effects of metabolic risk factors on cardiovascular diseases and diabetes: a pooled analysis. *PLoS ONE* 2013;8(7):e65174.
12. Aune D, Sen A, Schlesinger S, et al. Body mass index, abdominal fatness, fat mass and the risk of atrial fibrillation: a systematic review and dose–response meta-analysis of prospective studies. *Eur J Epidemiol* 2017;32(3):181-92.
13. Aune D, Norat T, Vatten LJ. Body mass index, abdominal fatness and the risk of gallbladder disease. *Eur J Epidemiol* 2015;30(9):1009-19.

14. World Cancer Research Fund. Diet, Nutrition, Physical Activity and Cancer: a Global Perspective, Third Expert Report: World Cancer Research Fund; American Institute for Cancer Research, 2018.
15. Kyrgiou M, Kalliala I, Markozannes G, et al. Adiposity and cancer at major anatomical sites: umbrella review of the literature. *Bmj* 2017;356
16. Azizpour Y, Delpisheh A, Montazeri Z, et al. Effect of childhood BMI on asthma: a systematic review and meta-analysis of case-control studies. *BMC Pediatr* 2018;18(1):1-13.
17. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med* 2007;175(7):661-66.
18. Shiri R, Karppinen J, Leino-Arjas P, et al. The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
19. Silverwood V, Blagojevic-Bucknall M, Jinks C, et al. Current evidence on risk factors for knee osteoarthritis in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage* 2015;23(4):507-15.
20. Jiang L, Rong J, Wang Y, et al. The relationship between body mass index and hip osteoarthritis: a systematic review and meta-analysis. *Joint Bone Spine* 2011;78(2):150-55.
21. Amiri S, Behnezhad S, Nadinlui KB. Body Mass Index (BMI) and risk of depression in adults: A systematic review and meta-analysis of longitudinal studies. *Obesity Medicine* 2018;12:1-12.
22. Peters SA, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia* 2014;57(8):1542-51.
23. Peters SA, Huxley RR, Woodward M. Diabetes as a risk factor for stroke in women compared with men: a systematic review and meta-analysis of 64 cohorts, including 775 385 individuals and 12 539 strokes. *Lancet* 2014;383(9933):1973-80.
24. Sullivan PW, Slejko JF, Sculpher MJ, et al. Catalogue of EQ-5D scores for the United Kingdom. *Med Decis Making* 2011;31(6):800-04.
25. NHS Digital. Hospital Episode Statistics for England: Admitted Patient Care statistics, 2019-20 2020 [Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2019-20> accessed 21 April 2021.