

# Association of maternal air pollution exposure and infant lung function is modified by genetic propensity to oxidative stress.

**Authors:** Dwan Vilcins<sup>1</sup>, Wen Ray Lee<sup>1</sup>, Cindy Pham<sup>2,3</sup>, Sam Tanner, Luke D. Knibbs<sup>4</sup>, Dave Burgner<sup>5</sup>, Tamara Blake<sup>1</sup>, Toby Mansell, Anne-Louise Ponsonby<sup>6</sup>, Peter D Sly<sup>1</sup>, Barwon Infant Study Investigator group.

## **Affiliations:**

<sup>1</sup> Child Health Research Centre, The University of Queensland, South Brisbane, QLD 4101

<sup>2</sup> Murdoch Children's Research Institute, Royal Children's Hospital, Parkville VIC 3052

<sup>3</sup> Melbourne School of Population and Global Health, University of Melbourne, Parkville VIC 3052

<sup>5</sup> Murdoch Children's Research Institute, Royal Children's Hospital, The University of Melbourne; Melbourne, Australia.

<sup>6</sup> The Florey Institute of Neuroscience and Mental Health, Murdoch Children's Research Institute

## Supplementary materials

### 1 Directed acyclic graph




Figure 1: Directed acyclic graph depicted proposed relationships exposure, outcome and covariates. Green shapes and arrows indicate exposure, blue shapes indicate outcome and its antecedents, including potential mediators. White shapes indicate variables which were included as covariates, while grey shapes are variables proposed to have a relationship on the causal network, but which are unobserved in our models. Maternal TRAP refers to maternal exposure to traffic related air pollution.

## 2 Summary of exposure, outcome and mediators

Table S1: Summary of exposures and outcomes in the Barwon Infant Study

Variable	mean	sd	median	min	max	IQR
FRC (mL / kg)	87.7	14.9	86.7	52.6	144.0	4.27
LCI (turnovers)	6.79	0.43	6.79	5.410	8.2	0.52
Maternal NO <sub>2</sub> (ppb)	5.58	1.97	5.44	1.983	13.9	2.27
Maternal PM <sub>2.5</sub> (µg/m <sup>3</sup> )	7.50	0.66	7.42	5.932	9.9	0.88
Maternal 8-OHdG (ng/mL) (log)	1.3	0.8	1.3	-1.3	4.1	0.94
Maternal 8-OHGua (ng/mL) (log)	2.9	1.1	3.0	-0.7	6.5	1.09
gPFSox	6.8	0.9	7.00	4.5	9.5	1.5

## 3 Sensitivity analysis

### 3.1 SES

Adding socio-economic status to the model (using maternal education level as a proxy measure) of FRC did not significantly change any of the estimates and was not itself significant ( $\beta=0.6$  mls, 95% CI (-1.7, 2.8),  $p=0.63$ ).

### 3.2 Air pollution and maternal smoking interaction (adjusted models)

Table S2: Estimates for lung function at 4 weeks of age (FRC) for maternal exposure to PM<sub>2.5</sub> and the interaction with maternal smoking in the Barwon Infant Study

<i>Predictors</i>	<b>median FRC</b>		
	<i>Estimates</i>	<i>CI</i>	<i>p</i>
Maternal PM <sub>2.5</sub> in year before birth (ug/m3)	1.2	-1.5 – 4.0	0.374
PM <sub>2.5</sub> * maternal smoking	1.6	-5.2 – 8.3	0.650

Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking

### 3.3 Child's passive smoke exposure (adjusted)

Table S3: Estimates for lung function at 4 weeks of age (LCI) for maternal exposure to PM<sub>2.5</sub>, maternal smoking, and a child's exposure to environmental tobacco smoke at 4 weeks of age in the Barwon Infant Study

<i>Predictors</i>	<b>median LCI</b>		
	<i>Estimates</i>	<i>CI</i>	<i>p</i>
PM <sub>2.5</sub> (ug/m3)	-0.03	-0.10 – 0.05	0.446
Maternal smoking	-0.04	-0.17 – 0.10	0.612
Child passive smoke exposure	0.01	-0.29 – 0.31	0.947

Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking

Table S4: Estimates for lung function at 4 weeks of age (LCI) for maternal exposure to NO<sub>2</sub>, maternal smoking, and a child's exposure to environmental tobacco smoke at 4 weeks of age in the Barwon Infant Study

<i>Predictors</i>	<b>median LCI</b>		
	<i>Estimates</i>	<i>CI</i>	<i>p</i>
NO <sub>2</sub> (ppb)	0.01	-0.02 – 0.03	0.580
Maternal smoking	-0.03	-0.17 – 0.10	0.619
Child passive smoke exposure	0.02	-0.27 – 0.32	0.871

Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking

### 3.4 Child's exposure to PM<sub>2.5</sub> and lung function (adjusted)

Table S5: Effect estimates for lung function at 4 weeks of age (FRC) for a child's exposure to PM<sub>2.5</sub> at different timepoints, including an interaction with the oxidative stress genotype, in the Barwon Infant Study.

<i>Predictors</i>	<b>median FRC</b>		
	<i>Estimates</i>	<i>CI</i>	<i>p</i>
PM <sub>2.5</sub> (mcg/m <sup>3</sup> ) at birth	1.2	-1.3 – 3.6	0.339
PM <sub>2.5</sub> mcg/m <sup>3</sup> ) at 1 month	0.6	-1.8 – 3.0	0.596
PM <sub>2.5</sub> mcg/m <sup>3</sup> ) at 1 month * oxidative stress genotype	4.1	-1.9 – 10.0	0.178

Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking

Table S6: Effect estimates for lung function at 4 weeks of age (FRC) for a child's exposure to NO<sub>2</sub> at different timepoints, including an interaction with the oxidative stress genotype, in the Barwon Infant Study.

<i>Predictors</i>	<b>median LCI</b>		
	<i>Estimates</i>	<i>CI</i>	<i>p</i>
PM <sub>2.5</sub> (mcg/m <sup>3</sup> ) at birth	0.002	-0.069 – 0.072	0.959
PM <sub>2.5</sub> (mcg/m <sup>3</sup> ) at 1 month	0.012	-0.058 – 0.081	0.742
PM <sub>2.5</sub> (mcg/m <sup>3</sup> ) at 1 month * oxidative stress genotype	-0.073	-0.246 – 0.099	0.404

Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking

### 3.5 Air pollutants in tertiles

Table S7: Results of linear regression models of maternal exposure to air pollution and infant lung function at 4 weeks of age in the Barwon Infant Study, sensitivity using air pollution variable in tertiles

<b>Whole cohort</b>				
		<i>Estimates</i>	<i>CI</i>	<i>p</i>
<b>LCI</b> <i>n</i> =275	Medium PM <sub>2.5</sub>	-0.02	-0.14 – 0.10	0.78
	High PM <sub>2.5</sub>	-0.02	-0.14 – 0.10	0.73
	Medium NO <sub>2</sub>	-0.09	-0.21 – 0.03	0.13
	High NO <sub>2</sub>	-0.01	-0.13 – 0.11	0.91
<b>FRC</b> <i>n</i> =275	Medium PM <sub>2.5</sub>	1.33	-2.81 – 5.47	0.53
	High PM <sub>2.5</sub>	3.14	-1.04 – 7.32	0.14
	Medium NO <sub>2</sub>	2.58	-1.58 – 6.73	0.22
	High NO <sub>2</sub>	1.34	-2.85 – 5.53	0.53
<b>tPTEF/tE</b> <i>n</i> =355	Medium PM <sub>2.5</sub>	-0.01	-0.03 – 0.02	0.47
	High PM <sub>2.5</sub>	0.00	-0.02 – 0.03	0.92
	Medium NO <sub>2</sub>	-0.02	-0.05 – 0.00	0.10
	High NO <sub>2</sub>	0.001	-0.03 – 0.03	0.97
<b>Children at high risk of oxidative stress</b>				
		<i>Estimates</i>	<i>CI</i>	<i>p</i>
<b>LCI</b> <i>n</i> =48	Medium PM <sub>2.5</sub>	0.14	-0.17 – 0.46	0.36
	High PM <sub>2.5</sub>	-0.02	-0.34 – 0.30	0.91
	Medium NO <sub>2</sub>	0.22	-0.11 – 0.54	0.19
	High NO <sub>2</sub> <sup>*</sup>	0.46	0.10 – 0.82	<b>0.01</b>
<b>FRC</b> <i>n</i> =48	Medium PM <sub>2.5</sub>	0.6	-11.0 – 12.1	0.92
	High PM <sub>2.5</sub>	-0.6	-12.2 – 11.1	0.92
	Medium NO <sub>2</sub>	-0.9	-12.6 – 10.8	0.88
	High NO <sub>2</sub> <sup>^</sup>	-13.1	-26.1 – -0.1	<b>0.05</b>
<b>tPTEF/tE</b> <i>n</i> =65	Medium PM <sub>2.5</sub>	-0.04	-0.10 – 0.03	0.31
	High PM <sub>2.5</sub>	-0.02	-0.09 – 0.05	0.64
	Medium NO <sub>2</sub>	-0.03	-0.10 – 0.03	0.33
	High NO <sub>2</sub>	0.02	-0.05 – 0.09	0.51
Models adjusted for sex, birthweight, maternal age, maternal pre-pregnancy BMI and maternal smoking				
* Estimates significantly different to model of low risk OS kids <i>p</i> =0.007				
^ Estimates significantly different to model of low risk OS kids <i>p</i> =0.02				