

Plasma GPI and PGD are associated with vascular normalization and may serve as novel prognostic biomarkers for lung adenocarcinoma: Multi-omics and multi-dimensional analysis

Yiran Liu^{1,*}, Yanchi Wang^{2,*}, Liping Mao^{2,*}, Yang Hu³, Rongrong Zhao⁴, Wendi Zhang¹,
Huiwen Xu¹, Buyun Xu¹, Yutong Wu¹, Qiong Chen¹, Xiaobo Tao¹, Shufan Xu¹, Xun Zhuang¹,
Yulong Lian¹, Yihua Lu¹, Lei Zhang¹, Tian Tian¹, Jiahua Cui¹, Guangyu Tian^{4,#}, Minjie
Chu^{1,#}

¹Department of Epidemiology, School of Public Health, Nantong University, Nantong, Jiangsu, China

²Affiliated Nantong Hospital of Shanghai University (The Sixth People's Hospital of Nantong), Nantong, Jiangsu, China

³Department of nutrition, Hai 'an City People's Hospital, Nantong, Jiangsu, China

⁴Department of Oncology, Jiangdu People's Hospital of Yangzhou, Yangzhou, Jiangsu, China

[#]Correspondence:

Minjie Chu

Department of Epidemiology, School of Public Health, Nantong University, Nantong, Jiangsu, China, 226019

E-mail: chuminjie@ntu.edu.cn

Guangyu Tian

Department of Oncology, Jiangdu People's Hospital of Yangzhou, Yangzhou, Jiangsu,

China, 225202

E-mail: jdryzlzx@hotmail.com

The authors have withdrawn their manuscript owing to adjustments have been made to the content. Therefore, the authors do not wish this work to be cited as reference for the project. If you have any questions, please contact the corresponding author.