

Table 1. Sensory properties and overall taste for active and placebo powders across concentrations

Active Solutions				
Sensory Property of Samples	S1 (1 g/L) (N=35)	S2 (2g/L) (N=35)	S3 (3 g/L) (N=35)	p-value
Odour	8 (6-9) ^a	8 (5-9) ^{ab}	6 (4-8) ^c	p<0.05
Colour	9 (8-9) ^a	9 (8-9) ^{ab}	8 (7-9) ^c	p<0.05
Salty	8 (6-9) ^a	5 (3-8) ^b	4 (3-6) ^c	p<0.05
Sweet	8 (7-9) ^a	7 (6-9) ^{ab}	6 (4-9) ^c	p=0.003
Bitter	8 (5-9) ^a	5 (4-8) ^b	4 (3-6) ^c	p<0.05
Umami	9 (6-9)	8 (5-9)	8 (4-9)	p=0.169
Sour	9 (8-9) ^a	6 (4-9) ^b	4 (3-5) ^c	p<0.05
Placebo Solutions				
Sensory Property of Samples	S4 (1 g/L) (N=35)	S5 (2g/L) (N=35)	S6 (3 g/L) (N=35)	p-value
Odour	8 (5-9) ^a	6 (4-8) ^a	4 (3-6) ^b	p<0.05
Colour	8 (7-9) ^a	6 (4-8) ^b	4 (3-7) ^c	p<0.05
Salty	6 (4-7) ^a	4 (2-5) ^b	2 (2-4) ^c	p<0.05
Sweet	7 (6-9) ^a	6 (4-7) ^b	6 (3-8) ^b	p=0.003
Bitter	7 (4-9) ^a	4 (2-5) ^b	3 (2-5) ^b	p<0.05
Umami	7 (4-9)	5 (4-7)	4 (2-6)	p=0.07
Sour	7 (4-9) ^a	4 (3-5) ^b	3 (1-4) ^c	p<0.05

Hedonic scores of participants' (N=35) evaluation of each sensory property compared within groups for active samples (S1, S2, S3) and placebo samples (S4, S5, S6) at concentrations of 1g/L, 2g/L, and 3g/L. Hedonic scores range from 1-9, where a score of 1=Disliked extremely, 9=Liked extremely, and 5=Neither like nor dislike. Data are median (IQR). Groups with different letters are significantly different (Friedmans two-way analysis of variance by ranks test and Wilcoxon signed-rank test with Bonferroni correction for multiple comparisons, p<0.05).