

1 SUPPLEMENTARY MATERIAL

2 **The spread of a validated molecular marker of artemisinin partial resistance *pfkelch13***

3 **R622I and association with *pfhrp2/3* deletions in Eritrea**

4 Selam Mihreteab¹, Karen Anderson^{2,3}, Irene Molina - de la Fuente^{4,5}, Colin J. Sutherland⁶,
5 David Smith^{2,3}, Jane Cunningham⁷, Khalid B Beshir^{6*}, Qin Cheng^{2,3*}

6
7 ¹National Malaria Control Program, Ministry of Health, Asmara, Eritrea; ²Drug Resistance
8 and Diagnostics, Australian Defence Force Malaria and Infectious Disease Institute, Brisbane,
9 Australia; ³QIMR-Berghofer Medical Research Institute, Brisbane, Australia; ⁴ Carlos III
10 Institute of Health, Madrid, Spain; ⁵University of Alcalá, Madrid, Spain; ⁶Faculty of Infectious
11 Diseases, London School of Hygiene & Tropical Medicine, UK; ⁷Global Malaria Programme,
12 World Health Organization, Geneva, Switzerland.

13 *Corresponding authors

14 Dr Qin Cheng, Australian Defence Force Malaria and Infectious Disease Institute, Weary
15 Dunlop Drive, Gallipoli Barracks, Enoggera, QLD 4051, Australia. email:

16 qin.cheng@defence.gov.au.

17 Dr Khalid B Beshir, Faculty of Infectious Diseases, London School of Hygiene & Tropical
18 Medicine, UK. Email: Khalid.Beshir@lshtm.ac.uk.

19

20

Supplemental Table 1. Number of samples sequenced for *pfk13*, mutations identified and mutation frequency in different health facilities.

Zone	Health Centre	No. sequenced	E605K		R622I		N657K		K658E		S679L		WT	
			n	%	n	%	n	%	n	%	n	%	n	%
Gash Barka	Agordat	95	0	0.0%	13	13.7%	1	1.1%	0	0.0%	0	0.0%	81	85.3%
	Shambuko	61	1	1.6%	7	11.5%	0	0.0%	1	1.6%	0	0.0%	52	85.2%
	Tesseney	61	0	0.0%	7	11.5%	0	0.0%	0	0.0%	0	0.0%	54	88.5%
	Tokombia	62	0	0.0%	7	11.3%	0	0.0%	0	0.0%	0	0.0%	55	88.7%
	Barentu	91	0	0.0%	10	11.0%	1	1.1%	0	0.0%	0	0.0%	80	87.9%
	<i>Subtotal</i>	<i>370</i>	<i>1</i>	<i>0.3%</i>	<i>44</i>	<i>11.9%</i>	<i>2</i>	<i>0.5%</i>	<i>1</i>	<i>0.3%</i>	<i>0</i>	<i>0.0%</i>	<i>322</i>	<i>87.0%</i>
Anseba	Keren	91	0	0.0%	9	9.9%	0	0.0%	0	0.0%	0	0.0%	82	90.1%
	Hagaz	25	0	0.0%	7	28.0%	0	0.0%	0	0.0%	0	0.0%	18	72.0%
	<i>Subtotal</i>	<i>116</i>	<i>0</i>	<i>0.0%</i>	<i>16</i>	<i>13.8%</i>	<i>0</i>	<i>0.0%</i>	<i>0</i>	<i>0.0%</i>	<i>0</i>	<i>0.0%</i>	<i>100</i>	<i>86.2%</i>
Debub	Mendefera	51	0	0.0%	3	5.9%	0	0.0%	0	0.0%	0	0.0%	48	94.1%
	Maymine	40	0	0.0%	5	12.5%	0	0.0%	0	0.0%	1	2.5%	34	85.0%
	A/D/E/M	10	0	0.0%	2	20.0%	0	0.0%	0	0.0%	0	0.0%	8	80.0%
	<i>Subtotal</i>	<i>101</i>	<i>0</i>	<i>0.0%</i>	<i>10</i>	<i>9.9%</i>	<i>0</i>	<i>0.0%</i>	<i>0</i>	<i>0.0%</i>	<i>1</i>	<i>1.0%</i>	<i>90</i>	<i>89.1%</i>
Total		587	1	0.2%	70	11.9%	2	0.3%	1	0.2%	1	0.2%	512	87.2%

Supplemental Table 2. Haplotypes observed among 26 samples carrying *pfk13* R622I mutation.


Zone	Location	SAMPLE ID	TA1	313	2490	383	POLY A	PK2	TA109	Haplo type
Gash Barka	Agordat	EA04	165	229	83	137	152	162	160	1
	Agordat	EA02	165	233	83	137	152	162	160	2
	Agordat	EA28	165	233	83	137	152	162	160	2
	Agordat	EA66	165	233	83	137	158	162	151	3
	Barentu	EB1988	159	253	83	125	152	162	172	4
	Barentu	EB2254	165	219	83	125	158	162	160	5
	Barentu	EB1793	165	219	83	127	158	171	160	6
	Barentu	EB1986	165	237	83	127	167	171	175	7
	Shambuko	ES65	165	233	83	137	152	162	151	8
	Shambuko	ES26	165	233	83	137	152	171	151	9
	Shambuko	ES61	165	235	83	137	158	162	175	10
	Tesseney	ETHOS03	165	229	83	137	158	162	151	11
	Tesseney	ETHOS30	165	247	83	136	152	162	154	12
	Tokombia	ETHC59	159	229	83	137	152	162	151	12
	Tokombia	ETHC60	165	233	83	137	173	162	175	14
Tokombia	ETHC42	165	235	83	137	158	162	165	15	
Anseba	Hagaz	EH07	165	229	83	137	149	162	160	16
	Hagaz	EH23	165	233	83	127	167	171	160	17
	Hagaz	EH04	165	233	83	137	170	162	163	18
	Hagaz	EH20	165	233	83	137	173	162	175	14
	Hagaz	EH29	165	233	83	137	173	162	175	14
	Hagaz	EH17	165	249	83	137	152	162	163	19
	Keren	EK67	165	233	83	137	152	162	163	20
Debut	Maimine	EMA02	159	233	83	127	176	186	172	21
	Maimine	EMA33	165	229	83	137	155	186	175	22
	Mendefera	EME31	165	229	83	137	158	162	160	23

Supplemental Table 3. Number of unique alleles and genetic diversity detected per microsatellite locus and population.

Microsatellite marker	Number of alleles		Gene diversity H_E per locus	
	Mutant	Wild type	Mutant	Wild type
TA1	2	8	0.212	0.647
313	8	8	0.745	0.542
2490	1	5	0.000	0.167
383	3	9	0.394	0.455
PolyA	8	13	0.785	0.828
PK2	3	9	0.394	0.674
TA109	7	10	0.812	0.728
Total	23	87	Mean (95% CI)	Mean (95% CI)
			0.4774 (0.1873 - 0.7676)	0.5773 (0.3760 - 0.7786)
P = 0.0500				

Supplemental Table 4. Number and proportion of samples with different *pfhrp2/3* status.

Zone	Health Centre	No. Tested	Single <i>pfhrp2</i> del		Single <i>pfhrp3</i> del		Dual <i>pfhrp2/3</i> del		No del	
			n	%	n	%	n	%	n	%
Gash Barka	Agordat	93	4	4.3%	43	46.2%	28	30.1%	18	19.4%
	Shambuko	59	1	1.7%	26	44.1%	18	30.5%	14	23.7%
	Tesseney	61	2	3.3%	25	41.0%	10	16.4%	24	39.3%
	Tokombia	62	1	1.6%	24	38.7%	11	17.7%	26	41.9%
	Barentu	90	3	3.3%	43	47.8%	21	23.3%	23	25.6%
	<i>Subtotal</i>	<i>365</i>	<i>11</i>	<i>3.0%</i>	<i>161</i>	<i>44.1%</i>	<i>88</i>	<i>24.1%</i>	<i>105</i>	<i>28.8%</i>
Anseba	Keren	89	5	5.6%	42	47.2%	26	29.2%	16	18.0%
	Hagaz	25	0	0.0%	11	44.0%	6	24.0%	8	32.0%
	<i>Subtotal</i>	<i>114</i>	<i>5</i>	<i>4.4%</i>	<i>53</i>	<i>46.5%</i>	<i>32</i>	<i>28.1%</i>	<i>24</i>	<i>21.1%</i>
Debub	Mendefera	51	0	0.0%	17	33.3%	19	37.3%	15	29.4%
	Maimine	38	0	0.0%	20	52.6%	3	7.9%	15	39.5%
	A/D/E/M	10	1	10.0%	4	40.0%	3	30.0%	2	20.0%
	<i>Subtotal</i>	<i>99</i>	<i>1</i>	<i>1.0%</i>	<i>41</i>	<i>41.4%</i>	<i>25</i>	<i>25.3%</i>	<i>32</i>	<i>32.3%</i>
Total		578	17	2.9%	255	44.1%	145	25.1%	161	27.9%



Supplementary Figure 1. Prevalence K13 mutations in samples with different *pfhrp2/3* status (A) and proportion of *pfhrp2/3*-deleted parasites among K13 R622I mutant and 622R wild type (B).