

Supplementary Materials.

Supplementary Methods.

Linking pharmacy claims with outpatient visits. Outpatient claims included up to two diagnoses coded according to the International Classification of Diseases (ICD). We aggregated these diagnoses into broader classes (“conditions”) using mappings provided by the Agency for Healthcare Research and Quality’s (AHRQ’s) Clinical Classification Software (CCS). To identify individual outpatient visits, we extracted the first outpatient encounter associated with each person on a given day. Frequently, there are many claims on a given day associated with a single encounter, capturing various procedures that are all associated with a common diagnosis or set of diagnoses. Then, for each person, we extracted all pharmacy claims and restricted to antibiotics on the basis of National Drug Code (NDC) codes. For each antibiotic claim, we found the most recent outpatient visit that occurred for that person within the preceding seven days (including the day of prescription fill). This constituted a “link” between the prescription and an outpatient visit. If there was no outpatient visit in the seven days preceding the prescription fill, the prescription was marked as “unlinked”, affecting less than 15% of prescriptions. Mappings from ICD diagnoses to CCS conditions and from NDC codes to antibiotics are included in the online code repository associated with this manuscript: <https://github.com/gradlab/AbxGeography>.

Sensitivity analysis of all conditions. As in the main analysis, we also measured the relative contribution of outpatient visit rates and per-visit prescribing to the geographic variation in antibiotic prescribing rates, by calculating the coefficient of determination (R^2) for the regression of antibiotic prescriptions per person-year on (a) outpatient visits per person-year and (b) antibiotic prescriptions per visit, with higher R^2 indicating a greater amount of variation explained (Supplementary Figure S1). Additionally, we measured how the amount of variation explained by outpatient visit rates and per-visit prescribing rates was related to the total prescribing volume across all conditions in the dataset. Specifically, we conducted an additional regression of the R^2 values for each of the two variables against the \log_{10} -transformed total volume of antibiotic prescriptions associated with each condition, to assess whether the explanatory power of outpatient visits and per-visit prescribing rates varied systematically by the total volume of antibiotics prescribed for a given condition (Supplementary Figure S2).

Supplementary Table 1. Characteristics of the study population and reference values from the 2020 Census

Variable	Category	Number of people (%) by year				Reference values
		2015	2016	2017	2018	2020 Census
Total	–	18989691 (100%)	18897347 (100%)	15574072 (100%)	16419151 (100%)	
Sex	Male	9139343 (48.1%)	9113395 (48.2%)	7594325 (48.8%)	8061316 (49.1%)	49.1%
	Female	9850348 (51.9%)	9783952 (51.8%)	7979747 (51.2%)	8357835 (50.9%)	50.9%
Census Region	Northeast	3401805 (17.9%)	3071460 (16.3%)	2351782 (15.1%)	3173820 (19.3%)	17.2%
	South	8393206 (44.2%)	8679832 (45.9%)	6671373 (42.8%)	6425726 (39.1%)	38.4%
	Midwest	3916502 (20.6%)	3862332 (20.4%)	3863799 (24.8%)	3965199 (24.1%)	20.7%
	West	3278178 (17.3%)	3283723 (17.4%)	2687118 (17.3%)	2854406 (17.4%)	23.7%
Age group (years)	0-4	1018741 (5.4%)	1023975 (5.4%)	866737 (5.6%)	902236 (5.5%)	6.0%
	5-9	1244135 (6.6%)	1217948 (6.4%)	1008299 (6.5%)	1041415 (6.3%)	6.2%
	10-14	1413095 (7.4%)	1392233 (7.4%)	1164661 (7.5%)	1215861 (7.4%)	6.4%
	15-19	1526902 (8.0%)	1376856 (7.3%)	1143650 (7.3%)	1191431 (7.3%)	6.4%
	20-24	1607390 (8.5%)	1589339 (8.4%)	1315453 (8.4%)	1385926 (8.4%)	6.5%
	25-29	1037819 (5.5%)	1107728 (5.9%)	937160 (6.0%)	995527 (6.1%)	7.1%
	30-34	1316622 (6.9%)	1313278 (6.9%)	1108050 (7.1%)	1179238 (7.2%)	6.8%
	35-39	1419642 (7.5%)	1436661 (7.6%)	1212464 (7.8%)	1298883 (7.9%)	6.6%
	40-44	1588943 (8.4%)	1508533 (8.0%)	1245861 (8.0%)	1309912 (8.0%)	6.1%
	45-49	1673758 (8.8%)	1683088 (8.9%)	1394427 (9.0%)	1464421 (8.9%)	6.0%
	50-54	1876276 (9.9%)	1819797 (9.6%)	1456097 (9.3%)	1522409 (9.3%)	6.1%
	55-59	1880816 (9.9%)	1883292 (10%)	1509362 (9.7%)	1605771 (9.8%)	6.5%
	60-64	1385552 (7.3%)	1544619 (8.2%)	1211851 (7.8%)	1306121 (8.0%)	6.5%










Supplementary Table 2. Prioritization scheme for assigning antibiotic prescriptions to outpatient visits. If a person had multiple diagnoses associated with the most recent visit preceding an antibiotic prescription, the condition with the highest rank in this table was marked as the condition associated with that prescription. For streptococcal pharyngitis and sinusitis (in CCD9), there is no corresponding CCS condition, so these were extracted explicitly using ICD codes. Asterisks denote a wildcard symbol, so that ICD code 461* contains all ICD codes that begin with 461 and end with a single additional number.

Condition	CCS9 code	CCS10 code
Pneumonia	122	RSP002
UTI	159	GEN004
Intestinal infection	135	DIG001
Strep pharyngitis	ICD= 4610	ICD= J020
Sinusitis	ICD=461*, 473*	RSP001
URI (other)	126	RSP006
SSTI	197	SKN001
Otitis media	92	EAR001
Asthma	128	RSP009
Bacterial infection (other)	3	INF003
Tonsillitis	124	RSP004
Bronchitis (acute)	125	RSP005
Influenza	123	RSP003
Viral infection	7	INF008
Essential hypertension	98	CIR007

Supplementary Table 3. Antibiotic prescribing rates associated with various medical conditions.

Condition	Antibiotic prescriptions per 1,000 people	Percent of antibiotic prescriptions nationally	Visits per 1,000 people	Percent of visits with an antibiotic prescription
Sinusitis	97.1	19.3	139.0	69.9
URI (other)	71.5	14.2	208.1	34.4
Otitis media	44.7	8.9	75.8	59.0
Strep pharyngitis	26.6	5.3	32.2	82.5
Bronchitis (acute)	24.0	4.8	41.8	57.5
SSTI	19.7	3.9	54.9	35.9
UTI	17.7	3.5	58.3	30.4
Pneumonia	7.5	1.5	17.9	42.2
Essential hypertension	6.9	1.4	241.4	2.9
Asthma	5.6	1.1	60.7	9.1
Bacterial infection (other)	5.3	1.0	16.1	33.0
Tonsillitis	4.7	0.9	13.1	35.8
Viral infection	3.3	0.7	85.6	3.8
Influenza	1.4	0.3	16.5	8.3
Intestinal infection	1.3	0.2	11.5	11.0
Motor vehicle traffic	0	0	0.2	1.3
Other	164.7	32.8	6881.1	2.4
Total	502.1	100	7954.2	6.3

Supplementary Figure 1. Variation in outpatient visits, per-visit prescribing, and per capita antibiotic prescribing rates across the US.



Supplementary Figure 2. Fraction of the variation in *per capita* antibiotic prescribing explained by per-visit prescribing and visits. Lines depict the least-squares linear regression between per-visit prescriptions (blue, panel A) and outpatient visits (red, panel B) against the \log_{10} -transformed volume of antibiotic prescriptions per 1000 people associated with each condition.

