

Supplementary Table 1.

Discovery GWASs of psychiatric diseases, behavioral and related traits used to construct the GPSs

Trait	n cases	n controls	n total ^a	ref
Attention-Deficit Hyperactivity Disorder	20,183	35,191	51,306	[1]
Anxiety	16,730	101,021	57,412	http://www.nealelab.is/uk-biobank/
Bipolar Disorder	41,917	371,549	150,670	[2]
Depression	170,756	329,443	449,856	[3]
Post-Traumatic Stress Disorder	9,354	25,175	27,280	[4]
Schizophrenia	67390	94015	157,013	[5]
Risk tolerance	-	-	466,571	[6]
Suicide attempt	26,590	492,022	100,907	[7]
Educational attainment	-	-	766,345	[8]
Well being	-	-	2,083,151	[9]

^a For binary traits effective sample size was calculated with the formula $4/(1/n \text{ cases} + 1/n \text{ controls})$

Supplementary Table 2a. Association between the GPS for ADHD and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,03 , -3,00E-04	4,53E-02
Age at onset of SUD ^c	1280	Linear	-0,01	-0,02 , 0,01	0,39
Years between substance use and SUD	1280	Linear	1,04	0,95 , 1,14	0,36
Years of substance use as proportion of lifespan	1199	Linear	1,07	0,03 , 2,1	4,42E-02
Number of substances consumed	879	Negative binomial	1,03	0,99 , 1,07	0,19
Number of therapeutic community interventions	1345	Negative binomial	1,01	0,92 , 1,11	0,80
Number of inpatient detoxifications	1327	Negative binomial	1,03	0,92 , 1,15	0,62
Number of outpatient treatments	1252	Negative binomial	1,05	0,99 , 1,11	0,09
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,12	0,92 , 1,36	0,28
Major depressive disorder	828	Logistic	1,10	0,96 , 1,27	0,17
Antisocial personality disorder	560	Logistic	1,24	1,04 , 1,47	1,60E-02
Psychotic disorder	593	Logistic	1,02	0,79 , 1,3	0,90
Anxiety disorder	654	Logistic	1,03	0,88 , 1,2	0,71
Attention deficit hyperactivity disorder	700	Logistic	1,24	1,06 , 1,45	7,35E-03
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	0,07	-0,29 , 0,43	0,72
Aggression Hostility personality factor	667	Linear	0,15	-0,08 , 0,38	0,21
Sociability personality factor	632	Linear	-0,18	-0,44 , 0,08	0,18
Impulsive sensation seeking personality factor	666	Linear	0,01	-0,3 , 0,32	0,94
Activity personality factor	665	Linear	0,02	-0,24 , 0,28	0,88
Suicide attempt	618	Logistic	1,03	0,89 , 1,19	0,67
Suicide ideation	731	Logistic	0,98	0,84 , 1,15	0,80
Psychotic symptoms	1281	Logistic	1,00	0,89 , 1,12	0,98
Sleeping disturbances	1228	Logistic	1,02	0,91 , 1,14	0,76

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,00	0,87 , 1,16	0,97
Employment status	984	Ordinal	1,08	0,97 , 1,2	0,18
Medical status	982	Ordinal	1,03	0,92 , 1,17	0,60
Psychiatric status	984	Ordinal	1,01	0,91 , 1,13	0,86
Drug use	984	Ordinal	1,06	0,95 , 1,19	0,28
Alcohol use	984	Ordinal	0,96	0,86 , 1,08	0,54
Family/Social relationships	981	Ordinal	1,10	0,99 , 1,23	0,09
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,52	-1,23 , 0,19	0,15
Mental health	751	Linear	0,49	-0,45 , 1,44	0,31
Criminal record	713	Logistic	1,13	0,98 , 1,32	0,10
Unemployment	1057	Logistic	1,03	0,91 , 1,17	0,61
Number of psychiatric hospitalizations	760	Negative binomial	1,05	0,86 , 1,29	0,61
Psychiatric family history	840	Logistic	0,97	0,85 , 1,11	0,66
Lifetime medical conditions	1334	Logistic	0,99	0,88 , 1,11	0,84
Substance use family history	818	Logistic	1,10	0,96 , 1,27	0,16
Educational attainment	1333	Ordinal	0,85	0,93 , 0,77	1,20E-03

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2b. Association between the GPS for anxiety and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,02 , 0,01	0,36
Age at onset of SUD ^c	1280	Linear	0,00	-0,02 , 0,01	0,77
Years between substance use and SUD	1280	Linear	1,04	0,95 , 1,13	0,45
Years of substance use as proportion of lifespan	1199	Linear	-0,03	-1,07 , 1,02	0,96
Number of substances consumed	879	Negative binomial	1,01	0,97 , 1,05	0,79
Number of therapeutic community interventions	1345	Negative binomial	1,10	1 , 1,21	0,05
Number of inpatient detoxifications	1327	Negative binomial	1,01	0,9 , 1,13	0,84
Number of outpatient treatments	1252	Negative binomial	1,09	1,03 , 1,15	2,67E-03
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,02	0,84 , 1,24	0,86
Major depressive disorder	828	Logistic	1,03	0,9 , 1,19	0,63
Antisocial personality disorder	560	Logistic	1,09	0,92 , 1,29	0,30
Psychotic disorder	593	Logistic	1,49	1,16 , 1,92	1,93E-03
Anxiety disorder	654	Logistic	1,04	0,89 , 1,21	0,63
Attention deficit hyperactivity disorder	700	Logistic	0,86	0,74 , 1,01	0,06
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	0,23	-0,12 , 0,59	0,20
Aggression Hostility personality factor	667	Linear	0,07	-0,16 , 0,3	0,56
Sociability personality factor	632	Linear	-0,08	-0,34 , 0,17	0,53
Impulsive sensation seeking personality factor	666	Linear	0,02	-0,29 , 0,33	0,91
Activity personality factor	665	Linear	0,04	-0,22 , 0,29	0,79
Suicide attempt	618	Logistic	1,07	0,93 , 1,23	0,37
Suicide ideation	731	Logistic	1,01	0,87 , 1,18	0,88
Psychotic symptoms	1281	Logistic	1,06	0,95 , 1,19	0,28
Sleeping disturbances	1228	Logistic	0,98	0,88 , 1,1	0,77

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	0,92	0,8 , 1,07	0,28
Employment status	984	Ordinal	0,97	0,87 , 1,08	0,57
Medical status	982	Ordinal	1,08	0,96 , 1,22	0,23
Psychiatric status	984	Ordinal	1,08	0,97 , 1,2	0,16
Drug use	984	Ordinal	1,08	0,97 , 1,21	0,17
Alcohol use	984	Ordinal	0,95	0,85 , 1,06	0,36
Family/Social relationships	981	Ordinal	1,07	0,96 , 1,19	0,26
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,93	-1,64 , -0,22	1,07E-02
Mental health	751	Linear	0,06	-0,89 , 1,01	0,90
Criminal record	713	Logistic	0,94	0,81 , 1,1	0,47
Unemployment	1057	Logistic	1,05	0,93 , 1,19	0,44
Number of psychiatric hospitalizations	760	Negative binomial	1,03	0,84 , 1,27	0,77
Psychiatric family history	840	Logistic	1,16	1,01 , 1,33	4,16E-02
Lifetime medical conditions	1334	Logistic	1,06	0,95 , 1,19	0,31
Substance use family history	818	Logistic	1,02	0,88 , 1,17	0,80
Educational attainment	1333	Ordinal	0,92	1,02 , 0,83	0,10

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2c. Association between the GPS for bipolar disorder and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	0,01	-0,01 , 0,02	0,33
Age at onset of SUD ^c	1280	Linear	0,00	-0,02 , 0,02	0,90
Years between substance use and SUD	1280	Linear	1,01	0,93 , 1,11	0,78
Years of substance use as proportion of lifespan	1199	Linear	0,20	-0,85 , 1,24	0,71
Number of substances consumed	879	Negative binomial	1,02	0,98 , 1,07	0,26
Number of therapeutic community interventions	1345	Negative binomial	0,92	0,84 , 1,01	0,09
Number of inpatient detoxifications	1327	Negative binomial	1,03	0,92 , 1,16	0,59
Number of outpatient treatments	1252	Negative binomial	0,99	0,94 , 1,05	0,85
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,03	0,85 , 1,26	0,74
Major depressive disorder	828	Logistic	0,94	0,81 , 1,08	0,37
Antisocial personality disorder	560	Logistic	1,02	0,86 , 1,21	0,85
Psychotic disorder	593	Logistic	1,08	0,84 , 1,4	0,53
Anxiety disorder	654	Logistic	0,98	0,83 , 1,15	0,78
Attention deficit hyperactivity disorder	700	Logistic	1,04	0,89 , 1,22	0,59
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	-0,07	-0,45 , 0,3	0,71
Aggression Hostility personality factor	667	Linear	0,05	-0,18 , 0,29	0,65
Sociability personality factor	632	Linear	0,16	-0,11 , 0,43	0,24
Impulsive sensation seeking personality factor	666	Linear	0,05	-0,28 , 0,38	0,76
Activity personality factor	665	Linear	-0,09	-0,36 , 0,18	0,53
Suicide attempt	618	Logistic	1,04	0,9 , 1,2	0,62
Suicide ideation	731	Logistic	1,05	0,9 , 1,23	0,54
Psychotic symptoms	1281	Logistic	1,14	1,02 , 1,28	2,58E-02
Sleeping disturbances	1228	Logistic	1,00	0,9 , 1,12	0,95

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,09	0,95 , 1,26	0,22
Employment status	984	Ordinal	0,98	0,88 , 1,09	0,73
Medical status	982	Ordinal	1,08	0,95 , 1,22	0,23
Psychiatric status	984	Ordinal	1,00	0,9 , 1,11	0,94
Drug use	984	Ordinal	0,93	0,83 , 1,04	0,19
Alcohol use	984	Ordinal	1,04	0,93 , 1,16	0,48
Family/Social relationships	981	Ordinal	1,01	0,9 , 1,12	0,92
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,62	-1,36 , 0,12	0,10
Mental health	751	Linear	0,32	-0,67 , 1,31	0,53
Criminal record	713	Logistic	1,08	0,93 , 1,25	0,32
Unemployment	1057	Logistic	1,14	1,01 , 1,29	3,72E-02
Number of psychiatric hospitalizations	760	Negative binomial	1,26	1,03 , 1,53	2,23E-02
Psychiatric family history	840	Logistic	1,12	0,98 , 1,28	0,11
Lifetime medical conditions	1334	Logistic	0,99	0,89 , 1,11	0,87
Substance use family history	818	Logistic	1,15	1 , 1,32	4,89E-02
Educational attainment	1333	Ordinal	0,97	1,06 , 0,88	0,51

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2d. Association between the GPS for depression and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	0,01	-0,03 , 2,00E-03	0,09
Age at onset of SUD ^c	1280	Linear	0,00	-0,02 , 0,01	0,54
Years between substance use and SUD	1280	Linear	1,01	1,01 , 1,21	3,49E-02
Years of substance use as proportion of lifespan	1199	Linear	0,20	-1,24 , 0,86	0,72
Number of substances consumed	879	Negative binomial	1,02	0,95 , 1,04	0,80
Number of therapeutic community interventions	1345	Negative binomial	0,92	0,9 , 1,08	0,76
Number of inpatient detoxifications	1327	Negative binomial	1,03	0,91 , 1,15	0,69
Number of outpatient treatments	1252	Negative binomial	0,99	1,03 , 1,15	3,04E-03
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,03	0,93 , 1,38	0,22
Major depressive disorder	828	Logistic	0,94	0,88 , 1,17	0,87
Antisocial personality disorder	560	Logistic	1,02	0,83 , 1,17	0,82
Psychotic disorder	593	Logistic	1,08	0,85 , 1,41	0,50
Anxiety disorder	654	Logistic	0,98	0,88 , 1,2	0,73
Attention deficit hyperactivity disorder	700	Logistic	1,04	0,92 , 1,25	0,41
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	-0,07	4,00E-03 , 0,72	4,91E-02
Aggression Hostility personality factor	667	Linear	0,05	0,04 , 0,5	2,18E-02
Sociability personality factor	632	Linear	0,16	-0,41 , 0,11	0,25
Impulsive sensation seeking personality factor	666	Linear	0,05	-0,22 , 0,41	0,56
Activity personality factor	665	Linear	-0,09	-0,29 , 0,23	0,83
Suicide attempt	618	Logistic	1,04	1,02 , 1,37	3,04E-02
Suicide ideation	731	Logistic	1,05	0,95 , 1,3	0,20
Psychotic symptoms	1281	Logistic	1,14	0,98 , 1,23	0,12
Sleeping disturbances	1228	Logistic	1,00	0,93 , 1,17	0,48

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,09	0,87 , 1,16	0,92
Employment status	984	Ordinal	0,98	0,96 , 1,19	0,25
Medical status	982	Ordinal	1,08	0,99 , 1,27	0,06
Psychiatric status	984	Ordinal	1,00	1 , 1,23	0,06
Drug use	984	Ordinal	0,93	0,87 , 1,09	0,68
Alcohol use	984	Ordinal	1,04	0,88 , 1,11	0,86
Family/Social relationships	981	Ordinal	1,01	0,99 , 1,23	0,08
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,62	-1,35 , 0,09	0,09
Mental health	751	Linear	0,32	-1,09 , 0,84	0,80
Criminal record	713	Logistic	1,08	1,09 , 1,48	1,82E-03
Unemployment	1057	Logistic	1,14	0,92 , 1,17	0,58
Number of psychiatric hospitalizations	760	Negative binomial	1,26	0,95 , 1,4	0,15
Psychiatric family history	840	Logistic	1,12	1,01 , 1,32	3,70E-02
Lifetime medical conditions	1334	Logistic	0,99	0,91 , 1,15	0,70
Substance use family history	818	Logistic	1,15	0,99 , 1,31	0,06
Educational attainment	1333	Ordinal	0,97	1,09 , 0,88	0,69

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2e. Association between the GPS for post-traumatic stress disorder and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,03 , -1,00E-03	3,72E-02
Age at onset of SUD ^c	1280	Linear	-0,01	-0,03 , 0,01	0,23
Years between substance use and SUD	1280	Linear	0,98	0,9 , 1,08	0,70
Years of substance use as proportion of lifespan	1199	Linear	0,16	-0,88 , 1,21	0,76
Number of substances consumed	879	Negative binomial	0,99	0,96 , 1,04	0,79
Number of therapeutic community interventions	1345	Negative binomial	1,01	0,92 , 1,11	0,89
Number of inpatient detoxifications	1327	Negative binomial	1,13	1,01 , 1,27	4,15E-02
Number of outpatient treatments	1252	Negative binomial	1,05	1 , 1,11	0,08
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,04	0,85 , 1,27	0,69
Major depressive disorder	828	Logistic	0,92	0,79 , 1,06	0,23
Antisocial personality disorder	560	Logistic	1,11	0,93 , 1,31	0,24
Psychotic disorder	593	Logistic	1,00	0,78 , 1,29	0,97
Anxiety disorder	654	Logistic	0,96	0,82 , 1,13	0,63
Attention deficit hyperactivity disorder	700	Logistic	1,15	0,99 , 1,35	0,07
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	0,15	-0,22 , 0,52	0,43
Aggression Hostility personality factor	667	Linear	0,19	-0,05 , 0,43	0,12
Sociability personality factor	632	Linear	-0,16	-0,43 , 0,11	0,24
Impulsive sensation seeking personality factor	666	Linear	-0,09	-0,41 , 0,23	0,59
Activity personality factor	665	Linear	0,05	-0,22 , 0,32	0,73
Suicide attempt	731	Logistic	1,04	0,88 , 1,22	0,68
Suicide ideation	618	Logistic	1,00	0,86 , 1,16	0,99
Psychotic symptoms	1281	Logistic	1,09	0,97 , 1,22	0,15
Sleeping disturbances	1228	Logistic	1,08	0,97 , 1,22	0,17

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,00	0,86 , 1,15	0,96
Employment status	984	Ordinal	1,00	0,9 , 1,12	0,99
Medical status	982	Ordinal	1,13	0,99 , 1,28	0,07
Psychiatric status	984	Ordinal	1,00	0,9 , 1,12	0,99
Drug use	984	Ordinal	0,96	0,86 , 1,08	0,50
Alcohol use	984	Ordinal	1,00	0,9 , 1,13	0,94
Family/Social relationships	981	Ordinal	1,06	0,95 , 1,19	0,28
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	0,32	-0,65 , 1,29	0,52
Mental health	751	Linear	-0,21	-0,94 , 0,52	0,57
Criminal record	713	Logistic	1,12	0,96 , 1,3	0,15
Unemployment	1057	Logistic	1,23	1,09 , 1,4	1,00E-03
Number of psychiatric hospitalizations	760	Negative binomial	0,86	0,7 , 1,06	0,15
Psychiatric family history	840	Logistic	1,15	1 , 1,32	0,05
Lifetime medical conditions	1334	Logistic	1,09	0,97 , 1,22	0,13
Substance use family history	818	Logistic	1,12	0,97 , 1,29	0,11
Educational attainment	1333	Ordinal	0,87	0,97 , 0,79	9,48E-03

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2f. Association between the GPS for schizophrenia and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	0,01	3,00E-04 , 0,03	4,61E-02
Age at onset of SUD ^c	1280	Linear	0,01	-0,01 , 0,02	0,44
Years between substance use and SUD	1280	Linear	0,99	0,9 , 1,08	0,80
Years of substance use as proportion of lifespan	1199	Linear	-0,11	-1,15 , 0,94	0,84
Number of substances consumed	879	Negative binomial	1,01	0,97 , 1,06	0,58
Number of therapeutic community interventions	1345	Negative binomial	1,04	0,95 , 1,14	0,43
Number of inpatient detoxifications	1327	Negative binomial	1,06	0,95 , 1,19	0,28
Number of outpatient treatments	1252	Negative binomial	0,98	0,93 , 1,04	0,49
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,00	0,82 , 1,22	0,99
Major depressive disorder	828	Logistic	0,93	0,8 , 1,07	0,31
Antisocial personality disorder	560	Logistic	1,04	0,87 , 1,23	0,67
Psychotic disorder	593	Logistic	1,37	1,05 , 1,78	1,97E-02
Anxiety disorder	654	Logistic	0,88	0,75 , 1,03	0,12
Attention deficit hyperactivity disorder	700	Logistic	0,98	0,84 , 1,15	0,84
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	0,07	-0,3 , 0,44	0,71
Aggression Hostility personality factor	667	Linear	0,18	-0,06 , 0,42	0,14
Sociability personality factor	632	Linear	0,01	-0,26 , 0,28	0,94
Impulsive sensation seeking personality factor	666	Linear	-0,17	-0,49 , 0,16	0,31
Activity personality factor	665	Linear	-0,01	-0,27 , 0,26	0,97
Suicide attempt	618	Logistic	0,90	0,78 , 1,04	0,17
Suicide ideation	731	Logistic	0,89	0,76 , 1,05	0,17
Psychotic symptoms	1281	Logistic	1,15	1,03 , 1,29	1,40E-02
Sleeping disturbances	1228	Logistic	0,93	0,83 , 1,04	0,21

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,06	0,92 , 1,23	0,42
Employment status	984	Ordinal	1,10	0,99 , 1,23	0,08
Medical status	982	Ordinal	1,02	0,9 , 1,16	0,71
Psychiatric status	984	Ordinal	1,03	0,92 , 1,15	0,60
Drug use	984	Ordinal	0,96	0,86 , 1,08	0,48
Alcohol use	984	Ordinal	1,03	0,92 , 1,16	0,61
Family/Social relationships	981	Ordinal	1,01	0,91 , 1,13	0,82
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,53	-1,26 , 0,21	0,16
Mental health	751	Linear	-0,10	-1,08 , 0,89	0,85
Criminal record	713	Logistic	1,07	0,92 , 1,24	0,41
Unemployment	1057	Logistic	1,13	1 , 1,28	4,71E-02
Number of psychiatric hospitalizations	760	Negative binomial	1,10	0,9 , 1,34	0,35
Psychiatric family history	840	Logistic	1,01	0,88 , 1,15	0,93
Lifetime medical conditions	1334	Logistic	0,99	0,88 , 1,11	0,86
Substance use family history	818	Logistic	1,03	0,9 , 1,19	0,64
Educational attainment	1333	Ordinal	1,01	1,11 , 0,91	0,87

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2g. Association between the GPS for risk tolerance and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,02 , 3,00E-03	0,12
Age at onset of SUD ^c	1280	Linear	-0,01	-0,03 , 4,00E-03	0,14
Years between substance use and SUD	1280	Linear	0,99	0,9 , 1,08	0,77
Years of substance use as proportion of lifespan	1199	Linear	0,42	-0,63 , 1,47	0,43
Number of substances consumed	879	Negative binomial	1,02	0,98 , 1,06	0,31
Number of therapeutic community interventions	1345	Negative binomial	1,02	0,93 , 1,12	0,69
Number of inpatient detoxifications	1327	Negative binomial	1,01	0,9 , 1,13	0,88
Number of outpatient treatments	1252	Negative binomial	1,06	1 , 1,12	4,33E-02
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	0,97	0,8 , 1,18	0,76
Major depressive disorder	828	Logistic	0,89	0,77 , 1,02	0,09
Antisocial personality disorder	560	Logistic	0,99	0,83 , 1,17	0,87
Psychotic disorder	593	Logistic	1,08	0,84 , 1,38	0,55
Anxiety disorder	654	Logistic	1,03	0,89 , 1,21	0,69
Attention deficit hyperactivity disorder	700	Logistic	1,08	0,93 , 1,26	0,30
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	-0,51	-0,88 , -0,14	6,90E-03
Aggression Hostility personality factor	667	Linear	0,17	-0,07 , 0,41	0,16
Sociability personality factor	632	Linear	0,09	-0,18 , 0,36	0,51
Impulsive sensation seeking personality factor	666	Linear	0,32	0 , 0,65	4,98E-02
Activity personality factor	665	Linear	0,10	-0,17 , 0,36	0,48
Suicide attempt	618	Logistic	1,11	0,95 , 1,29	0,18
Suicide ideation	731	Logistic	1,04	0,88 , 1,23	0,62
Psychotic symptoms	1281	Logistic	1,00	0,89 , 1,12	0,98
Sleeping disturbances	1228	Logistic	1,09	0,97 , 1,22	0,16

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,18	1,02 , 1,36	2,29E-02
Employment status	984	Ordinal	0,95	0,85 , 1,06	0,35
Medical status	982	Ordinal	1,03	0,91 , 1,16	0,64
Psychiatric status	984	Ordinal	0,97	0,87 , 1,08	0,59
Drug use	984	Ordinal	1,01	0,91 , 1,13	0,84
Alcohol use	984	Ordinal	1,03	0,92 , 1,15	0,58
Family/Social relationships	981	Ordinal	1,01	0,9 , 1,12	0,92
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	0,08	-0,65 , 0,81	0,83
Mental health	751	Linear	0,30	-0,67 , 1,27	0,55
Criminal record	713	Logistic	1,16	1 , 1,36	0,06
Unemployment	1057	Logistic	0,99	0,87 , 1,11	0,83
Number of psychiatric hospitalizations	760	Negative binomial	0,87	0,71 , 1,07	0,18
Psychiatric family history	840	Logistic	1,01	0,88 , 1,17	0,84
Lifetime medical conditions	1334	Logistic	1,05	0,94 , 1,18	0,37
Substance use family history	818	Logistic	1,11	0,96 , 1,28	0,15
Educational attainment	1333	Ordinal	1,04	1,15 , 0,94	0,42

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2h. Association between the GPS for suicide attempt and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,03 , 2,00E-03	0,11
Age at onset of SUD ^c	1280	Linear	-0,02	-0,04 , -0,01	1,12E-02
Years between substance use and SUD	1280	Linear	0,92	0,84 , 1,01	0,08
Years of substance use as proportion of lifespan	1199	Linear	0,03	-1,04 , 1,09	0,96
Number of substances consumed	879	Negative binomial	1,00	0,96 , 1,05	0,85
Number of therapeutic community interventions	1345	Negative binomial	0,96	0,87 , 1,06	0,40
Number of inpatient detoxifications	1327	Negative binomial	1,02	0,9 , 1,14	0,79
Number of outpatient treatments	1252	Negative binomial	1,06	1 , 1,12	4,10E-02
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,10	0,9 , 1,35	0,34
Major depressive disorder	828	Logistic	1,08	0,93 , 1,25	0,33
Antisocial personality disorder	560	Logistic	1,18	0,99 , 1,4	0,07
Psychotic disorder	593	Logistic	1,09	0,85 , 1,42	0,50
Anxiety disorder	654	Logistic	1,03	0,88 , 1,21	0,69
Attention deficit hyperactivity disorder	700	Logistic	1,09	0,93 , 1,27	0,30
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	0,09	-0,29 , 0,48	0,63
Aggression Hostility personality factor	667	Linear	0,26	0,02 , 0,51	3,60E-02
Sociability personality factor	632	Linear	0,04	-0,24 , 0,31	0,80
Impulsive sensation seeking personality factor	666	Linear	0,00	-0,33 , 0,34	0,99
Activity personality factor	665	Linear	0,24	-0,03 , 0,52	0,09
Suicide attempt	618	Logistic	1,01	0,87 , 1,18	0,85
Suicide ideation	731	Logistic	1,16	0,98 , 1,37	0,08
Psychotic symptoms	1281	Logistic	1,15	1,03 , 1,29	1,69E-02
Sleeping disturbances	1228	Logistic	0,97	0,87 , 1,09	0,65

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	1,18	1,01 , 1,37	3,27E-02
Employment status	984	Ordinal	1,03	0,92 , 1,16	0,56
Medical status	982	Ordinal	1,14	1 , 1,29	4,93E-02
Psychiatric status	984	Ordinal	1,13	1,01 , 1,27	3,88E-02
Drug use	984	Ordinal	1,05	0,93 , 1,18	0,44
Alcohol use	984	Ordinal	1,02	0,91 , 1,15	0,75
Family/Social relationships	981	Ordinal	1,13	1 , 1,26	0,05
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	-0,47	-1,21 , 0,27	0,22
Mental health	751	Linear	-0,19	-1,18 , 0,8	0,71
Criminal record	713	Logistic	1,14	0,98 , 1,32	0,11
Unemployment	1057	Logistic	1,05	0,93 , 1,19	0,40
Number of psychiatric hospitalizations	760	Negative binomial	1,08	0,88 , 1,32	0,47
Psychiatric family history	840	Logistic	0,94	0,82 , 1,08	0,40
Lifetime medical conditions	1334	Logistic	1,13	1,01 , 1,27	3,84E-02
Substance use family history	818	Logistic	1,11	0,96 , 1,28	0,15
Educational attainment	1333	Ordinal	0,90	1 , 0,81	4,30E-02

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2i. Association between the GPS for educational attainment and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	0,02	0,01 , 0,03	8,11E-03
Age at onset of SUD ^c	1280	Linear	0,02	4,00E-03 , 0,04	1,48E-02
Years between substance use and SUD	1280	Linear	1,07	0,98 , 1,17	0,15
Years of substance use as proportion of lifespan	1199	Linear	-0,73	-1,77 , 0,31	0,17
Number of substances consumed	879	Negative binomial	1,00	0,96 , 1,04	0,89
Number of therapeutic community interventions	1345	Negative binomial	0,89	0,81 , 0,98	2,28E-02
Number of inpatient detoxifications	1327	Negative binomial	0,89	0,8 , 1	0,06
Number of outpatient treatments	1252	Negative binomial	0,91	0,87 , 0,97	1,64E-03
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	0,96	0,79 , 1,18	0,70
Major depressive disorder	828	Logistic	0,91	0,79 , 1,06	0,24
Antisocial personality disorder	560	Logistic	0,90	0,75 , 1,08	0,24
Psychotic disorder	593	Logistic	1,04	0,8 , 1,37	0,76
Anxiety disorder	654	Logistic	0,97	0,83 , 1,14	0,71
Attention deficit hyperactivity disorder	700	Logistic	0,89	0,76 , 1,05	0,16
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	-0,39	-0,78 , -0,01	4,50E-02
Aggression Hostility personality factor	667	Linear	-0,18	-0,42 , 0,07	0,16
Sociability personality factor	632	Linear	0,10	-0,18 , 0,38	0,47
Impulsive sensation seeking personality factor	666	Linear	-0,05	-0,39 , 0,28	0,75
Activity personality factor	665	Linear	0,17	-0,11 , 0,45	0,24
Suicide attempt	618	Logistic	1,03	0,89 , 1,2	0,66
Suicide ideation	731	Logistic	1,00	0,85 , 1,17	0,96
Psychotic symptoms	1281	Logistic	0,91	0,82 , 1,02	0,12
Sleeping disturbances	1228	Logistic	0,91	0,81 , 1,02	0,09

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	0,96	0,82 , 1,11	0,55
Employment status	984	Ordinal	0,90	0,81 , 1,01	0,07
Medical status	982	Ordinal	1,00	0,88 , 1,13	0,99
Psychiatric status	984	Ordinal	0,91	0,81 , 1,01	0,08
Drug use	984	Ordinal	0,95	0,84 , 1,06	0,36
Alcohol use	984	Ordinal	0,98	0,87 , 1,1	0,75
Family/Social relationships	981	Ordinal	0,86	0,77 , 0,96	1,00E-02
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	0,53	-0,24 , 1,3	0,18
Mental health	751	Linear	0,59	-0,43 , 1,62	0,26
Criminal record	713	Logistic	0,67	0,57 , 0,78	8,03E-07
Unemployment	1057	Logistic	0,82	0,72 , 0,92	1,34E-03
Number of psychiatric hospitalizations	760	Negative binomial	1,15	0,94 , 1,41	0,17
Psychiatric family history	840	Logistic	1,08	0,94 , 1,23	0,28
Lifetime medical conditions	1334	Logistic	0,99	0,88 , 1,11	0,88
Substance use family history	818	Logistic	0,80	0,7 , 0,92	2,20E-03
Educational attainment	1333	Ordinal	1,39	1,54 , 1,25	3,34E-10

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 2j. Association between the GPS for well-being and 39 clinical variables from the SUD phenome. In bold nominal significant results.

Phenotypes	<i>n</i> ^a	Regression	Estimate ^b	95% CI	<i>p</i>
SUD-related phenotypes					
Age at onset of substance use ^c	1308	Linear	-0,01	-0,02 , 3,00E-03	0,13
Age at onset of SUD ^c	1280	Linear	-0,01	-0,03 , 0,01	0,25
Years between substance use and SUD	1280	Linear	0,98	0,9 , 1,07	0,69
Years of substance use as proportion of lifespan	1199	Linear	0,65	-0,4 , 1,7	0,23
Number of substances consumed	879	Negative binomial	0,99	0,95 , 1,04	0,81
Number of therapeutic community interventions	1345	Negative binomial	0,99	0,9 , 1,08	0,77
Number of inpatient detoxifications	1327	Negative binomial	0,99	0,88 , 1,11	0,87
Number of outpatient treatments	1252	Negative binomial	0,91	0,86 , 0,96	7,00E-04
Comorbidity and personality traits					
<i>Mental disorders in DSM-IV</i>					
Borderline personality disorder	447	Logistic	1,07	0,87 , 1,31	0,52
Major depressive disorder	828	Logistic	0,99	0,86 , 1,15	0,90
Antisocial personality disorder	560	Logistic	1,06	0,9 , 1,26	0,49
Psychotic disorder	593	Logistic	0,85	0,65 , 1,11	0,23
Anxiety disorder	654	Logistic	0,90	0,76 , 1,06	0,20
Attention deficit hyperactivity disorder	700	Logistic	0,92	0,79 , 1,07	0,28
<i>Zuckerman–Kuhlman Personality Questionnaire (ZKPQ)</i>					
Neuroticism Anxiety personality factor	663	Linear	-0,34	-0,72 , 0,04	0,08
Aggression Hostility personality factor	667	Linear	0,05	-0,2 , 0,29	0,71
Sociability personality factor	632	Linear	0,27	-0,01 , 0,54	0,06
Impulsive sensation seeking personality factor	666	Linear	0,25	-0,08 , 0,58	0,14
Activity personality factor	665	Linear	0,11	-0,16 , 0,38	0,43
Suicide attempt	618	Logistic	0,95	0,82 , 1,1	0,50
Suicide ideation	731	Logistic	0,93	0,79 , 1,09	0,36
Psychotic symptoms	1281	Logistic	0,93	0,83 , 1,04	0,20
Sleeping disturbances	1228	Logistic	0,91	0,82 , 1,02	0,12

Sociodemographic and health phenotypes

EuropASI

Legal status	984	Ordinal	0,95	0,82 , 1,1	0,51
Employment status	984	Ordinal	0,95	0,85 , 1,06	0,37
Medical status	982	Ordinal	0,95	0,83 , 1,07	0,40
Psychiatric status	984	Ordinal	0,87	0,78 , 0,97	1,43E-02
Drug use	984	Ordinal	0,98	0,87 , 1,09	0,67
Alcohol use	984	Ordinal	1,01	0,9 , 1,14	0,83
Family/Social relationships	981	Ordinal	0,83	0,74 , 0,93	1,00E-03
<i>36-Item Short Form Survey (SF-36)</i>					
Physical health	751	Linear	0,64	-0,1 , 1,37	0,09
Mental health	751	Linear	0,51	-0,47 , 1,49	0,31
Criminal record	713	Logistic	0,90	0,78 , 1,05	0,17
Unemployment	1057	Logistic	0,88	0,77 , 0,99	3,59E-02
Number of psychiatric hospitalizations	760	Negative binomial	0,78	0,64 , 0,95	1,33E-02
Psychiatric family history	840	Logistic	0,90	0,79 , 1,03	0,13
Lifetime medical conditions	1334	Logistic	0,98	0,87 , 1,1	0,70
Substance use family history	818	Logistic	0,83	0,73 , 0,96	1,13E-02
Educational attainment	1333	Ordinal	1,06	1,18 , 0,96	0,23

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$

^b OR is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; IRR is reported for negative binomial regression

^c Logarithmic transformations were applied to continuous variables not following a normal distribution

Supplementary Table 3.

Interaction between GPSs and lifetime emotional, physical and/or sexual abuse in the SUD phenome

GPS	SUD-Phenome	<i>n</i>^a	Estimate	95% CI	<i>p</i>
Attention-deficit hyperactivity disorder	Age at onset of substance use ^b	701	-.002	-.04 , .04	.94
	Years of substance use as proportion of lifespan	631	.32	-2.57 , 3.2	.83
	Antisocial personality disorder	358	1.00	.66 , 1.51	.99
	Attention deficit hyperactivity disorder	434	.94	.63 , 1.4	.77
	Educational attainment	701	.96	.73 , 1.26	.77
Anxiety	Number of outpatient treatments	681	1.06	.92 , 1.22	.42
	Psychotic disorder	180	1.10	.61 , 1.99	.76
	SF36 Physical health	509	.43	-1.27 , 2.12	.62
	Psychiatric family history	524	1.03	.73 , 1.46	.86
Bipolar disorder	Psychotic symptoms	678	1.01	.74 , 1.39	.93
	Unemployment	568	1.08	.78 , 1.51	.64
	Number of psychiatric hospitalizations	504	.87	.53 , 1.42	.57
	Substance use family history	516	.98	.69 , 1.39	.91
Depression	Years between substance use and SUD	691	.96	.74 , 1.25	.78
	Number of outpatient treatments	681	1.07	.92 , 1.23	.39
	ZKPQ-Neuroticism Anxiety personality factor	492	.20	-.64 , 1.05	.64
	ZKPQ-Aggression Hostility personality factor	495	-.28	-.83 , .26	.31

	Suicide attempt	725	.93	.69 , 1.26	.65
	Criminal record	488	.76	.53 , 1.1	.14
	Psychiatric family history	524	1.31	.92 , 1.86	.13
Post-traumatic stress disorder	Age at onset of substance use ^b	701	-.02	-.06 , .03	.45
	Number of inpatient detoxifications	702	.91	.65 , 1.26	.56
	Unemployment	568	.85	.6 , 1.21	.38
	Educational attainment	701	1.08	.81 , 1.44	.58
Schizophrenia	Age at onset of substance use ^b	701	.03	-.01 , .08	.11
	Psychotic disorder	180	.85	.45 , 1.59	.60
	Psychotic symptoms	678	.97	.7 , 1.33	.83
	Unemployment	568	.76	.54 , 1.07	.12
Risk tolerance	Number of outpatient treatments	681	1.12	.96 , 1.31	.14
	ZKPQ-Neuroticism Anxiety personality factor	492	.29	-.6 , 1.18	.53
	ZKPQ-Impulsive sensation seeking personality factor	493	.39	-.37 , 1.14	.32
	EuropASI-Legal status	701	1.07	.75 , 1.52	.72
	Unemployment	568	.85	.6 , 1.21	.38
Suicide attempt	Age at onset of SUD ^b	699	.00	-.04 , .05	.89
	Number of outpatient treatments	681	.98	.84 , 1.13	.77
	ZKPQ-Aggression Hostility personality factor	495	-.02	-.58 , .55	.96
	Psychotic symptoms	678	1.03	.75 , 1.41	.86
	EuropASI-Legal status	701	1.08	.76 , 1.55	.66

	EuropASI-Medical status	699	1.13	.84 , 1.51	.41
	EuropASI-Psychiatric status	701	1.35	1.03 , 1.78	2.94E-02
	Lifetime medical conditions	705	.95	.7 , 1.31	.76
	Educational attainment	701	1.10	.83 , 1.46	.51
Educational attainment	Age at onset of substance use ^b	701	.01	-.03 , .06	.50
	Age at onset of SUD ^b	699	.01	-.03 , .06	.54
	Number of therapeutic community interventions	698	.86	.67 , 1.11	.26
	Number of outpatient treatments	681	1.06	.91 , 1.23	.47
	ZKPQ-Neuroticism Anxiety personality factor	492	.16	-.73 , 1.06	.72
	Criminal record	488	1.27	.86 , 1.89	.23
	Unemployment	568	.73	.51 , 1.04	.08
	Substance use family history	516	1.00	.7 , 1.45	.99
	Educational attainment	701	.94	.71 , 1.26	.69
Well being	Number of outpatient treatments	681	1.00	.86 , 1.17	.97
	EuropASI- Psychiatric status	701	1.01	.77 , 1.32	.96
	EuropASI-Family/social relationships	701	.95	.73 , 1.25	.22
	Unemployment	568	.87	.61 , 1.25	.46
	Number of psychiatric hospitalizations	504	1.23	.73 , 2.07	.44
	Substance use family history	516	1.05	.73 , 1.52	.80

Note. Odds Ratio (OR) is reported for logistic regression and ordinal regression; Beta is reported for lineal regression; Incidence Rate Ratio (IRR) is reported for negative binomial regression.

^a For binary traits sample size was calculated with the formula $4/(1/n1+1/n2)$.

^b Logarithmic transformations were applied to continuous variables not following a normal distribution.

References

1. Demontis D, Walters RK, Martin J, Mattheisen M, Als TD, Agerbo E, et al. Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet.* 2019;51:63–75.
2. Mullins N, Forstner AJ, O'Connell KS, Coombes B, Coleman JRI, Qiao Z, et al. Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat Genet.* 2021;53:817–829.
3. Howard DM, Adams MJ, Clarke TK, Hafferty JD, Gibson J, Shiri M, et al. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci.* 2019;22:343–352.
4. Nievergelt CM, Maihofer AX, Klengel T, Atkinson EG, Chen CY, Choi KW, et al. International meta-analysis of PTSD genome-wide association studies identifies sex- and ancestry-specific genetic risk loci. *Nat Commun.* 2019;10.
5. Trubetskoy V, Pardiñas AF, Qi T, Panagiotaropoulou G, Awasthi S, Bigdeli TB, et al. Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature.* 2022;604:502.
6. Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA, et al. Genome-wide association analyses of risk tolerance and risky behaviors in over 1 million individuals identify hundreds of loci and shared genetic influences. *Nat Genet.* 2019;51:245–257.
7. Docherty AR, Mullins N, Ashley-Koch AE, Qin XJ, Coleman J, Shabalín AA, et al. Genome-wide association study meta-analysis of suicide attempt in 43,871 cases identifies twelve genome-wide significant loci. *MedRxiv.* 2022. 2022. <https://doi.org/10.1101/2022.07.03.22277199>.
8. Lee JJ, Wedow R, Okbay A, Kong E, Maghzián O, Zacher M, et al. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet.* 2018;50:1112–1121.
9. Baselmans BML, Jansen R, Ip HF, van Dongen J, Abdellaoui A, van de Weijer MP, et al. Multivariate genome-wide analyses of the well-being spectrum. *Nat Genet.* 2019;51:445–451.