

Supplemental Table 1: Patients and Demographics


Patient Characteristics	HIV-NSCLC N=18		Uninfected N=19		P-value
	N	%	N	%	
Median age	54.1		57.2		0.22
Gender					0.06
Male	14	78%	9	47%	
Female	4	22%	10	53%	
Smoking Status					0.49
Non-smoker	0	0%	2	11%	
Smoking History	18	100%	17	89%	
Year of Diagnosis					0.7
1999-2004	4	22%	6	32%	
2005-2010	8	44%	6	32%	
2011-2016	6	33%	7	37%	
Histology					0.77
Adenocarcinoma	8	44%	10	53%	
Squamous Cell	8	44%	8	42%	
NSCLC (unspecified)	2	11%	1	5%	
Stage at diagnosis					0.96
Stage I	6	33%	6	32%	
Stage II	1	6%	2	11%	
Stage III	4	22%	4	21%	
Stage IV	7	39%	7	37%	
Years of HIV Diagnosis					
Median	17	-			
Range	2-36	-			
AIDS Diagnosis	8	44%			
HIV VL (copies/mL)					
<400	8	44%			
>400	10	56%			
CD4 count (cells/ μ L)					
Median	440	-			
Range	26-872	-			

Supplemental Table 2: Number of T cells within TME


	P value	Mean (HIV)	Mean (non-HIV)	Difference	SE of difference	t ratio	df	q value
CK+	0.170988	1236	1564	328.8	237.3	1.386	60	0.703621
CD8+	0.083642	170.2	72	-98.16	55.54	1.767	47	0.697019
CD4+	0.253303	57.74	94.84	37.11	32.1	1.156	49	0.703621
CD68+	0.371457	115.8	82.43	-33.32	36.97	0.9014	53	0.773868

Supplementary Figure S1: Tumor Killing Assay with Circulating Cells from PWH and PWOH (A549)

T-cell dysfunction marker



Apoptotic cell death marker



T-cell activation marker

