

SUPPLEMENTARY INFORMATION

Non-random Mating Patterns in Education, Mental, and Somatic Health: A Population Study on Within- and Cross-Trait Associations

FA Torvik, HF Sunde, R Cheesman, NH Eftedal, MC Keller, E Ystrom, EM Eilertsen

Corresponding author: Farstein Ask Torvik, farstein.ask.torvik@fhi.no

Table of contents

Supplemental information for prospective analyses

Supplemental Figure S1	S2
Supplemental Figure S2	S3
Supplemental Figure S3	S4

Supplemental information for prospective analyses with odds ratios


Supplemental Figure S4	S5
Supplemental Figure S5	S6
Supplemental Table S1	S7

Supplemental information for cross-sectional analyses


Supplemental Table S2	S8
Supplemental Table S3	S9
Supplemental Figure S6	S10
Supplemental Figure S7	S11
Supplemental Figure S8	S12
Supplementary Figure S9	S13
Supplementary Figure S10	S14
Supplemental Figure S11	S15

Supplemental information for methods


Supplemental Figure S12	S16
Supplemental Script S1	S17
Supplemental Script S2	S18




Supplemental Figure S1. Within person correlations for educational outcomes, 10 mental health conditions, and 10 somatic health conditions, 10 to 5 years before first child. Adjusted for age and sex.




Supplemental Figure S2. Within and across-trait partner correlations for 10 mental health conditions, and 10 somatic health conditions, 10 to 5 years before first child. Adjusted for age and educational attainment.



Supplemental Figure S3. Within and across-trait partner correlations for 10 mental health conditions, and 10 somatic health conditions, 10 to 5 years before first child. Adjusted for age and grade point average.



Supplemental Figure S4. Odds ratios for within person associations between educational outcomes, 10 mental health conditions, and 10 somatic health conditions, 10 to 5 years before first child. Adjusted for age and sex.



Supplemental Figure S5. Odds ratios for within and across-trait partner associations between educational outcomes, 10 mental health conditions, and 10 somatic health conditions, 10 to 5 years before first child, all adjusted for age.

Supplemental Table S1. Results from multiple binary logistic regressions of each of the educational outcomes and the 10 mental and 10 somatic health conditions on the corresponding traits in partners, siblings, and in-laws. Health conditions were measured 10 to 5 years before a couple had their first child. Results expressed as adjusted odds ratios (OR), including 95% confidence intervals. Significant adjusted associations with siblings-in-law indicate deviations from direct assortment.

	OR(partner)	OR(sibling)	OR(in-law)
GPA among top 20%	2.76 [2.55, 3.00]	4.57 [4.22, 4.94]	1.54 [1.42, 1.68]
University education	5.35 [5.11, 5.59]	3.58 [3.44, 3.73]	1.76 [1.69, 1.83]
Depressive disorder	1.77 [1.62, 1.94]	2.36 [2.17, 2.56]	1.27 [1.15, 1.39]
Anxiety disorder	1.59 [1.29, 1.96]	2.68 [2.28, 3.16]	1.18 [0.94, 1.47]
Phobias	1.73 [1.10, 2.71]	2.17 [1.49, 3.15]	1.38 [0.85, 2.25]
PTSD	1.46 [1.31, 1.63]	2.72 [2.48, 2.98]	1.13 [1.01, 1.26]
Sleep problems	1.58 [1.38, 1.82]	1.89 [1.66, 2.15]	1.05 [0.90, 1.24]
Alcohol use disorder	1.69 [1.16, 2.46]	2.04 [1.39, 2.98]	1.41 [0.91, 2.19]
Substance use disorder	16.07 [12.83, 20.12]	4.91 [3.61, 6.68]	3.74 [2.75, 5.08]
ADHD	2.73 [2.17, 3.43]	6.40 [5.37, 7.62]	1.37 [1.03, 1.81]
Psychotic disorders	2.91 [1.53, 5.51]	3.41 [2.08, 5.59]	0.96 [0.39, 2.32]
Personality disorder	1.99 [0.49, 8.06]	3.23 [1.19, 8.74]	3.39 [1.25, 9.18]
Headaches	1.11 [1.03, 1.19]	1.49 [1.39, 1.59]	1.05 [0.98, 1.12]
Neck/back	1.25 [1.19, 1.31]	1.49 [1.42, 1.57]	1.14 [1.08, 1.20]
Abdominal pain	1.11 [1.04, 1.18]	1.41 [1.33, 1.50]	1.06 [1.00, 1.13]
Fractures	1.27 [1.14, 1.42]	1.42 [1.27, 1.58]	1.19 [1.06, 1.33]
Acne	1.14 [1.00, 1.30]	2.33 [2.10, 2.59]	0.96 [0.83, 1.11]
Injury musculoskeletal	1.29 [1.17, 1.41]	1.63 [1.49, 1.79]	1.13 [1.03, 1.25]
Asthma	1.12 [0.99, 1.27]	2.50 [2.28, 2.75]	1.04 [0.92, 1.18]
Allergic rhinitis	1.09 [1.00, 1.18]	2.03 [1.89, 2.18]	1.02 [0.94, 1.12]
Laceration/cut	1.18 [1.08, 1.28]	1.21 [1.11, 1.32]	1.04 [0.96, 1.14]
Naevus/mole	1.18 [1.09, 1.27]	1.61 [1.50, 1.73]	1.06 [0.98, 1.14]

Supplemental Table S2. List of ICPC-2 codes for the mental and somatic health condition, prevalence in the sample (including education), and prevalence among partners and relatives of affected individuals, measured cross sectionally in 2015-2019.

Variable	ICPC-2 codes	Index		Partner of affected		Sibling of affected		In-law of affected	
		n	%	n	%	n	%	n	%
GPA among top 20%		26,824	20.15	8,058	37.43	7,654	42.38	5,085	32.38
University education		93,303	49.84	64,110	68.89	48,682	61.23	46,001	58.16
Depressive disorder	P76	21,811	11.61	4,126	18.92	2,966	17.59	2,352	13.47
Anxiety disorder/anxiety state	P74	8,202	4.36	648	7.90	560	8.83	353	5.34
Phobia/compulsive disorder	P79	3,702	1.97	146	3.94	128	4.44	93	3.12
Acute stress reaction	P02	24,996	13.30	5,442	21.77	3,320	16.81	2,564	12.67
Sleep disturbance	P06	16,807	8.94	2,424	14.42	1,619	12.17	1,312	9.63
Alcohol use disorders	P15, P16	2,505	1.33	158	6.31	91	4.83	64	3.29
Substance use disorders	P18, P19	2,899	1.54	724	24.97	163	8.22	143	6.76
Hyperkinetic disorder	P81	5,216	2.78	618	11.85	542	14.03	223	5.60
Psychotic disorders	P72, P98, P73	1,965	1.05	80	4.07	95	6.26	29	1.88
Personality disorder	P80	1,652	0.88	64	3.87	53	4.39	25	1.99
Headaches	N89, N90, N01, N95	32,011	17.03	5,148	16.08	4,808	18.59	4,136	15.64
Neck/back symptom/complaint	L01, L02, L03	44,851	23.87	12,446	27.75	8,342	23.27	7,633	20.90
Abdominal pain/cramps general	D01	36,242	19.29	6,878	18.98	5,427	18.73	4,809	16.17
Fractures	L72, L73, L74, L75, L76	10,641	5.66	764	7.18	694	8.01	556	6.40
Acne	S96	7,721	4.11	326	4.22	638	10.00	356	5.53
Injury musculoskeletal	L81	12,271	6.53	1,094	8.92	911	9.16	800	7.96
Asthma	R96	11,415	6.07	800	7.01	1,069	11.51	598	6.35
Allergic rhinitis	R97	23,335	12.42	3,224	13.82	3,447	17.81	2,222	11.38
Laceration/cut	S18	15,377	8.18	1,388	9.03	1,299	10.15	1,152	9.11
Naevus/mole	S82	29,454	15.67	5,030	17.08	4,535	18.26	3,614	14.47


Supplemental Table S3. Correlations between relatives in educational outcomes and 10 mental and 10 somatic health conditions measured cross-sectionally in 2015-2019, including 95% confidence intervals, along with tests of deviations from direct assortment. Adjusted for sex and year of birth.

Variable	r(partner)	r(sibling)	r(inlaw)	Inlaw inf.	Deviation from direct assortment, p-value
Grade point average	0.42 [0.42, 0.43]	0.52 [0.52, 0.53]	0.29 [0.29, 0.30]	1.33	<1.00e-99
Educational attainment	0.48 [0.47, 0.48]	0.40 [0.40, 0.40]	0.29 [0.28, 0.29]	1.50	<1.00e-99
Depressive disorder	0.25 [0.23, 0.27]	0.17 [0.16, 0.19]	0.07 [0.05, 0.08]	1.57	3.39e-04
Anxiety disorder	0.17 [0.14, 0.20]	0.18 [0.16, 0.20]	0.05 [0.02, 0.07]	1.53	0.200
Phobias	0.14 [0.09, 0.18]	0.13 [0.10, 0.17]	0.07 [0.03, 0.11]	3.68	0.015
PTSD	0.25 [0.23, 0.26]	0.19 [0.18, 0.20]	0.06 [0.05, 0.08]	1.36	0.012
Sleep problems	0.17 [0.15, 0.19]	0.13 [0.11, 0.14]	0.05 [0.03, 0.06]	2.26	0.001
Alcohol use disorder	0.28 [0.23, 0.33]	0.20 [0.15, 0.24]	0.11 [0.07, 0.16]	2.09	0.012
Substance use disorder	0.67 [0.65, 0.70]	0.32 [0.29, 0.36]	0.27 [0.24, 0.31]	1.25	0.003
ADHD	0.33 [0.30, 0.36]	0.38 [0.36, 0.40]	0.13 [0.10, 0.16]	1.03	0.805
Psychotic disorders	0.25 [0.19, 0.31]	0.23 [0.19, 0.27]	0.02 [-0.04, 0.07]	0.31	0.171
Personality disorder	0.24 [0.17, 0.30]	0.24 [0.19, 0.29]	0.10 [0.05, 0.16]	1.83	0.145
Headaches	0.06 [0.04, 0.07]	0.11 [0.10, 0.12]	0.03 [0.01, 0.04]	3.95	0.002
Neck/back	0.12 [0.11, 0.13]	0.11 [0.10, 0.12]	0.05 [0.04, 0.06]	3.35	4.29e-10
Abdominal pain	0.09 [0.07, 0.10]	0.09 [0.08, 0.10]	0.02 [0.01, 0.03]	2.97	0.009
Fractures	0.07 [0.04, 0.10]	0.08 [0.06, 0.10]	0.02 [-0.00, 0.04]	2.70	0.350
Acne	0.04 [0.01, 0.08]	0.16 [0.14, 0.19]	0.02 [-0.01, 0.05]	2.95	0.305
Injury musculoskeletal	0.09 [0.07, 0.12]	0.08 [0.07, 0.10]	0.04 [0.02, 0.06]	4.71	0.002
Asthma	0.04 [0.01, 0.06]	0.19 [0.18, 0.21]	0.02 [-0.00, 0.04]	2.67	0.230
Allergic rhinitis	0.04 [0.02, 0.06]	0.18 [0.17, 0.19]	-0.00 [-0.01, 0.01]	-0.07	0.267
Laceration/cut	0.06 [0.04, 0.08]	0.08 [0.06, 0.09]	0.04 [0.02, 0.05]	7.79	1.27e-04
Naevus/mole	0.07 [0.06, 0.09]	0.13 [0.11, 0.14]	0.01 [0.00, 0.03]	1.58	0.384


Note: The p-value arises from comparing a constrained model, where in-law correlation is the product of partner and sibling correlations, to an unconstrained model with independent estimates for each relationship type. A low p-value signifies a poor fit for direct assortment.




Supplemental Figure S6. Prevalence of 10 mental and 10 somatic health conditions among males and females with unaffected and affected partners, measured cross sectionally in 2015-2019.




Supplemental Figure S7. Correlations between female and male partners for educational outcomes and 10 mental health and 10 somatic health phenotypes 10 to 5 years before they had their first child and cross-sectionally in 2015-2019. Including adjustment for educational attainment and grade point average in both assessments.




Supplemental Figure S8. Within person correlations for educational outcomes, 10 mental health conditions, and 10 somatic health conditions, measured cross sectionally in 2015-2019. Adjusted for age and sex.




Supplementary Figure S9. Within and across-trait partner correlations for educational outcomes, 10 mental health conditions, and 10 somatic health conditions, measured cross sectionally in 2015–2019. Adjusted for age.



Supplementary Figure S10. Within and across-trait partner correlations for 10 mental health conditions, and 10 somatic health conditions, measured cross sectionally in 2015-2019. Adjusted for age and educational attainment.



Supplementary Figure S11. Within and across-trait partner correlations for 10 mental health conditions, and 10 somatic health conditions, measured cross sectionally in 2015-2019. Adjusted for age and grade point average attainment.



Supplemental Figure S12. The correlation between observed phenotypes (squares) can be derived by path tracing rules allowing for copaths, so that e.g., $r_{\text{partner}} = a*m*a + x*m*x + a*x^2 + s^2$; $r_{\text{sibling}} = a*f*a + x*m*f*a + s*s$; $r_{\text{inlaw}} = a*m*f*a + x*f*a + s*s + a*m*f*m*x + x*f*m*x$, and the inlaw inflation factor $= r_{\text{inlaw}}/(r_{\text{partner}} * r_{\text{sibling}})$. With direct assortment, $a=1$, $x=0$, and $s=0$, this reduces to $r_{\text{partner}} = m$, $r_{\text{sibling}} = f$, $r_{\text{inlaw}} = m*f$, and the inlaw inflation factor $= r_{\text{inlaw}}/(r_{\text{partner}} * r_{\text{sibling}}) = m*f/(m*f)=1$. With indirect assortment ($a<1$; assuming $x=0$; $s=0$), one would expect all correlations between individuals' phenotypes to be deflated by the same degree; the in-laws correlation as much as the partner correlation and the sibling correlation. If a is the correlation between the assorted variable and the observed variable, then $r_{\text{partner}}=ma^2$ and $r_{\text{sibling}}=fa^2$. Then $r_{\text{inlaw}} = mra^2$ would be higher than the product $r_{\text{partner}} * r_{\text{sibling}} = ma^2fa^2$. Hence, the in-law correlation – which is deflated once – should be larger than the product of the sibling and the partner correlations – which are deflated one time each. Note that measurement error is one example of indirect assortment – people assort on some true value, although what we observe is the phenotype with measurement error. Hence, with indirect assortment, one expects the in-law inflation factor to be above 1. With social homogamy ($s>0$; assuming $x=0$), the strength of the social homogamy is added to the correlations between all relatives, so that the observed $r_{\text{partner}}=a^2m+s^2$, $r_{\text{sibling}}=a^2f+s^2$, and $r_{\text{inlaw}}=a^2mf+s^2$. This typically leads to in-law inflation factor above 1.00, although in cases with implausibly high phenotypic resemblance ($m+f+s^2 > 1$), the in-law inflation factor can be less than 1.00. With convergence ($x>0$), the partner correlation would share variation that does not influence correlations with siblings and inlaws as much. If x is the degree of convergence, $r_{\text{partner}}=a^2m+2xa+x^2m + s^2$, $r_{\text{sibling}}=a^2f+xmfa + s^2$, and $r_{\text{inlaw}}=a^2mf + xfa(1+m^2) + x^2fm + s^2$. All else being equal, $x > 0$ leads to an in-law inflation factor less than 1.00.

```

library(polykor)

# dichotomized normally distributed variables
n = 10000
vc = rnorm(n)
va = vc+rnorm(n)
vb = vc+rnorm(n)

va_dich = ifelse(va<2,0,1)
vb_dich = ifelse(vb<2,0,1)

# pearson correlation between original variables
cor(va,vb)

# pearson correlation between dichotomized variables (underestimated)
cor(va_dich, vb_dich)

# polychoric correlation between dichotomized variables (correct)
polychor(va_dich,vb_dich)

# dichotomized skewed variables
n = 100000
vc = rnorm(n)
va = (vc+rnorm(n))^2
vb = (vc+rnorm(n))^2

hist(va)

va_dich = ifelse(va<5,0,1)
vb_dich = ifelse(vb<5,0,1)

# pearson correlation between original variables
cor(va,vb)

# pearson correlation between dichotomized variables
cor(va_dich, vb_dich)

# polychoric correlation between dichotomized variables (overestimated)
polychor(va_dich,vb_dich)

```

Supplemental Script S1. Tetrachoric correlations for skewed variables. Tetrachoric correlations get overestimated when based on non-normal variables.

```

library(tidyverse)
library(polycor)

(randvar=runif(1))

# normally distributed variables

n = 100000
common1 = rnorm(n)
person_i = common1 + rnorm(n)*randvar    # index
person_ip = common1 + rnorm(n)*randvar    # index' partner
person_ips = person_ip/2 + rnorm(n)*randvar # index' partner's sibling (inlaw)

# correlation matrix
cor(tibble(person_i, person_ip, person_ips))

# observed correlations between indirectly associated variables (e.g. between in-laws)
cor(person_i, person_ips)

# observed product of correlations (matches the expectation)
cor(person_i, person_ip)*cor(person_ip, person_ips)

# skewed variables
person_i2 = person_i^2
person_ip2 = person_ip^2
person_ips2 = person_ips^2

# correlation matrix
cor(tibble(person_i2, person_ip2, person_ips2))

# observed correlations between indirectly associated variables (e.g. between in-laws)
cor(person_i2, person_ips2)

# observed product of correlations (still matches the expectation)
cor(person_i2, person_ip2)*cor(person_ip2, person_ips2)

# dichotomized skewed variables
threshold=3
person_i2_dich = ifelse(person_i2<threshold,0,1)
person_ip2_dich = ifelse(person_ip2<threshold,0,1)
person_ips2_dich = ifelse(person_ips2<threshold,0,1)

# correlation matrix
cor(tibble(person_i2_dich, person_ip2_dich, person_ips2_dich))

# observed pearson correlations between indirectly associated variables (e.g. between in-laws)
cor(person_i2_dich, person_ips2_dich)

# observed product of pearson correlations (lower than the expectation)
cor(person_i2_dich, person_ip2_dich)*cor(person_ip2_dich, person_ips2_dich)

# observed polychoric correlations between indirectly associated variables (e.g. between in-laws)
polychor(person_i2_dich, person_ips2_dich)

# observed product of polychoric correlations (matches the expectation again)
polychor(person_i2_dich, person_ip2_dich)*polychor(person_ip2_dich, person_ips2_dich)

```

Supplemental Script S2. The product of tetrachoric correlations for skewed variables. This script shows that if the product of two Pearson correlations matches with a third correlation, this is also the case for tetrachoric correlations after dichotomization, even if the underlying variables are skewed.