

Supplemental material

Table 1. Acute test symptoms of dysautonomia on standing, presenting in Long COVID patients who could not complete the lean test because of excessive symptoms.

Symptom	In Long COVID patients who could not complete lean test (n=46)
<i>Symptoms of orthostatic intolerance:</i>	
Dizziness / light-headedness	23 (50%)
Palpitations	5 (11%)
Chest pain / discomfort	2 (4%)
Tremors	1 (2%)
<i>Other symptoms of dysautonomia:</i>	
Tingling / numbness in skin	11 (24%)
Fatigue	11 (24%)
Sweating / clammy / hot	5 (11%)
Breathlessness	2 (4%)
Headache	4 (9%)
Muscle / joint pain	2 (4%)
Cognitive dysfunction	1 (2%)
Visual disturbances	1 (2%)

Supplemental Table 2. Associations between patient characteristics, pre-existing comorbidities and prior symptoms with criteria for POTS, both unadjusted and adjusted for age and sex, in Long COVID patients.

	POTS (n=20)	No POTS (n=257)	Unadjusted OR (95% CI)	Adjusted OR* (95% CI)
Patient characteristics (%):				
Age ≥50 years	5 (25%)	139 (54%)	0.3 (0.1, 0.8)	0.3 (0.1, 0.8)
Female gender	6 (30%)	98 (38%)	1.4 (0.5, 3.9)	1.3 (0.5, 3.6)
Body mass index ≥30 kg/m ²	4 (27%)	79 (41%)	0.5 (0.2, 1.7)	0.7 (0.2, 2.2)
Non-white ethnicity	2 (10%)	44 (18%)	0.5 (0.1, 2.3)	0.4 (0.1, 1.6)
Long COVID duration ≥12 months	18 (90%)	226 (88%)	1.2 (0.3, 5.6)	1.0 (0.2, 4.7)
Pre-existing comorbidities (%):				
Allergies or autoimmune conditions	4 (20%)	40 (15%)	1.4 (0.4, 4.3)	1.6 (0.5, 5.2)
Other inflammatory conditions	1 (5%)	11 (4%)	1.2 (0.1, 9.6)	3.0 (0.3, 27.8)
Hypertension	2 (10%)	26 (10%)	1.0 (0.2, 4.5)	1.4 (0.3, 7.1)
Other heart conditions	2 (10%)	20 (8%)	1.3 (0.3, 6.1)	1.5 (0.3, 7.0)
Type 2 diabetes mellitus	0 (0%)	16 (6%)	-	-
Mental health condition	6 (30%)	37 (14%)	2.5 (0.9, 7.1)	2.2 (0.8, 6.3)
History suggestive of orthostatic intolerance (%):				
Dizziness or light headedness	14 (70%)	91 (35%)	4.3 (1.6, 11.5)	3.9 (1.4, 10.8)
Palpitations	7 (35%)	63 (25%)	1.7 (0.6, 4.3)	1.4 (0.5, 3.7)
Chest pain or discomfort	3 (15%)	24 (9%)	1.7 (0.5, 6.3)	1.2 (0.3, 4.7)
Tremors	2 (10%)	9 (4%)	3.1 (0.6, 15.2)	4.2 (0.8, 22.9)
Any of dizziness, palpitations, chest pain, tremors	14 (70%)	116 (45%)	2.8 (1.1, 7.6)	2.3 (0.8, 6.3)
Other prior symptoms (%):				
Fatigue	3 (15%)	27 (11%)	1.5 (0.4, 5.5)	1.5 (0.4, 5.7)
Aches or pains	2 (10%)	22 (9%)	1.2 (0.3, 5.5)	1.0 (0.2, 5.0)
Breathlessness	2 (10%)	14 (5%)	1.9 (0.4, 9.2)	2.3 (0.5, 11.2)

* Odds ratios adjusted for age and sex. Age and sex only adjusted for each other.

Supplemental Table 3. Associations between patient characteristics, pre-existing comorbidities, and prior symptoms with criteria for orthostatic hypotension, both unadjusted and adjusted for age and sex, in Long COVID patients.

	Orthostatic hypotension (n=21)	No Orthostatic hypotension (n=256)	Unadjusted OR (95% CI)	Adjusted OR* (95% CI)
Patient characteristics (%):				
Age ≥50 years	19 (90%)	125 (49%)	10.0 (2.3, 43.6)	10.5 (2.4, 46.3)
Female gender	8 (38%)	96 (38%)	1.0 (0.4, 2.4)	1.4 (0.5, 3.6)
Body mass index ≥30 kg/m ²	5 (31%)	78 (41%)	0.7 (0.2, 2.0)	0.4 (0.1, 1.4)
Non-white ethnicity	2 (10%)	44 (18%)	0.5 (0.1, 2.3)	0.8 (0.2, 3.9)
Long COVID duration ≥12 months	19 (90%)	225 (88%)	1.3 (0.3, 5.9)	1.7 (0.4, 8.1)
Pre-existing comorbidities (%):				
Allergies or autoimmune conditions	3 (14%)	41 (16%)	0.9 (0.2, 3.1)	1.0 (0.3, 3.6)
Other inflammatory conditions	1 (5%)	11 (4%)	1.1 (0.1, 9.1)	0.7 (0.1, 6.1)
Hypertension	4 (19%)	24 (9%)	2.3 (0.7, 7.3)	1.9 (0.6, 6.4)
Other heart conditions	1 (5%)	21 (8%)	0.6 (0.1, 4.4)	0.6 (0.1, 4.8)
Type 2 diabetes mellitus	4 (19%)	12 (5%)	4.8 (1.4, 16.4)	2.6 (0.7, 9.5)
Mental health condition	2 (10%)	41 (16%)	0.6 (0.1, 2.5)	0.7 (0.1, 3.2)
History suggestive of orthostatic intolerance (%):				
Dizziness or light headedness	7 (33%)	98 (38%)	0.8 (0.3, 2.1)	1.1 (0.4, 3.1)
Palpitations	5 (24%)	65 (25%)	0.9 (0.3, 2.6)	1.3 (0.4, 4.0)
Chest pain or discomfort	0 (0%)	27 (11%)	-	-
Tremors	0 (0%)	11 (4%)	-	-
Any of dizziness, palpitations, chest pain, tremors	7 (33%)	123 (48%)	0.5 (0.2, 1.4)	0.8 (0.3, 2.2)
Other prior symptoms (%):				
Fatigue	6 (29%)	24 (9%)	3.9 (1.4, 10.9)	4.3 (1.4, 13.2)
Aches or pains	1 (5%)	23 (9%)	0.5 (0.1, 3.9)	0.6 (0.1, 5.1)
Breathlessness	1 (5%)	15 (6%)	0.8 (0.1, 6.4)	1.2 (0.1, 10.1)

* Odds ratios adjusted for age and sex. Age and sex only adjusted for each other.

Supplemental Table 4. – Numbers and percentages of participants presenting during the lean test with sustained increased heart rate, orthostatic hypotension, or narrow pulse pressure, with and without acute symptoms during the lean test.

Lean test measures	Long COVID patients (n=277)				Healthy controls (n=64)			
	History of Long COVID symptoms suggestive of orthostatic intolerance §				History of Long COVID symptoms suggestive of orthostatic intolerance §			
	No (n=147)		Yes (n=130)		No (n=50)		Yes (n=0)	
	n (%)	(95% CI)	n (%)	(95% CI)	n (%)	(95% CI)	n (%)	(95% CI)
Orthostatic tachycardia								
(Increased heart rate* with no drop in blood pressure†)								
<i>With no acute symptoms during lean test</i>	5 (3%)	(1, 8%)	3 (2%)	(0, 7%)	0 (0%)	(0, 7%)	-	-
<i>With acute symptoms during lean test (POTS)‡</i>	6 (4%)	(2, 9%)	14 (11%)	(6, 17%)	0 (0%)	(0, 7%)	-	-
Orthostatic hypotension								
(Fall in blood pressure†)								
<i>With no acute symptoms during lean test</i>	6 (4%)	(2, 9%)	4 (3%)	(0, 8%)	5 (10%)	(3, 22%)	-	-
<i>With acute symptoms during lean test‡</i>	8 (5%)	(2, 10%)	3 (2%)	(0, 7%)	0 (0%)	(0, 7%)	-	-
Narrow pulse pressure								
(Pulse pressure <25% of systolic blood pressure)								
<i>With no acute symptoms during lean test</i>	27 (18%)	(12, 26%)	24 (18%)	(12, 26%)	10 (20%)	(10, 34%)	-	-
<i>With acute symptoms during lean test‡</i>	31 (21%)	(15, 29%)	40 (31%)	(23, 39%)	0 (0%)	(0, 7%)	-	-

* Heart rate increased by ≥30bpm (age 18+) or ≥40 bpm (age <18) sustained over two consecutive time points

† Fall in systolic blood pressure ≥20mmHg or diastolic blood pressure ≥10mmHg within first four minutes

‡ Symptoms of dysautonomia included dizziness, palpitations, chest pain or tremors, fatigue, muscle or joint pain, cognitive dysfunction, headache, visual disturbances, breathlessness, tingling or numbness in the skin, sweating or clamminess.

§ History of Long COVID symptoms suggestive of orthostatic intolerance prior to the lean test was taken as dizziness, palpitations, chest pain or tremors