

Supplementary material.

Supplementary table 1. Association between the MetaboHealth score and cognitive function – stratified for treatment group.

Cognitive test	MetaboHealth Score (per SD)						
	Lower third Beta (95% CI)	Middle third Placebo Beta (95% CI)	Middle third Pravastatin Beta (95% CI)	Upper third Placebo Beta (95% CI)	Upper third Pravastatin Beta (95% CI)	Continuous Placebo Beta (95% CI)	Continuous Pravastatin Beta (95% CI)
Minimally adjusted¹							
Stroop, seconds	Ref	4.60 (1.95; 7.24)	3.27 (0.64; 5.89)	8.32 (5.49; 11.14)	10.09 (7.49; 12.69)	9.47 (6.92; 12.02)	9.75 (7.48; 12.03)
LDCT, digits coded	Ref	-1.71 (-2.40; -1.02)	-1.40 (-2.15; -0.65)	-2.36 (-3.09; -1.63)	-3.33 (-4.07; -2.59)	-2.64 (-3.30; -1.98)	-3.17 (-3.82; -2.53)
PLTi, pictures remembered	Ref	-0.12 (-0.30; 0.05)	-0.19 (-0.37; -0.01)	-0.44 (-0.63; -0.25)	-0.46 (-0.64; -0.28)	-0.45 (-0.61; -0.28)	-0.45 (-0.61; -0.30)
PLTd, pictures remembered	Ref	-0.14 (-0.40; 0.12)	-0.09 (-0.36; 0.17)	-0.53 (-0.80; -0.26)	-0.40 (-0.66; -0.14)	-0.58 (-0.82; -0.34)	-0.49 (-0.72; -0.27)
Barthel, index	Ref	0.07 (-0.01; 0.15)	0.04 (-0.05; 0.14)	0.04 (-0.05; 0.12)	-0.10 (-0.19; 0.00)	-0.09 (-0.16; -0.02)	-0.20 (-0.28; -0.12)
IADL, points	Ref	0.07 (-0.05; 0.19)	0.04 (-0.09; 0.17)	0.01 (-0.11; 0.14)	-0.18 (-0.31; -0.06)	-0.24 (-0.35; -0.14)	-0.34 (-0.45; -0.24)
Fully adjusted²							
Stroop, seconds	Ref	4.62 (1.95; 7.28)	3.48 (0.84; 6.13)	7.37 (4.50; 10.24)	9.80 (7.15; 12.46)	8.68 (6.08; 11.27)	9.38 (7.04; 11.71)
LDCT, digits coded	Ref	-1.68 (-2.38; -0.99)	-1.49 (-2.24; -0.74)	-2.18 (-2.93; -1.43)	-3.34 (-4.10; -2.58)	-2.44 (-3.21; -1.76)	-3.13 (-3.79; -2.47)
PLTi, pictures remembered	Ref	-0.14 (-0.32; 0.04)	-0.19 (-0.37; -0.01)	-0.43 (-0.63; -0.24)	-0.44 (-0.63; -0.26)	-0.43 (-0.61; -0.26)	-0.44 (-0.60; -0.28)
PLTd, pictures remembered	Ref	-0.14 (-0.40; 0.12)	-0.10 (-0.36; 0.17)	-0.49 (-0.77; -0.21)	-0.38 (-0.65; -0.11)	-0.55 (-0.80; -0.30)	-0.47 (-0.70; -0.24)
Barthel, index	Ref	0.09 (0.01; 0.17)	0.04 (-0.06; 0.13)	0.06 (-0.03; 0.15)	-0.11 (-0.20; -0.01)	-0.07 (-0.15; -0.00)	-0.20 (-0.28; -0.12)
IADL, points	Ref	0.09 (-0.03; 0.21)	0.02 (-0.11; 0.15)	0.04 (-0.09; 0.17)	-0.19 (-0.32; -0.06)	-0.22 (-0.33; -0.12)	-0.34 (-0.45; -0.23)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, systolic blood pressure, apolipoprotein E.

Supplementary table 2. Association between the MetaboHealth score and cognitive decline – stratified for treatment group.

Cognitive test	MetaboHealth Score (per SD)						
	Lower third	Middle third Placebo Estimate (95% CI)	Middle third Pravastatin Estimate (95% CI)	Upper third Placebo Estimate (95% CI)	Upper third Pravastatin Estimate (95% CI)	Continuous Placebo Estimate (95% CI)	Continuous Pravastatin Estimate (95% CI)
	Estimate (95% CI)						
Minimally adjusted¹							
Stroop, seconds	Ref	0.58 (0.11; 1.05)	0.59 (0.14; 1.04)	0.65 (0.16; 1.14)	0.57 (0.12; 1.02)	0.42 (-0.03; 0.87)	0.63 (0.23; 1.02)
LDCT, digits coded	Ref	-0.02 (-0.12; 0.08)	-0.02 (-0.12; 0.08)	-0.10 (-0.21; 0.00)	0.00 (-0.10; 0.10)	-0.07 (-0.17; 0.02)	-0.07 (-0.16; 0.02)
PLTi, pictures remembered	Ref	-0.01 (-0.05; 0.03)	-0.04 (-0.08; 0.00)	-0.07 (-0.11; -0.02)	-0.06 (-0.10; -0.02)	-0.05 (-0.90; -0.01)	-0.05 (-0.08; -0.01)
PLTd, pictures remembered	Ref	-0.04 (-0.10; 0.02)	-0.11 (-0.17; -0.05)	-0.11 (-0.17; -0.05)	-0.12 (-0.18; -0.06)	-0.08 (-0.13; -0.02)	-0.11 (-0.16; -0.06)
Barthel, index	Ref	-0.01 (-0.04; 0.02)	0.01 (-0.01; 0.04)	-0.06 (-0.09; -0.03)	-0.04 (-0.07; -0.02)	-0.06 (-0.09; -0.04)	-0.06 (-0.09; -0.04)
IADL, points	Ref	-0.01 (-0.05; 0.02)	0.04 (0.00; -0.08)	-0.10 (-0.14; -0.06)	-0.02 (-0.06; 0.01)	-0.10 (-0.13; -0.06)	-0.07 (-0.10; -0.04)
Fully adjusted²							
Stroop, seconds	Ref	0.49 (0.03; 0.96)	0.64 (0.19; 1.10)	0.60 (0.11; 1.09)	2.63 (0.40; 4.87)	0.36 (-0.08; 0.81)	0.66 (0.26; 1.06)
LDCT, digits coded	Ref	-0.02 (0.12; 0.08)	-0.02 (-0.12; 0.08)	-0.11 (-0.21; 0.00)	-0.01 (-0.11; 0.10)	-0.07 (-0.17; 0.02)	-0.07 (-0.16; 0.02)
PLTi, pictures remembered	Ref	-0.02 (-0.06; 0.02)	-0.04 (-0.08; 0.00)	-0.07 (-0.11; -0.02)	-0.06 (-0.10; -0.02)	-0.05 (-0.09; -0.01)	-0.04 (-0.08; -0.01)
PLTd, pictures remembered	Ref	-0.04 (-0.10; 0.02)	0.26 (0.03; 0.50)	-0.11 (-0.18; -0.05)	-0.12 (-0.18; -0.06)	-0.08 (-0.14; -0.02)	-0.11 (-0.16; -0.05)
Barthel, index	Ref	-0.02 (-0.05; 0.02)	0.03 (-0.08; 0.13)	-0.07 (-0.10; -0.04)	-0.06 (-0.08; -0.02)	-0.07 (-0.10; -0.04)	-0.07 (-0.09; -0.04)
IADL, points	Ref	-0.02 (-0.06; 0.02)	0.03 (-0.01; 0.07)	-0.10 (-0.14; -0.06)	-0.04 (-0.08; 0.00)	-0.10 (-0.13; -0.06)	-0.07 (-0.10; -0.04)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, systolic blood pressure, apolipoprotein E.

Supplementary table 3. Association between MetaboHealth score and cognitive function – stratified for history of cardiovascular disease.

Cognitive test	MetaboHealth Score (per SD)						
	Lower third		Middle third		Upper third		Continuous
	Beta (95% CI)	No Beta (95% CI)	Yes Beta (95% CI)	No Beta (95% CI)	Yes Beta (95% CI)	No Beta (95% CI)	Yes Beta (95% CI)
Minimally adjusted¹							
Stroop, seconds	Ref	4.45 (1.91; 7.00)	3.37 (0.66; 6.08)	9.65 (6.96; 12.34)	8.69 (5.99; 11.40)	8.84 (6.47; 11.21)	10.45 (8.04; 12.86)
LDCT, digits coded	Ref	-1.56 (-2.24; -0.88)	-1.59 (-2.35; -0.83)	-2.79 (-3.50; -2.08)	-2.94 (-3.70; -2.18)	-2.78 (-3.40; -2.15)	-3.07 (-3.75; -2.39)
PLTi, pictures remembered	Ref	-0.14 (-0.30; 0.03)	-0.18 (-0.37; 0.02)	-0.40 (-0.58; -0.23)	-0.49 (-0.69; -0.30)	-0.42 (-0.57; -0.27)	-0.48 (-0.66; -0.31)
PLTd, pictures remembered	Ref	-0.15 (-0.39; 0.09)	-0.07 (-0.35; 0.22)	-0.40 (-0.65; -0.16)	-0.49 (-0.77; -0.20)	-0.50 (-0.72; -0.29)	-0.53 (-0.79; -0.28)
Barthel, index	Ref	0.02 (-0.06; 0.10)	0.10 (0.00; 0.21)	0.03 (-0.04; 0.11)	-0.09 (-0.20; 0.01)	-0.05 (-0.11; 0.02)	-0.27 (-0.36; -0.18)
IADL, points	Ref	-0.01 (-0.10; 0.09)	0.14 (-0.01; 0.30)	0.01 (-0.10; 0.11)	-0.17 (-0.32; -0.01)	-0.13 (-0.21; -0.05)	-0.50 (-0.63; -0.36)
Fully adjusted²							
Stroop, seconds	Ref	4.53 (1.97; 7.09)	3.42 (0.67; 6.17)	8.73 (6.02; 11.44)	8.00 (5.22; 10.79)	8.21 (5.82; 10.60)	9.74 (7.24; 12.24)
LDCT, digits coded	Ref	-1.56 (-2.25; -0.87)	-1.63 (-2.39; -0.86)	-2.68 (-3.40; -1.95)	-2.79 (-3.57; -2.01)	-2.65 (-3.29; -2.01)	-2.89 (-3.59; -2.18)
PLTi, pictures remembered	Ref	-0.14 (-0.30; 0.03)	-0.19 (-0.39; 0.02)	-0.40 (-0.57; -0.22)	-0.46 (-0.67; -0.26)	-0.42 (-0.57; -0.26)	-0.45 (-0.63; -0.27)
PLTd, pictures remembered	Ref	-0.15 (-0.39; 0.09)	-0.06 (-0.35; 0.23)	-0.39 (-0.65; -0.14)	-0.42 (-0.71; -0.13)	-0.50 (-0.73; -0.28)	-0.47 (-0.73; -0.20)
Barthel, index	Ref	0.03 (-0.04; 0.11)	0.11 (0.00; 0.21)	0.04 (-0.04; 0.12)	-0.09 (-0.19; 0.02)	-0.04 (-0.10; 0.03)	-0.27 (-0.36; -0.18)
IADL, points	Ref	0.01 (-0.09; 0.10)	0.13 (-0.03; 0.28)	0.01 (-0.09; 0.11)	-0.16 (-0.32; 0.00)	-0.12 (-0.21; -0.04)	-0.48 (-0.61; -0.34)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of diabetes, history of myocardial infarct, smoking, alcohol intake, treatment arm, systolic blood pressure, apolipoprotein E.

Supplementary table 4. Association between the MetaboHealth score and cognitive decline – stratified for history of cardiovascular disease.

Cognitive test	MetaboHealth Score (per SD)						
	Lower third		Middle third		Upper third		Continuous
	No	Estimate (95% CI)	Yes	Estimate (95% CI)	No	Estimate (95% CI)	No
Minimally adjusted¹							
Stroop, seconds	Ref	0.52 (0.08; 0.96)	0.59 (-1.89; 3.07)	0.56 (0.09; 1.02)	0.69 (0.22; 1.16)	0.55 (0.14; 0.96)	0.52 (0.09; 0.95)
LDCT, digits coded	Ref	0.03 (-0.06; 0.12)	-0.67 (-1.30; -0.04)	-0.09 (-0.19; 0.01)	0.00 (-0.11; 0.10)	-0.13 (-0.21; 0.04)	0.00 (-0.10; 0.10)
PLTi, pictures remembered	Ref	-0.03 (-0.07; 0.01)	-0.02 (-0.07; 0.02)	-0.06 (-0.10; -0.02)	-0.06 (-0.11; -0.02)	-0.06 (-0.09; -0.02)	-0.04 (-0.08; 0.00)
PLTd, pictures remembered	Ref	-0.05 (-0.10; 0.01)	-0.11 (-0.18; -0.04)	-0.13 (-0.19; -0.07)	-0.10 (-0.16; -0.03)	-0.12 (-0.17; -0.06)	-0.07 (-0.13; -0.01)
Barthel, index	Ref	0.01 (-0.01; 0.04)	-0.01 (-0.05; 0.02)	-0.05 (-0.07; -0.02)	-0.06 (-0.10; -0.03)	-0.07 (-0.09; -0.04)	-0.06 (-0.09; -0.03)
IADL, points	Ref	0.01 (-0.02; 0.05)	0.01 (-0.03; 0.06)	-0.05 (-0.08; -0.01)	-0.07 (-0.11; -0.03)	-0.07 (-0.10; -0.04)	-0.08 (-0.12; -0.05)
Fully adjusted²							
Stroop, seconds	Ref	0.48 (0.04; 0.92)	0.69 (0.21; 1.16)	0.53 (0.07; 1.00)	0.77 (0.29; 1.25)	0.51 (0.10; 0.92)	0.56 (0.13; 1.00)
LDCT, digits coded	Ref	0.04 (-0.06; 0.13)	-0.10 (-0.21; 0.01)	-0.08 (-0.18; 0.02)	-1.25 (-1.89; -0.62)	-0.12 (-0.21; -0.03)	0.00 (-0.10; 0.09)
PLTi, pictures remembered	Ref	-0.03 (-0.07; 0.01)	-0.03 (-0.08; 0.01)	-0.06 (-0.19; -0.01)	-0.07 (-0.11; -0.02)	-0.06 (-0.09; -0.02)	-0.04 (-0.08; 0.00)
PLTd, pictures remembered	Ref	-0.05 (-0.10; 0.01)	-0.12 (-0.18; -0.05)	-0.13 (-0.19; -0.07)	-0.10 (-0.17; -0.03)	-0.11 (-0.17; -0.06)	-0.06 (-0.12; 0.00)
Barthel, index	Ref	0.01 (-0.02; 0.03)	-0.01 (-0.05; 0.02)	-0.06 (-0.08; -0.03)	-0.06 (-0.10; -0.03)	-0.07 (-0.09; -0.04)	-0.06 (-0.09; -0.03)
IADL, points	Ref	0.00 (-0.03; 0.03)	0.02 (-0.03; 0.06)	-0.06 (-0.10; -0.03)	-0.07 (-0.12; -0.03)	-0.07 (-0.10; -0.05)	-0.09 (-0.13; -0.05)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of diabetes, history of myocardial infarct, smoking, alcohol intake, treatment arm, systolic blood pressure, apolipoprotein E.

Supplementary table 5. Cross-sectional associations of MetaboHealth score and cognitive function – individuals with disease during follow-up excluded.

Cognitive test	MetaboHealth Score (per SD)			
	Lower third (N _{max} =1444) Beta (95% CI)	Middle third (N _{max} =1329) Beta (95% CI)	Upper third (N _{max} =1236) Beta (95% CI)	Continuous (N _{max} =4094) Beta (95% CI)
Minimally adjusted¹				
Stroop, seconds	Ref	5.05 (3.04; 7.07)	9.11 (6.99; 11.22)	8.33 (6.47; 10.20)
LDCT, digits coded	Ref	-1.93 (-2.50; -1.36)	-3.07 (-3.67; -2.47)	-2.97 (-3.50; -2.44)
PLTi, pictures remembered	Ref	-0.18 (-0.32; -0.04)	-0.47 (-0.62; -0.32)	-0.41 (-0.54; -0.28)
PLTd, pictures remembered	Ref	-0.20 (-0.41; 0.01)	-0.48 (-0.69; -0.26)	-0.50 (-0.70; -0.31)
Barthel, index	Ref	0.01 (-0.05; -0.06)	-0.07 (-0.12; -0.01)	-0.08 (-0.14; -0.03)
IADL, points	Ref	-0.01 (-0.09; 0.08)	-0.14 (-0.23; -0.06)	-0.20 (-0.28; -0.12)
Fully adjusted²				
Stroop, seconds	Ref	4.89 (2.87; 6.91)	8.42 (6.28; 10.56)	7.85 (5.96; 9.74)
LDCT, digits coded	Ref	-1.86 (-2.44; -1.28)	-2.93 (-3.54; -2.32)	-2.87 (-3.42; -2.33)
PLTi, pictures remembered	Ref	-0.20 (-0.34; -0.05)	-0.47 (-0.63; -0.32)	-0.41 (-0.54; -0.27)
PLTd, pictures remembered	Ref	-0.22 (-0.43; -0.01)	-0.47 (-0.69; -0.25)	-0.50 (-0.70; -0.30)
Barthel, index	Ref	0.02 (-0.03; 0.08)	-0.05 (-0.11; 0.01)	-0.07 (-0.12; -0.02)
IADL, points	Ref	0.01 (-0.08; 0.09)	-0.11 (-0.20; -0.03)	-0.18 (-0.26; -0.10)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, treatment arm, systolic blood pressure, apolipoprotein E.

Supplementary table 6. Association between the MetaboHealth score and cognitive decline – individuals with disease during follow-up excluded.

Cognitive test	MetaboHealth Score (per SD)			
	Lower third (N _{max} =1444) Estimate (95% CI)	Middle third (N _{max} =1329) Estimate (95% CI)	Upper third (N _{max} =1236) Estimate (95% CI)	Continuous (N _{max} =4094) Estimate (95% CI)
Minimally adjusted¹				
Stroop, seconds	Ref	0.58 (0.24; 0.91)	0.58 (0.23; 0.93)	0.59 (0.29; 0.90)
LDCT, digits coded	Ref	0.00 (-0.08; 0.08)	-0.05 (-0.13; 0.03)	-0.08 (-0.15; -0.01)
PLTi, pictures remembered	Ref	-0.03 (-0.06; 0.00)	-0.06 (-0.09; -0.03)	-0.05 (-0.08; -0.02)
PLTd, pictures remembered	Ref	-0.06 (-0.10; -0.01)	-0.11 (-0.16; -0.06)	-0.09 (-0.13; -0.05)
Barthel, index	Ref	-0.01 (-0.02; 0.01)	-0.06 (-0.07; -0.04)	-0.04 (-0.06; -0.03)
IADL, points	Ref	0.00 (-0.02; 0.03)	-0.05 (-0.08; -0.03)	-0.04 (-0.07; -0.02)
Fully adjusted²				
Stroop, seconds	Ref	0.58 (0.24; 0.92)	0.61 (0.26; 0.97)	0.59 (0.28; 0.91)
LDCT, digits coded	Ref	0.00 (-0.07; 0.08)	-0.05 (-0.13; 0.03)	-0.08 (-0.16; -0.01)
PLTi, pictures remembered	Ref	-0.03 (-0.06; 0.00)	-0.06 (-0.09; -0.02)	-0.05 (-0.08; -0.02)
PLTd, pictures remembered	Ref	-0.05 (-0.10; 0.00)	-0.10 (-0.15; -0.05)	-0.09 (-0.13; -0.04)
Barthel, index	Ref	-0.01 (-0.03; 0.01)	-0.06 (-0.08; -0.04)	-0.05 (-0.06; -0.03)
IADL, points	Ref	0.00 (-0.02; 0.02)	-0.05 (-0.08; -0.03)	-0.05 (-0.07; -0.02)

Abbreviations: LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education.

²=sex, age, country, education, body mass index, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, treatment arm, systolic blood pressure, apolipoprotein E.

Supplementary table 7. Association between individual metabolites of the MetaboHealth score and cognitive decline.

	Metabolites Z-score (per SD)														
	XXL-VLDL-L Estimate (95% CI)	S-HDL-L Estimate (95% CI)	VLDL-D Estimate (95% CI)	PUFA/FA Estimate (95% CI)	Glc Estimate (95% CI)	Lac Estimate (95% CI)	His Estimate (95% CI)	Ile Estimate (95% CI)	Leu Estimate (95% CI)	Val Estimate (95% CI)	Phe Estimate (95% CI)	AcAce Estimate (95% CI)	Alb Estimate (95% CI)	GlycA Estimate (95% CI)	
Cognitive test															
Minimally adjusted¹															
Stroop, seconds	-0.03 (- 0.16; 0.11)	-0.24 (- 0.38; - 0.10)	-0.10 (- 0.23; 0.04)	-0.08 (- 0.22; 0.06)	0.03 (- 0.11; 0.17)	0.00 (- 0.14; 0.14)	-0.23 (- 0.37; - 0.09)	-0.05 (- 0.19; 0.09)	-0.10 (- 0.24; 0.04)	-0.20 (- 0.34; - 0.06)	-0.10 (- 0.24; 0.04)	-0.05 (- 0.20; 0.10)	-0.09 (- 0.23; 0.04)	0.03 (- 0.11; 0.17)	
LDCT, digits coded	-0.01 (- 0.04; 0.02)	0.02 (- 0.01; 0.02)	-0.01 (- 0.04; 0.02)	-0.02 (- 0.05; 0.01)	0.02 (- 0.01; 0.05)	0.02 (- 0.01; 0.05)	0.04	0.04	0.05	0.04	-0.02 (- 0.05; 0.02)	0.04	-0.01 (- 0.05; 0.02)	-0.01 (- 0.04; 0.02)	
PLTi, pictures remembered	-0.003 (- 0.02; 0.01)	0.00 (- 0.01; 0.01)	0.00 (- 0.00; 0.03)	0.02 (- 0.01; 0.02)	-0.01 (- 0.02; 0.00)	0.01 (- 0.02; 0.02)	0.00 (- 0.00; 0.01)	0.00 (- 0.01; 0.01)	0.00 (- 0.01; 0.01)	0.01 (- 0.02; 0.02)	-0.01 (- 0.02; 0.01)	0.00 (- 0.02; 0.01)	0.00 (- 0.01; 0.01)	-0.01 (- 0.02; 0.00)	
PLTd, pictures remembered	0.01 (- 0.003; 0.03)	0.00 (- 0.02; 0.02)	0.02 (- 0.00; 0.04)	0.01 (- 0.00; 0.03)	-0.02 (- 0.01; 0.00)	0.03	0.03	0.03	0.03	0.00 (- 0.02; 0.05)	0.00 (- 0.02; 0.05)	-0.01 (- 0.02; 0.01)	-0.01 (- 0.03; 0.01)	-0.01 (- 0.03; 0.01)	
Barthel, index	-0.01 (- 0.02; - 0.01)	0.01	0.00 (- 0.01; 0.01)	0.00 (- 0.01; 0.04)	-0.01 (- 0.01; 0.01)	-0.01 (- 0.02; 0.00)	0.01	0.00 (- 0.01; 0.01)	0.01	0.02	0.00 (- 0.01; 0.01)	-0.01 (- 0.01; - 0.01)	0.00 (- 0.01; 0.01)	-0.02 (- 0.03; - 0.01)	
IADL, points	-0.01 (- 0.02; 0.00)	0.00 (- 0.01; 0.02)	0.01 (- 0.00; 0.02)	-0.02 (- 0.03; - 0.01)	0.01 (- 0.00; 0.02)	0.02 (- 0.01; 0.03)	0.02 (- 0.02; 0.03)	0.01	0.04	0.03	0.01	-0.02 (- 0.02; - 0.04)	0.00 (- 0.01; 0.01)	-0.02 (- 0.03; - 0.01)	
Fully adjusted²															
Stroop, seconds	-0.03 (- 0.17; 0.11)	-0.24 (- 0.38; - 0.10)	-0.10 (- 0.24; 0.04)	-0.09 (- 0.23; 0.05)	-0.001 (- 0.15; 0.15)	0.03 (- 0.11; 0.17)	-0.24 (- 0.38; - 0.10)	-0.04 (- 0.18; 0.10)	-0.10 (- 0.24; 0.05)	-0.19 (- 0.33; - 0.05)	-0.11 (- 0.25; 0.03)	-0.04 (- 0.19; 0.10)	-0.09 (- 0.23; 0.05)	0.03 (- 0.11; 0.17)	
LDCT, digits coded	-0.01 (- 0.04; 0.02)	0.02 (- 0.01; 0.05)	-0.01 (- 0.02; 0.04)	-0.02 (- 0.05; 0.02)	0.02 (- 0.01; 0.06)	0.02 (- 0.01; 0.05)	0.04	0.04	0.05	0.04	-0.01 (- 0.04; 0.02)	0.04	-0.01 (- 0.05; 0.02)	-0.01 (- 0.04; 0.02)	
PLTi, pictures remembered	-0.003 (- 0.02; 0.01)	0.00 (- 0.02; 0.01)	0.00 (- 0.02; 0.03)	0.02 (- 0.01; 0.01)	-0.01 (- 0.02; 0.00)	0.01 (- 0.01; 0.02)	0.00 (- 0.00; 0.01)	0.00 (- 0.01; 0.01)	0.00 (- 0.01; 0.01)	0.01 (- 0.02; 0.02)	-0.01 (- 0.02; 0.02)	0.00 (- 0.02; 0.02)	0.00 (- 0.01; 0.01)	-0.01 (- 0.02; 0.00)	
PLTd, pictures remembered	0.01	0.01	0.02	0.03	0.01	-0.02 (- 0.01; 0.03)	0.02	0.03	0.03	0.03	0.00 (- 0.02; 0.05)	0.00 (- 0.02; 0.05)	-0.01 (- 0.02; 0.02)	-0.01 (- 0.03; 0.01)	-0.01 (- 0.03; 0.01)
Barthel, index	-0.01 (- 0.02; 0.01)	0.01	0.00 (- 0.02; 0.01)	0.00 (- 0.01; 0.04)	-0.01 (- 0.01; 0.03)	-0.01 (- 0.02; 0.00)	0.01	0.00 (- 0.01; 0.01)	0.01	0.02	0.00 (- 0.01; 0.01)	-0.01 (- 0.02; 0.01)	0.00 (- 0.01; 0.01)	-0.02 (- 0.03; - 0.01)	
IADL, points	-0.01 (- 0.02; 0.00)	0.01 (- 0.01; 0.02)	0.01 (- 0.00; 0.02)	-0.02 (- 0.04; - 0.01)	0.01 (- 0.00; 0.02)	0.02	0.01	0.03	0.03	0.01	-0.01 (- 0.02; - 0.04)	0.00 (- 0.01; 0.02)	-0.02 (- 0.03; - 0.01)	-0.01 (- 0.03; 0.01)	

Abbreviations: XXL-VLDL-L = total lipids in chylomicrons and extremely large VLDL; S-HDL-L = total lipids in small HDL; VLDL-D = mean diameter for VLDL particles, PUFA/FA = ratio of polyunsaturated fatty acids to total fatty acids (%); Glc = glucose; Lac = lactate; His = histidine; Ile = isoleucine; Leu = leucine; Val = valine; Phe = phenylalanine; AcAce = acetoacetate; Alb = albumin; GlycA = glycoprotein acetyls; LDCT = Letter-Digit Coding Test; PLTi = Picture-Word Learning test immediate; PLTd = Picture-Word Learning Test delayed; IADL = Instrumental activities of daily living. ¹=sex, age, country, education. ²=sex, age, country, education,

body mass index, history of cardiovascular disease, history of diabetes, history of myocardial infarct, smoking, alcohol intake, treatment arm, systolic blood pressure, apolipoprotein E. Items written in bold indicate that the 95% confidence interval does not contain zero.