

Supplementary Table 1: Spectral Biomarkers identified by corroboration of independent statistical tests to identify osteoarthritis vs control class discriminating peaks. Features in agreement under 3 and 4 tests (no. of hits) are displayed alongside the corresponding spectral position and attributed chemical vibration. ^{18,19 18,28 53}

Hits	Spectromics Data Point	Wavenumber (cm ⁻¹)	Modality	1 st Derivative Attributed Peak	2 nd Derivative Attributed Peak	Assignment
4*	373 – 381	1342.64 – 1334.18	Raman	-	-	CH, GAGs at 1342 cm ⁻¹
4*	733 – 738	946.56 – 940.82	Raman	-	-	C–C deformation of aggrecan / C–O–C stretching of GAGs at 937 – 941 cm ⁻¹
4*	806 – 808	862.16 – 859.83	Raman	-	-	C–C Stretching, Proline, Collagen at 856 – 859 cm ⁻¹
4*	1011 / 1015	617.056 / 10819.9	Raman / NIR-SWIR	-	-	Spectromics Artefact: Raman to NIR-SWIR transition
4*	1164, 1168	7158.55, 7094.38	NIR-SWIR	7174.78	7062.73	O–H Stretching (1st Overtone) at 7280 - 6040, 7460 - 6780 cm ⁻¹
4*	1207 – 1211	6524.79 – 6471.57	NIR-SWIR	-	6254.84	N–H stretch (-CONH, 1st overtone) at 6352 cm ⁻¹
4*	1313 – 1316	5356.95 – 5333.09	NIR-SWIR	5342.02	5280.17	Bound & Free water at 5200 cm ⁻¹
4*	1443 – 1449	4402.48 – 4366.62	NIR-SWIR	4408.17, 4420.64	4372.55, 4384.47	C–H bend (protein, 2nd overtone), at 4350 cm ⁻¹
4*	1500 – 1503	4084.29 – 4068.83	NIR-SWIR	4073.97, 4058.60	4068.83, 4073.97	No precedent
3*	106, 107	1616.69, 1615.69	Raman	-	-	Amide I at 1612–1696 cm ⁻¹
3*	245 – 247	1475.99 – 1473.94	Raman	-	-	CH₂ deformation/scissoring; protein & lipids at 1441 – 1460 cm ⁻¹
3*	264 – 269	1456.43 – 1451.27	Raman	-	-	CH₂/CH₃ scissoring; collagen & other protein at 1451 cm ⁻¹
3*	455 – 463	1255.22 – 1246.61	Raman	-	-	C–N stretching (Amide III) at 1230 – 1280 cm ⁻¹
3*	575 – 577	1124.38 – 1122.17	Raman	-	-	Pyranose ring at 1127 – 1163 cm ⁻¹
3*	629, 630	1064.33, 1063.21	Raman	-	-	SO₃⁻ stretching in sulphated GAGs, PGs at 1060 – 1064 cm ⁻¹
3*	1083, 1084	8767.51, 8744.18	NIR-SWIR	8719.98	8577.59	C–H stretching (2nd overtone) at 8820 – 8060, 8695 - 8197 cm ⁻¹