

1 Appendix 1: Search strategy

Medline Search, conducted 08/02/2023

("Attitude"[MeSH Terms] OR "Attitude to Health"[MeSH Terms] OR ("knowledge"[MeSH Terms] OR "knowledge"[All Fields] OR "knowledge s"[All Fields] OR "knowledgeability"[All Fields] OR "knowledgeable"[All Fields] OR "knowledgeably"[All Fields] OR "knowledges"[All Fields]) OR ("Help"[All Fields] AND ("seeking"[All Fields] OR "seeks"[All Fields])) OR "Self-Management"[MeSH Terms] OR "adaptation, psychological"[MeSH Terms] OR ("illness behaviour"[All Fields] OR "illness behavior"[MeSH Terms] OR ("illness"[All Fields] AND "behavior"[All Fields]) OR "illness behavior"[All Fields]) OR ("experience"[All Fields] OR "experience s"[All Fields] OR "experiences"[All Fields]) OR ("qualities"[All Fields] OR "quality"[All Fields] OR "quality s"[All Fields]) OR "Role"[MeSH Terms] OR "Self Concept"[MeSH Terms] OR "Self Efficacy"[MeSH Terms] OR ("belief s"[All Fields] OR "culture"[MeSH Terms] OR "culture"[All Fields] OR "belief"[All Fields] OR "beliefs"[All Fields]) OR ("believability"[All Fields] OR "believable"[All Fields] OR "believed"[All Fields] OR "believer"[All Fields] OR "believers"[All Fields] OR "believing"[All Fields] OR "culture"[MeSH Terms] OR "culture"[All Fields] OR "believe"[All Fields] OR "believes"[All Fields]) OR (("illness"[All Fields] OR "illness s"[All Fields] OR "illnesses"[All Fields]) AND ("percept"[All Fields] OR "perceptibility"[All Fields] OR "perceptible"[All Fields] OR "perception"[MeSH Terms] OR "perception"[All Fields] OR "perceptions"[All Fields] OR "perceptual"[All Fields] OR "perceptive"[All Fields] OR "perceptiveness"[All Fields] OR "percepts"[All Fields])) OR (("patient s"[All Fields] OR "patients"[MeSH Terms] OR "patients"[All Fields] OR "patient"[All Fields] OR "patients s"[All Fields]) AND ("perspective"[All Fields] OR "perspective s"[All Fields] OR "perspectives"[All Fields])) OR ("coherence"[All Fields] OR "coherences"[All Fields] OR "coherencies"[All Fields] OR "coherency"[All Fields] OR "coherent"[All Fields] OR "coherently"[All Fields]) OR (("therapeutics"[MeSH Terms] OR "therapeutics"[All Fields] OR "treatments"[All Fields] OR "therapy"[MeSH Subheading] OR "therapy"[All Fields] OR "treatment"[All Fields] OR "treatment s"[All Fields]) AND "initiation"[All Fields]) OR ("expect"[All Fields] OR "expectable"[All Fields] OR "expectance"[All Fields] OR "expectant"[All Fields] OR "expectative"[All Fields] OR "expected"[All Fields] OR "expecting"[All Fields] OR "expects"[All Fields] OR "motivation"[MeSH Terms] OR "motivation"[All Fields] OR "expectancies"[All Fields] OR "expectancy"[All Fields] OR "expectation"[All Fields] OR "expectations"[All Fields]) OR ("expect"[All Fields] OR "expectable"[All Fields] OR "expectance"[All Fields] OR "expectant"[All Fields] OR "expectative"[All Fields] OR "expected"[All Fields] OR "expecting"[All Fields] OR "expects"[All Fields] OR "motivation"[MeSH Terms] OR "motivation"[All Fields] OR "expectancies"[All Fields] OR "expectancy"[All Fields] OR "expectation"[All Fields] OR "expectations"[All Fields]) OR ("Patient Preference"[MeSH Terms] OR ("decision making, shared"[MeSH Terms] OR ("decision"[All Fields] AND "making"[All Fields] AND "shared"[All Fields]) OR "shared decision making"[All Fields] OR ("shared"[All Fields] AND "decision"[All Fields] AND "making"[All Fields])) OR ("physical examination"[MeSH Terms] OR ("physical"[All Fields] AND "examination"[All Fields]) OR "physical examination"[All Fields] OR "physical"[All Fields] OR "physically"[All Fields] OR "physicals"[All Fields]) OR "somat*"[All Fields] OR "non-specific"[All Fields] OR ("complaint"[All Fields] OR "complained"[All Fields] OR "complaints"[All Fields]) OR ("psychosomatic"[All Fields] OR "psychosomatacal"[All Fields] OR "psychosomatically"[All Fields] OR "psychosomatics"[All Fields] OR ("psychophysiologic disorders"[MeSH Terms] OR ("psychophysiologic"[All Fields] AND "disorders"[All Fields]) OR "psychophysiologic disorders"[All Fields])) OR ("pain"[MeSH Terms] OR "pain"[All Fields]) OR ("diagnosis"[MeSH Subheading] OR "diagnosis"[All Fields] OR "symptoms"[All Fields] OR "diagnosis"[MeSH Terms] OR "symptom"[All Fields] OR "symptom s"[All Fields] OR

"symptomes"[All Fields])) AND ("General Practice"[MeSH Terms] OR "General Practitioners"[MeSH Terms] OR "Family Practice"[MeSH Terms] OR "Primary Health Care"[MeSH Terms] OR "physicians, primary care"[MeSH Terms] OR "Ambulatory Care"[MeSH Terms]) AND ("Depressive Disorder"[MeSH Terms] OR "depressive disorder, major"[MeSH Terms] OR "Depression"[MeSH Terms])

Psycinfo Search, conducted 08/02/2023

(general practice or GP or family practice or primary health care or physicians primary care or ambulatory care)

AND

(depression or depressive disorder or depressive symptoms or major depressive disorder)

AND

(attitude or knowledge or help seeking or self-management or adaptation or illness behavior or experience or belief or illness perception or patient perspective or coherence or treatment initiation or expect or shared decision making or patient preference or symptom or complaint)

3 Appendix 2:

Authors/ Year	Study design	Number of participants	Population: Definition of depression	Population: Special Group	Context: Region	Context: Primary care	Concept
Johnson, 1973	Longitudinal, Interview	73	Diagnosis by GP	.	UK	GP	Number of consultations Satisfaction Relationship
Widmer, 1978	Longitudinal, Case-control-matching, Patient records	154	Diagnosis by GP	.	US	FP	Consultation frequency Chief complaint
Widmer RB, Cadoret RJ, 1979	Case-control-matching, Patient records	43	Diagnosis by GP	.	US	FP	Consultation frequency
Cadoret et al., 1980	Case-control-matching, Medical records	234	Diagnosis by GP	.	US	FPC	Consultation frequency
Johnson DA, 1981	Longitudinal, Interview	117	BDI \geq 11	.	UK	GP	Compliance Attitude to medication Doctor-patient-communication
Cremniter et al., 1982	Cross-sectional, Survey	682	Diagnosis by GP	.	F	GP	Presenting symptoms
Wilson et al., 1983	Longitudinal, Case-control-matching, Record audit	101	Diagnosis by GP	.	US	FPC	Consultation frequency
Oxman et al., 1983	Cross-sectional	126	Chart diagnosis	.	US	FPC	Number of calls
Katon et al., 1983	Cross-sectional	147	Zung depression test	.	US	Family medical center	Medical utilization
Foster JM, Gallagher D, 1986	Case-control-matching, Health and Daily Living Questionnaire	32	Diagnosis by GP	Elderly	US	VC	Coping
Diamond et al., 1987	Longitudinal, Patient charts	67	Diagnosis by GP	Women	US	FP	Reason for visit

Williamson PS, Yates WR, 1989	Cross-sectional, Review of symptoms, Patient chart	105	Inventory to diagnose depression, DSM III	.	US	FPC	Presenting complaints
Romans-Clarkson et al., 1990	Cross-sectional, Interview	314	Present State Examination (PSE)	Women	NZ	GP	Number of visits to GP Help seeking
Katon et al., 1990	RCT, Physician Review Form	124	National Institute of Mental Health Diagnostic Interview Schedule, DSM-III-R	.	US	PCC	Health care utilization
Gerber et al., 1992	Cross-sectional, Physician Recording Form	1055	HSCL-90 Depression Scale ≥ 7	.	US	GP	Presenting complaints
Scott AI, Freeman CP, 1992	RCT, Questionnaire	101	DSM-III, diagnostic interview	.	UK	GP	Satisfaction
Kirmayer et al., 1993	Observational, Interview	202	DIS, CES-D ≥ 16	.	Canada	FMC	Clinical presentation
Blanchard et al., 1994	Follow-up, Interview	96	Short-CARE, GMS-AGECAT	Older people	UK	GP	Declaration of symptoms
Callahan et al., 1994	Longitudinal, Interview	1711	CES-D ≥ 16	Elderly	US	GP	Health service use
Karlsson et al., 1995	Case control study	562	Present state examination interview	.	Finland	Primary care station	Frequent attendance
Lin et al., 1995	Longitudinal, Interview	155	Antidepressant prescription for depression	.	US	PCP	Adherence
Katon et al., 1995	RCT, Questionnaire	193	Diagnosis by GP	.	US	PCP	Adherence Quality rating
Tylee et al., 1995	Observational, Interview, Analyzed video-recordings	72	Diagnostic interview	Women	UK	GP	Mentions of symptoms

Brown et al., 1996	RCT, Health Locus of Control Scale	272	CES-D \geq 22, DIS	African American	US	PCP	Health locus of control
Gormley N, O'Leary D, 1998	Cross-sectional, Interview	100	HRSD \geq 16	.	Ireland	GP	Time to presentation
Cornwell J, Hull S, 1998	Cross-sectional, Data from records	90	Antidepressant prescription for depression	South Asian	UK	GP	Presenting complaints
Rubio Montañés et al., 1998	Case-control-matching, Questionnaire	209	Escala de Ansiedad y Depresión de Goldberg (EADG)	.	Spain	HCC	Over-attendance
Van Hook, 1999	Cross-sectional, Questionnaire	321	Medical Outcome Services (MOS), ICD-10	Women	US	PCC	Help seeking patterns
Kessler et al., 1999	Cross-sectional, Questionnaire	305	General Health Questionnaire \geq 3	.	UK	GP	Symptom interpretation
Arve et al, 1999	Cross-sectional	847	Zung depression test	70 years and older	Finland	GP	Self-perception of depression
Waza et al., 1999	Observational, Chart audit	189	Chart diagnosis	.	Japan/ US	FP	Complaints
Gilmore KA; Hargie O; 2000	Cross-sectional, Interview	8	Prior diagnosis	.	UK	GP	Satisfaction
Rost et al., 2000	Cross-sectional, Interview	240	Inventory to diagnose depression	.	US	PCP	Acceptability
Nichols GA, Brown JB, 2000	Cross-sectional, Questionnaire	1161	Positive screen/ chart diagnosis	.	US	FP	Being asked
Cooper et al., 2000	Cross-sectional, Survey	76	CESD \geq 16	.	US	PCC	Attitude to care Importance of different aspects of care
Katon et al., 2000	RCT, Automated databases	1599	SCID	.	US	PCC	Consultation frequency Adherence

Dwight-Johnson et al., 2000	Cross-sectional, Interview	1187	CIDI	.	US	PCC	Preference
Bedi et al., 2000	Partially randomised preference trial, Preference between antidepressants or psychotherapy given by patient, Questionnaire	323	Diagnosis by GP	.	UK	GP	Preference Attitude to treatment
Demyttenaere et al., 2001	Longitudinal, Antidepressant Compliance Questionnaire	272	Diagnosis by GP	.	US	GP	Compliance Informing physician
Andrew et al., 2001	Case-control-matching, Primary care records	90	Seasonal affective Disorder (SPAQ), DSM IV	.	UK	GP	Consultation rate
Meredith et al., 2001	Cross-sectional, Components of Primary Care Index, Medical Outcomes Study, Consumer Assessment of Health Plans Survey, Data from patient surveys	1104	CIDI	.	US	primary care visitors in managed care organizations	Satisfaction Relationship Quality of care indicators
Rogers et al., 2001	Qualitative interviews	27	Diagnosis by GP	.	UK	GP	Experience of care
Brown et al., 2001	Cross-sectional, Brief COPE Inventory, Consumer Assessment of Health Plans Survey, Modified Illness Perceptions Questionnaire	41	Diagnosis by GP	.	US	PCC	Coping Adherence Illness model Controllability

Andrews G, Carter GL, 2001	Cross-sectional, Interview, UK Survey of Psychiatric Morbidity	10641	CIDI	.	Australia	GP	Number of consultations Perceived need for care Report of received treatment
Dwight-Johnson et al., 2001	RCT, Patient assessment questionnaire	1187	CIDI	.	US	PCC	Preference
O'Connor et al., 2001	Cross-sectional, Interview	1021	Diagnosis by GP	Elderly patients	Australia	GP	Symptom disclosure
McKelvey et al., 2001	Cross-sectional, Summary sheets	2792	CES-D, DSI-SS	.	US	GP	Chief complaints
Webster et al., 2001	Cross-sectional, Questionnaire	574	EPDS	Women postpartum	Australia	GP	Visiting health care provider Satisfaction
Ronalds et al., 2002	Follow-up, Interview	148	Psychiatric Assessment Schedule	.	UK	GP	Consultation rate
Holopainen D, 2002	Cross-sectional, Interview	7	Recruited from support group	Women postpartum	Australia	GP	Experience of seeking help
Shin et al., 2002	Focus group	70	Self-identification as depressed	Korean immigrants	US	PC	Help seeking
Maidment et al., 2002	Cross-sectional	67	Antidepressant prescription for depression	65 and older	UK	GP	Adherence (Global adherence measure (GAQ)) Patient education (the questionnaire on patient education (QPE)) Insight (schedule for assessing the three components of insight) Beliefs about medication (beliefs about medicine questionnaire (BMQ))

Orlando M, Meredith LS, 2002	Longitudinal, Consumer Assessment of Health Plans Survey, Patient provider relationship ratings, Data from the patient surveys	697	CIDI	.	US	primary care visitors in managed care organizations	Satisfaction Relationship Quality of care indicators
Pollock K, Grime J, 2002	Follow-up, Interview	62	Diagnosis by GP, Depression Alliance	.	UK	GP	Entitlement to time
Manning C, Marr J, 2003	Cross-sectional, Survey	1010	Depression Alliance	.	UK	GP	Lifestyle choices Reason for stopping medication Experience of recurring depression
Gabbay et al., 2003	RCT, Questionnaire	464	Diagnosis by GP	.	UK	GP	Problem formulation Agreement with GP on problem
Solberg et al., 2003	Follow-up, Questionnaire, Shortform functional status survey	274	Chart diagnosis	.	US	PCC	Self-efficacy Adherence Satisfaction Experience of treatment
Gask et al., 2003	Cross-sectional, Interviews	27	Diagnosis by GP	.	UK	GP	Adherence Quality of care Experience of treatment
Wills CE, Holmes-Rovner M, 2003	Cross-sectional, Satisfaction With Decision Scale, Decision Conflict Scale, Physician Participatory Decision-Making Style Scale, Questionnaire	97	Antidepressant prescription for depression	.	US	PCP	Decision-making Satisfaction with decision Knowledge
Parker et al., 2003	Cross-sectional	638	18-item questionnaire	.	Australia	GP	Attribution style

Templeton et al., 2003	Longitudinal, Interviews, Focus groups	20	EPDS	Minority ethnic women	UK	GP	Experience of treatment
Corrigan et al., 2003	Cross-sectional, Survey	230	GDS-screen	Older adults	US	PCP	Discuss depression
Lin et al., 2003	RCT, Follow-up, Interview, Questionnaire	386	Antidepressant prescription for depression	.	US	PCC	Self-management Adherence Attitude to medication Confidence in managing side effects
Van Voorhees et al., 2003	Cross-sectional, Questionnaire	881	CIDI	.	US	PCP	Health care utilization Acceptance of care
Yeung et al., 2004	Cross-sectional, Explanatory Model of Interview Catalogue	40	CBDI	Chinese Americans	US	PCC	Chief complaint Help seeking Perceived causes Stigma
Ford et al., 2004	Case-control-matching, Medical record data	300	Chart diagnosis	.	US	HC	Health care utilization
Yamada et al., 2004	Cross-sectional	162	MINI	.	Japan	PCC	Complaints Consultation frequency
Shvartzman et al., 2005	Cross-sectional, Interview	2507	MINI	.	Israel	PCC	Health care utilization
Brook et al., 2005	RCT, Patient medication records, Electronic pill container	135	Antidepressant prescription for depression	.	The Netherlands	GP	Adherence
Pyne et al., 2005	RCT, Interview	211	IDD/ DSM IV	.	US	PCP	Acceptability of medication
Al-Windi et al., 2005	Cross-sectional	1055	Göteborg quality of life instrument	.	Sweden	GP	Visits to GP
Ruoff G, 2005	Follow-up, Patient charts	103	Diagnosis by GP	.	US	FP	Discontinuation of medication

Brown et al., 2005	Cross-sectional, Observational, Beliefs about Medicines Questionnaire	192	Antidepressant prescription for depression	.	US	FP	Beliefs about medication
Löwe et al., 2006	Cross-sectional, Interview	178	SCID /DSM IV	.	D	internal medicine outpatient clinic	Self-management Preference
Bogner et al., 2006	Two-stage sampling, Pill counts	228	SCID /DSM IV	Older adults	US	PCP	Adherence
Rhodes et al., 2006	Cross-sectional, Survey	36984	CIDI/ DSM IV	.	Canada	primary medical care	Health service use
Lawrence et al., 2006	Qualitative interviews	110	HADS	Older adults	UK	PCP	Concept of depression
Givens et al., 2006	Qualitative interviews	42	CES-D/ HDRS	Older adults	US	PCP	Attitude to medication
Gum et al., 2006	RCT, Interview	1602	SCID /DSM IV	Older adults	US	PCC	Preference
Menchetti et al., 2006	Cross-sectional, Two-phase, Clinical charts	1854	WHO-Checklist for ICD-10	Elderly patients	Italy	PCP	Attendance frequency
Akerblad et al., 2006	RCT, Sertraline-/Desmethylsertraline-Level, Self-report, Scheduled visits	940	Prescription of antidepressants for depression	.	Sweden	GP	Adherence
Backenstrass et al., 2006	Cross-sectional, Questionnaire	607	PHQ-9	.	D	GP	Preference
Finucane A, Mercer SW, 2006	Longitudinal, Interview	13	BDI-II, ICD-10 chart diagnosis	.	UK	GP	Preconceptions about treatment Expectations Acceptability Experience of treatment

Clever et al., 2006	Prospective cohort, Cross-sectional, Rating the involvement	1607	CIDI/ DSM IV	.	US	PCP	Involvement in decision-making
Schneider et al., 2006	Cross-sectional, Fragebogen zur Erhebung von Kontrollüberzeugung zu Krankheit und Gesundheit, Autonomy Preference Index	234	HADS	.	D	GP	Health locus of control Preference for involvement
Mellor et al., 2006	Cross-sectional, Interview, Depression Treatment Satisfaction Questionnaire	31	SCID	≥65 years old	Australia	aged-care facility	Experience of care Awareness of care Relationship Satisfaction
Eilat-Tsanani et al., 2006	Follow-up, Interview	527	EPDS	Women postpartum	Israel	FP	Consultation pattern
MsCracken et al., 2006	Two-phase interview, Client Service Receipt Inventory	427	BDI, ICD-10	.	Ireland Finland Norway Spain UK	PCP	Service use
von Knorring et al., 2006	RCT, Sertraline- /Desmethylsertraline- Level, Self-report, Scheduled visits	1031	Diagnosis by GP (DSM- IV)	.	Sweden	GP	Adherence
Johnson et al., 2006	Cross-sectional, Question	1140	CIDI/ DSM IV/ chart diagnosis	.	US	PCC	Preference
Badger F, Nolan P, 2006	Cross-sectional, Semi-structured questionnaire	60	Diagnosis by GP	.	UK	PCC	Medication concordance Beliefs about medication

							Role of relationship Experience of medication
Lawrence et al., 2006	Qualitative interviews	110	HADS	Older adults	UK	PCP	Help seeking
Mohr et al., 2006	Cross-sectional, Perceived Barriers to Psychotherapy Scale	290	PHQ >10	.	US	PCC	Perceived barriers to psychotherapy
Brook et al., 2006	Prospective cohort, Follow-up, Electronic pill container, Drug Attitude Inventory	119	Prescription for tricyclic antidepressant for depression	.	The Netherlan ds	GP	Adherence Drug attitude
Saver et al., 2007	Cross-sectional, Interview	15	Diagnosis by GP	.	US	PCC	Understanding of depression Sources of information Participation in treatment decisions Barriers to treatment and information
Loh et al., 2007	Follow-up, Patient participation scale, Scale regarding adherence	207	Diagnosis by GP	.	D	GP	Treatment adherence Participation in treatment decisions
Saur et al., 2007	Cross-sectional	105	SCID	Older adults	US	Academic group practice	Satisfaction with care
Sleath et al., 2007	Observational, Interview audiotapes	40	Prescribed antidepressants for depression	Veterans	US	general medicine outpatient clinic	Ask and talk about antidepressants
Badger F, Nolan P, 2007	Cross-sectional, Interview	60	Diagnosis by GP	.	UK	GP	Experience of treatment Attributing recovery
Pollock K, 2007	Cross-sectional, Interview	62	Diagnosis by GP	.	UK	GP	Maintaining face in medical consultations

Stecker et al., 2007	Cross-sectional	29	Chart diagnosis	.	US	Family medical center	Filling prescription Initiation of psychotherapy Stigma (The Stigma Questionnaire) Beliefs about psychotherapy (The Beliefs about Psychotherapy Scale)
Badger F, Nolan P, 2007	Cross-sectional, Interview	60	Diagnosis by GP	.	UK	GP	Self-chosen therapies
Weich, 2007	Cross-sectional, Interview, Attitudes to depression and it's treatment questionnaire	866	CIDI/ ICD-10	.	UK	GP	Adherence Attitudes to depression and its treatment
Cornford et al., 2007	Cross-sectional, Interview	23	HADS	.	UK	GP	Beliefs about depression
Johnston et al., 2007	Cross-sectional, Interview	111	Diagnosis by GP	.	UK	GP	Goals for the management of depression Acceptance of diagnosis
Dobscha et al., 2007	RCT, Questionnaire	314	PHQ	Veterans	US	VC	Preference
Leydon et al., 2007	Qualitative interviews	17	Prescription of antidepressants for depression	.	UK	GP	Barriers and facilitators to discontinuation of antidepressants
Ferrari et al., 2008	Case-control study, Database	50	SCID, DSM IV	.	Italy	PCC	Attendance frequency
Druss et al., 2008	Cross-sectional, Survey	30801	CIDI-SF, DSM-III-R	.	US	PC	Access to primary care
Frémont et al., 2008	Cross-sectional, survey	598	Diagnosis by GP	.	F	GP	Causal attribution Satisfaction Therapeutic alliance

							Acceptance of diagnosis and treatment Experience of treatment Communication behaviors
Ghods et al., 2008	Cross-sectional	108	Medical Outcomes Study SF-12	African Americans and Whites	US	Group practices and community health centers	
Glaesmer et al., 2008	Cross-sectional, Amount of doctor and hospital visits, Scale for the Assessment of Illness Behavior	1185	PHQ-9	≥50 years of age	D	GP	Health service use Illness behavior
Bazargan et al., 2008	Cross-sectional, Survey	415	PHQ-9	African American and hispanic patients	US	PCC	Use of complementary and alternative medicine
Dernovsek et al., 2008	Cross-sectional, Questionnaire	391	Prescription of antidepressants for depression	.	Slovenia	public health center	Compliance
Damush et al., 2008	Cross-sectional, Questionnaire	500	PHQ-9 ≥10	Musculoskeletal pain	US	PCC	Self-management
Dowrick et al., 2008	Cross-sectional, Interview	100	CES-D ≥16	.	UK	GP	Resilience
Bogner et al., 2008	Cross-sectional, Interview	33	CES-D	≥65 years of age	US	PCP	Illness model
Russell J, Kazantzis, 2008	Cross-sectional, Medication Adherence Report Scale, Beliefs about Medication Questionnaire	85	Diagnosis by GP	.	NZ	GP	Adherence Beliefs about medication

Turner et al., 2008	Cross-sectional, Interview	27	EPDS	Women	UK	GP	Views on medication Preference
Wittkamp et al., 2008	Cross-sectional, Interview	17	PHQ, SCID, HDRS	.	The Netherlands	GP	Views on screening and treatment Acceptance of diagnosis
Chew-Graham et al., 2009	RCT, Interview	28	Prescription of antidepressants for depression	Women postpartum	UK	GP	Disclosure of symptoms Understanding postnatal depression Barriers of disclosure
Bogner et al., 2009	Qualitative interviews	102	CES-D	65 and older	US	PCP	Views regarding antidepressant medication
van Geffen et al., 2009	Retrospective, Contact file, Pharmacy dispensing registration database	965	Prescription of antidepressant for depression	.	The Netherlands	GP	Contact with GP Initiation of antidepressant treatment and filing prescriptions
Hérique A, Kahn JP, 2009	Retrospective, Cross-sectional, Questionnaire	632	Diagnosis by GP	.	F	GP	Compliance to antidepressants
Wittink et al., 2009	Cross-sectional, Interview	47	CES-D	Older African Americans	US	PCP	A Faith-Based Explanatory Model of Depression
Beattie et al., 2009	Longitudinal, Interview	24	Diagnosis by GP	.	UK	GP	Completion of online-CBT Expectations of online CBT Experience of online CBT
Raue et al., 2009	Longitudinal, Randomized, Care manager records, Participants' reports, Questionnaire	60	SCID	.	US	PCP	Adherence Initiation Expectation Preference
Menchetti et al., 2009	Two-phase, PCP records, Patient chart	250	WHO ICD-10 Symptom Checklist for Depression	.	Italy	PCP	Visiting frequency Presenting complaints
Suija et al., 2009	Cross-sectional, Registration forms	1094	CIDI	.	Estonia	FP	Visits to FD

Danielsson et al., 2009	Cross-sectional, Interview	20	ICD-10, chart diagnosis	.	Sweden	health care center	Source and course of depression
Hodges et al., 2009	Cross-sectional, Questionnaire	100	Diagnosis by GP	Cancer patients	UK	GP	Preference
Prins et al., 2009	Cross-sectional, Perceived Need for Care Questionnaire	622	CIDI	.	The Netherlands	primary care center	Perceived need for care
Vega et al., 2010	Cross-sectional, Stigma Checklist	200	PHQ-9 score ≥ 10	Latino patients	US	PCC	Stigma
Dickinson et al., 2010	Cross-sectional, Interview	36	Antidepressant prescription for depression	≥ 75 years of age	UK	GP	Understanding depression Views on long term use of antidepressants Barriers to discontinuation
Malpass et al., 2010	Longitudinal, Interview	10	PHQ-9, diagnosis by GP	.	UK	GP	Patients' experiences of using the PHQ-9 in primary care consultations
Barg et al., 2010	Follow-up (randomized), Interview	24	CES-D	Older patients	UK	PCC	Experience of treatment
Hansson et al., 2010	Cross-sectional, Open-ended question	303	Diagnosis by GP	.	Sweden	health care center	Belief about the cause of depression
van Geffen et al., 2010	Follow-up, Observational, Data from dispensing records, Questionnaire	110	Antidepressant prescription for depression	.	The Netherlands	GP	Initiation and continuation of antidepressant treatment Health beliefs
Palmer et al., 2010	Longitudinal, Interview	474	CESD ≥ 16	.	Australia	GP	Desires towards depression treatment
Newman et al., 2010	Cross-sectional, Interview	40	PHQ-9	Gay men	Australia	GP	Ascribed role of the GP

Gask et al., 2011	Cross-sectional, Interview	15	Diagnosis by GP	Pakistani women	UK	GP	Understanding persistence of depression Views on depression treatment
Mergl et al., 2011	RCT, Fragebogen zur Messung der Psychotherapiemotivation (Item 36)	145	CIDI, DSM IV	.	D	PCP	Attitude to psychotherapy preference
Malpass et al., 2011	Longitudinal, Interview, Consultation records	10	Antidepressant prescription for depression	.	UK	GP	Unvoiced agenda
Saito et al., 2011	Cross-sectional, Interview	5	Diagnosis by GP	.	Japan	PCP	Perceived triggers for depression Opinions on treatment
Fortney et al., 2011	Cross-sectional	395	PHQ 9	Veterans	US	VA medical center	Adherence Acceptability of antidepressants Health beliefs (The Depression Health Beliefs Inventory) Reasons for non-adherence
Callister et al., 2011	Qualitative interviews	20	Postpartum Depression Screening Scale, Spanish Version	Hispanic women postpartum	US	Community health center	Help seeking
Körner et al., 2011	Cross-sectional, Interview	40	PHQ 9	Gay men	Australia	GP	Understanding of depression Experience of depression
Prins et al., 2011	Follow-up, Perceived Need for Care Questionnaire	568	CIDI, DSM IV	.	The Netherlands	GP	Perceived need for care
Deen et al., 2011	RCT, Depression Health Beliefs Inventory, One Item regarding acceptability,	360	PHQ9 ≥ 12	.	US	PCP	Health beliefs Satisfaction with care Acceptability of treatment Patient-centeredness of care

	One scale regarding satisfaction, Experience of Care and Health Outcomes Survey						
Lynch et al., 2011	Cross-sectional, Beliefs about Depression Questionnaire	334	Diagnosis by GP	.	UK	GP	Beliefs about depression
Bell et al., 2011	Follow-up, Perceived Barriers Index	1054	PHQ-9	.	US	PCP	Barriers of disclosure
Elwy et al., 2011	Cross-sectional, Interview	30	PHQ-9 ≥ 11	.	US	PCP	Illness perception
van den Boogaard et al., 2011	Cross-sectional, Medical records, Causal attributions inventory	120	CIDI	.	The Netherlands	GP	Visits to GP Causal attribution
Gensichen et al., 2012	Cross-sectional, Interview	41	Diagnosis by GP	.	D	GP	Perspective on collaborative care
Kwong et al., 2012	Cross-sectional, Stigma assessment scale, Interview	42	PHQ-9	Chinese American	US	community health center	Help seeking Perceived cause of depression Perceived stigma Reasons that prevent individuals from participating in a depression treatment program
Chew-Graham et al., 2012	Cross-sectional, Interview	19	Diagnosis by GP	Older people	UK	GP	Accessing help Reasons to not present with depression
Hansson et al., 2012	Follow-up, Questionnaire	184	Diagnosis by GP, HADS	.	Sweden	primary health care center	Perceived improvement factors
Calderón Gómez et al., 2012	Observational, Discussion groups,	31	Diagnosis by GP	.	Spain	FP	Views on treatment

	Interview						
Cornford et al., 2012	Cross-sectional, Interview, Focus group	24	Diagnosis by GP	Opioid dependency	UK	GP	Causes of depression Experience of depression
Hansen et al., 2012	Cross-sectional, Focus group	19	PHQ-9	Low-income Latinos with diabetes	US	PCC	Help seeking Barriers to help seeking
Dwight Johnson et al., 2013	Cross-sectional, Survey	63	SCID	Older mexican men	US	PCC	Preference for treatment
Simmonds et al., 2013	Cross-sectional, Interview	30	PHQ2	Patients with coronary heart disease	UK	GP	Links between CHD and depression Experience of CHD and depression
Busch et al., 2013	Cross-sectional, Survey	2890	EURO-D-Skala ≥ 4	≥ 50 years of age	D	GP	Health service utilization
Bennett et al., 2013	Cross-sectional, Interview	41	CIDI	Chronic depression	UK	GP and practice nurses	Experience of care Treatment initiation Adherence Barriers to care
Kales et al., 2013	Observational, Follow-up, Brief Medication Questionnaire	188	Antidepressant prescription for depression	Old age	US	PCC	Adherence
Izquierdo et al., 2014	Two-stage, Interview	47	CIDI	Older Latinos	US	PCC	Experience of visits to PC for mental health
Keeley et al., 2014	Cross-sectional, Interview	30	Diagnosis by PCP, ICD-9	.	US	PCP	Conceptualization of depression Expectations of depression care Experience of depression care
DeJesus et al., 2014	Cross-sectional, Questionnaire	125	Diagnosis by GP	.	US	primary care site	Self-management strategies Satisfaction with CCM for depression Opinions of care managers

							Attitudes towards care management
Magnezi et al., 2014	Cross-sectional, Patient Activation Measure	278	PHQ-9	.	Israel	PCC	Patient activation
Solberg et al., 2014	Cross-sectional, Questionnaire	1168	Diagnosis by GP, PHQ-9 ≥ 7	.	US	PCC	Shared decision-making
Osei-Bonsu et al., 2014	Observational, Brief COPE, Interview	23	PHQ-9	Veterans	US	VC	Coping strategies
Serrano et al., 2014	Observational, Longitudinal, Simplified Medication Adherence Questionnaire, Drug Attitude Inventory	29	DSMIV, HADS ≥ 17	.	Spain	primary care center	Adherence Attitude to drugs
Fosgerau CF, Davidsen AS, 2014	Observational, Conversation analysis	28	Diagnosis by GP, ICD-10	.	Denmark	GP	Patients' perspectives on antidepressant treatment
Edwards et al., 2014	Cross-sectional, Questionnaire	1478	Diagnosis by GP	.	UK	GP	Satisfaction with telehealth use Advantages of telehealth use
Jaffray et al., 2014	Cross-sectional, Interview	29	Diagnosis by GP	.	UK	GP	Attitude to antidepressants
Waltz et al., 2014	Cross-sectional, Questionnaire	761	PHQ-9	Veterans	US	VC	Satisfaction Treatment preference
Holloway et al., 2015	Longitudinal, Open-ended question	57	PHQ-9	Vision-impaired	Australia	GP	Uptake of GP referral
Chen et al., 2015	RCT, Explanatory Model Interview Catalogue	190	PHQ-9 ≥ 10	Chinese American	US	community health center	Chief complaint Name of problem/ perceived cause of depression Experience of help sought
Knowles et al., 2015	Cross-sectional, Interview	31	PHQ-9 ≥ 10	Physical long-term condition	UK	PCP	Experience of collaborative care
Lynch et al., 2015	Longitudinal,	227	Diagnosis by GP	.	UK	GP	Beliefs about depression

	Beliefs about Depression Questionnaire						
Lee King et al., 2015	Cross-sectional, Treatment Beliefs Scale, Medication Beliefs Scale, Agnew Relationship Measure, Goldring Patient–Provider Scale, Interview	198, 29	Chart diagnosis	.	US	PCP	Treatment beliefs Relationship Experience of depression
Horevitz et al., 2015	Two-phase, Chart review, Interview	431, 16	Chart diagnosis, PHQ-9	Latinos	US	health care center	Taking up referral to behavioral health Relationship Factors influencing decision to take up referral
Knowles et al., 2015	Cross-sectional, Interview	36	PHQ-9 ≥ 10	.	UK	collaborative care	Experience of computerized CBT
Patel et al., 2015	Case-control study	142	SCID	.	UK	GP and practice nurses	Attendance
Larsen et al., 2016	Longitudinal, National register	1332	HADS-D ≥ 8	Heart disease	Denmark	GP	Contact with GP
von Faber et al., 2016	Qualitative interviews	38	Geriatric Depression Scale (GDS-15)	75 and older	The Netherlands	GP	Coping strategies
Elwy et al., 2016	Prospective observational study	271	PHQ-9	Veterans	US	VA PCC	Illness perceptions (the illness perceptions questionnaire - revised (IPQ-R))
Campbell et al., 2016	Cross-sectional, Questionnaire	761	PHQ-9	Veterans	US	VC	Care engagement Stigma Agreement with depression label Treatment preference

							Report of provider behavior Received treatment
Keller et al., 2016	Cross-sectional, Interview	24	PHQ-8, history of depression	Women	US	PCP	Experience of disclosure Barriers and facilitators to disclosure
Vuorilehto et al., 2016	Longitudinal, Monitoring records, Questionnaire	100	SCID/ DSM IV	.	Finland	GP	Visits to GP Adherence to treatment Initiation of treatment
Chen et al., 2016	Follow-up, Explanatory Model Interview Catalogue	168	CB-PHQ-9 ≥ 10	Chinese immigrants	US	health care center	Stigma
Magnani et al., 2016	Cross-sectional, Question	170	HDRS ≥ 13	.	Italy	PCP	Preference
Davis et al., 2016	Follow-up, Questionnaire	761	PHQ-9 ≥ 10	Veterans	US	VA collaborat ive care	Satisfaction with care Preference Experience of PC-related mental health care
Rossom et al., 2016	Observational, Question regarding care quality, Modified Patient Assessment of Chronic Illness Care	792	PHQ-9	.	US	PCC	Care rating Experience of patient-centered care
Sternke et al., 2016	Focus groups data	18	PHQ-9 ≥ 10	Chronic pain	US	VC and PCC	Experience of empathy
Wikberg et al., 2016	Recorded focus groups	9	Diagnosis by GP	.	Sweden	GP	Patients' experience of the use of MADRS-S in primary care
Moise et al., 2017	Cross-sectional, Morisky Medication Adherence Scale, Control Preference Scale	195	PHQ-8 ≥ 10	Hypertension	US	PCC	Adherence Preference for decision-making

Robinson et al., 2017	Longitudinal, Interview	29	Diagnosis by GP	.	UK	PCP	Explanations for discrepancies between changes in PHQ-9 and GRC
Saint Aurnalt et al., 2017	Cross-sectional	209	CES-D	Japanese immigrants	US	PC	Illness interpretation
Green et al., 2017	Cross-sectional, Interview, Focus groups	28, 12	PHQ-9 ≥ 10	Latinos	US	collaborative care	Adherence to antidepressants Receiving information about antidepressants Experience of taking medication
Gunn et al., 2018	Longitudinal, Survey	789	CES-D ≥ 10	.	Australia	GP	Health service use Self help
Stark et al., 2018	Cross-sectional, Interview	12	Geriatric Depression Scale (GDS) ≥ 6	Older patients	D	GP	Attitudes concerning depression Views on depression treatment
Lopez et al., 2018	Cross-sectional, Stigma Concerns About Mental Health Care Measure, Social Distance Measure, Latino Scale for Antidepressant Stigma Measure, Depression Knowledge Measure	319	PHQ-9	Hispanic women	US	PCC	Stigma Knowledge on depression
Dorow et al., 2018	Cross-sectional	641	Diagnosis by GP	.	D	PCP	Preference for treatment
Taylor et al., 2018	Cross-sectional, Interview	13	MINI	Older patients	UK	collaborative care	Views on collaborative care for depression
Waitzfelder et al., 2018	Observational, Health record data	241251	Diagnosis by GP, ICD-9	.	US	primary care site	Initiation of treatment
Jones et al., 2018	Cross-sectional, Survey	27362	ICD-9, chart diagnosis	.	US	VC	Number of clinic visits Experience of care
Vannoy et al., 2018	Cross-sectional, Observational,	77	SCID	Older men	US	PCP	Attitudes towards discussing suicide

	Interview						Patients' perspectives on what primary care providers can do to help prevent late-life suicide
Pung et al., 2018	Cross-sectional, Interview	16	PHQ-2	.	Australia	PCP	Patterns of mobile app use among patients with depressive symptoms
Avey et al., 2018	Cross-sectional, Questionnaire	125	PHQ-9 ≥10	Alaskan Native, American Indian	US	PCC	Preference
Gordon et al., 2018	Cross-sectional, Interview	16	Diagnosis by GP	Older people	UK	GP	Understanding of depression
Carmassi et al., 2018	Cross-sectional	75	SCID	.	Italy	GP	Attendance Empowerment (Empowerment in patients with Affective and Schizophrenic Disorders)
Canty et al., 2019	Cross-sectional, Interview	17	EPDS	Women postpartum	US	PCC	Barriers and facilitators to mental health care postpartum Experience with depressive symptoms
Aznar-Lou et al., 2019	Cross-sectional, Beliefs About Medicines Questionnaire, The Brief Illness Perception Questionnaire	263	Diagnosis by GP, SCID	.	Spain	GP	Beliefs about illness and medicine
Schomerus et al., 2019	Longitudinal, Self-Identification as Having Mental Illness Scale, Social Distance Scale, Questionnaire	188	PHQ-9 ≥8	.	D	GP	Help seeking intention Self-identification as having a mental illness Prejudice and discrimination Literacy for depression Perceived need for care
Liu et al., 2019	Cross-sectional, Medical records	2976	PHQ-8	Older patients with osteoarthritis	US	group health cooperative	Health care use

Aragonès et al., 2019	RCT, Questionnaire, Medical records	274	SCID	Musculoskeletal pain	Spain	primary care center	Health care resource use Adherence to program Satisfaction
Richards et al., 2019	Cross-sectional, Interview	37	PHQ-2	.	US	primary care center	Experience of being asked about suicidality
Backhaus et al., 2020	Cross-sectional, Question	641	Diagnosis by GP, PHQ-9 \geq 5	.	D	GP	Treatment preference
Pitrou et al., 2020	Cross-sectional	1624	Questionnaire based on DSM-IV	65 years and older	Canada	PCP	Satisfaction with care
Pirard et al., 2020	Cross-sectional, Questionnaire	454	HADS-D \geq 8	Survivors of terrorist attacks	F	GP	Mental health support use
Kloppe et al., 2020	Cross-sectional, Interview	26	PHQ-9	60 years and older	D	GP	Relief seeking style
Gomà-I-Freixanet et al., 2020	Case-control-matching, Medical records	238	Goldberg Anxiety–Depression Scale \geq 2	.	Spain	PCP	Self-initiated consultations
Heinz et al., 2021	Cross-sectional, Questionnaire, Depression Stigma Scale	430	Depression screening questionnaire, ICD 10	.	D	GP	Reported complaints Stigma
Cornwell et al., 2021	Cross-sectional	92535	PHQ-2	Veterans	US	integrated primary care	Initiation of treatment
Poole et al., 2022	Cross-sectional, Interview	41	Diagnosis by GP	Older patients, Long-term conditions	UK	GP	Experience of depression
Gartland et al., 2022	Prospective cohort study	1507	CES-D	Mothers	Australia	GP	Visits to GP Trust in PCP (Health Care Relationship Trust (HCRT) Scale Willingness to seek help
Grung et al., 2022	Cross-sectional, Survey	250	Self-identification as depressed	.	Norway	GP	Experience of care

Af Winklerfelt Hammarberg et al., 2022	Cluster randomized controlled trial	376	Diagnosis by GP	.	Sweden	PCC	Self-efficacy Preferences
Taylor et al., 2022	Cross-sectional, Interview	14	PHQ-2	.	UK	GP	Self-help strategies Confidence in own ability
Harrison et al., 2023	Feasibility study	18	PHQ-9	.	UK	GP	Adherence to treatment
Jeffery et al., 2023	Cohort study	73808	Chart diagnosis	Patients with type 2 diabetes	UK	PCP	Discontinuation of antidepressants

4

Appendix 3: List of abbreviations

Abbreviation	Full phrase
BDI	Beck Depression Inventory
CBDI	Chinese translation of the Beck Depression Inventory
CB-PHQ-9	Chinese Bilingual version of the 9-item Patient Health Questionnaire
CBT	Cognitive Behavioral Therapy
CES-D	Center for Epidemiologic Studies – Depression Scale
CIDI	Composite International Diagnostic Interview
CIDI-SF	Composite International Diagnostic Interview – Short Form
CPG	Clinical practice guidelines
DIS	Diagnostic Interview Schedule
DSI-SS	Depressive Symptom Inventory – Suicidality Subscale
DSM	Diagnostic and Statistical Manual of Mental Disorders
EPDS	Edinburgh Postnatal Depression Scale
FD	Family doctor
FP	Family practitioner
GDS	Geriatric Depression Scale
GHQ-12	12-item General Health Questionnaire
GP	General practitioner
GRC	Global Rating of Change
HADS	Hospital Anxiety and Depression Scale
HAMD	Hamilton Rating Scale for Depression
HDRS	Hamilton Depression Rating Scale
HRSD	Hamilton Rating Scale for Depression
HSCL-90	Hopkins Symptom Checklist
ICD	International Statistical Classification of Diseases and Related Health Problems
IDD	Inventory to Diagnose Depression
MINI	Mini-International Neuropsychiatric Interview
N/A	Not applicable
OECD	Organisation for Economic Co-operation and Development
PC	Primary care
PCP	Primary care professionals
PHQ	Patient Health Questionnaire
RCT	Randomized controlled trial
RQ	Research question
SCID	Structured Clinical Interview for DSM-IV
Short-CARE	Short Comprehensive Assessment and Referral Evaluation
WHO	World Health Organization