

Supplementary Table 1: Comparison of FILA levels between OA and control plasma lubricin glycoforms in individuals with BMI<30 and for all individuals

Glycoforms	^a All individuals		^b BMI<30	
	Median (Control / OA)	p-value	Median (Control / OA)	p-value
Gal-3	4.195 / 4.519	0.3037	3.991 / 4.167	0.7441
MAA	1.007 / 1.010	0.9360	1.071 / 0.8970	0.2007
PNA	0.2998 / 0.3345	0.9912	0.2891 / 0.3179	0.6017
SNA	1.191 / 1.053	0.0610	1.198 / 0.8950	0.0023**
MGL	0.00160 / 0.00158	0.4436	0.001681 / 0.001604	0.5524
Gal-3/MGL	2820 / 2567	0.6174	2744 / 2488	0.4276
MAA/MGL	610.0 / 635.0	0.7379	599.2 / 615.2	0.3703
PNA/MGL	240.4 / 217.1	0.3009	229.3 / 189.0	0.2007
SNA/MGL	793.2 / 618.6	0.0931	756.0 / 517.2	0.0178*

^a (total n=224), (Controls: n=41), (OA: n=183)

^b (total n=146), (Controls: n=38), (OA n=108)

Supplementary Table 2: Correlation between glycoforms and BMI

Correlation between BMI and lectins in control and OA samples						
		Gal-3	MAA	PNA	SNA	MGL
OA (n=183)	Spearman r	0.1794	0.2453	0.1373	0.2449	-0.00615
	p-value	0.0151 *	0.0008 ***	0.0638	0.0008 ***	0.9342
Controls (n=41)	Spearman r	0.1235	0.04906	0.09891	0.1522	-0.06903
	p-value	0.4418	0.7607	0.5384	0.3420	0.6681
All (n=224)	Spearman r	0.1653	0.2039	0.1255	0.1963	-0.01051
	p-value	0.0133 *	0.0022 **	0.0607	0.0032 **	0.8757

Supplementary Table 3: Correlation between glycoforms and age

Correlation between Age and lectins in control and OA samples						
		Gal-3	MAA	PNA	SNA	MGL
OA (n = 183)	Spearman r	-0.1084	-0,1500	-0,02542	-0,1470	0,1769
	P-value	0.1441	0,0426*	0,7327	0,0471*	0,0166*
Controls (n = 41)	Spearman r	0.2459	-0.05855	0.2807	0.01311	0.06493
	P-value	0.1212	0.7161	0.0755	0.9352	0.6867
All (n = 224)	Spearman r	-0.03545	-0.01338	0.02981	-0.0149	0.1683
	P-value	0.5976	0.0455*	0.6572	0.0258*	0.0117*