

Supplemental Material

Disturbed sleep after lung transplantation is associated with worse patient-reported outcomes and chronic lung allograft dysfunction

Aric A. Prather, PhD, Ying Gao MS, Legna Betancourt BS, Rose C Kordahl BS, Anya Sriram BS, Chiung-Yu Huang PhD, Steven R Hays MD, Jasleen Kukreja MD, Daniel R. Calabrese MD, Aida Venado MD, Bhavya Kapse PhD, John R Greenland MD PhD, Jonathan P Singer, MD MS

Appendix Table 1.

	Continuous scale SPI MID = 8 Insomnia MID = 12		Worst Quartile	
	Sleep Problem Index (SPI)			
Predictor	Parameter estimate (95% CI)	p-value	Odds ratio (95% CI)	p-value
Severe PGD	3.80 (-2.66, 10.27)	0.25	0.97 (0.39, 2.45)	0.95
Length of hospital stay >30 days	12.18 (3.38, 20.99)	0.01	7.09 (2.10, 23.95)	<0.01
Delirium	-1.14 (-10.59, 8.30)	0.81	2.40 (0.69, 8.38)	0.17
	Insomnia			
Severe PGD	1.51 (-7.63, 10.64)	0.74	0.72 (0.27, 1.92)	0.51
Length of hospital stay >30 days	14.56 (2.08, 27.04)	0.02	1.69 (0.47, 6.04)	0.42
Delirium	-0.27 (-13.28, 12.74)	0.97	0.67 (0.13, 3.40)	0.63

Appendix Table 2.

		Time from survey to CLAD (Hazards Ratio)	Time from survey to Death (Hazards Ratio)
Sleep Problem Index (SPI) MID = 8	Per MID worsening	1.09 (0.93, 1.27) p = 0.29	1.27 (1.02, 1.57) p = 0.03
	Worst quartile	1.76 (0.90, 3.45) p = 0.10	2.06 (0.82, 5.18) p = 0.12
Insomnia MID = 12	Per MID worsening	1.15 (0.98, 1.36) p = 0.09	1.10 (0.88, 1.38) p = 0.41
	Worst quartile	1.73 (0.87, 3.47) p = 0.12	0.64 (0.21, 1.96) p = 0.43